crystals of silver nitrate gradually, shaking well after each addition; when the silver nitrate is dissolved, add the mucilage and shake thoroughly once more.

The resulting mixture will represent approximately 3 per cent. of silver iodid. This strength may be increased or decreased as desired. If prepared in this way, the weaker mixtures will require a correspondingly smaller amount of mucilage to hold the powder in suspension and mixtures below 1 per cent. may be dispensed without the addition of any foreign material.

Of the drugs that are frequently used for internal administration in the second or sub-acute stage of the disease the Pharmacopeia includes, in the order of their present popularity:

OLEUM SANTALI.—U. S.—Oil of santal, or oil of sandalwood, is a volatile oil distilled from the wood of *Santalum album*. The Pharmacopeia directs that this oil contain not less than 90 per cent. of santalol. This requirement is readily met by the better grades of oil of santal now on the market, so that there is little or no need to resort to proprietary specialties containing, or purporting to contain, true oil of santal.

Average dose: 0.5 c.c. (8 minims).

COPAIBA.—U. S.—Copaiba, balsam of copaiba, as it is more popularly called, is derived from one or more South American species of copaiba. It was introduced into Europe by the Portuguese, during the latter half of the seventeenth century, and has long enjoyed the reputation of being of especial use in connection with specific urethritis in the male.

Average dose.—1 c.c. (15 minims).

OLEUM COPAIBÆ.—U. S.—Oil of copaiba is officially described as a volatile oil distilled from copaiba.

Average dose: 0.5 c.c. (8 minims).

OLEORESINÆ CUBEÆ.—U. S.—Oleoresin or cubeb represents the alcohol soluble portion of the dried, unripe, fruit of *Piper Cubeba*.

Average dose: 0.5 c.c. (7½ grains).

OLEUM CUBEÆ.—U. S.—Oil of cubeb is a volatile oil distilled from cubeb.

Average dose: 0.5 c.c. (8 minims).

Of the more active astringents that are used in the third, or chronic stage of gonorrhea, in the male, and form such a prominent part in the treatment of the more common form of gonorrhea in the female, we may enumerate:

ZINCI SULPHAS.—U. S.—Zinc sulphate.

ZINCI ACETAS.—U. S.—Zinc acetate.

ZINCI CHLORIDUM.—U. S.—Zinc chlorid.

ZINCI PHENOLSULPHONAS.—U. S.—Zinc phenolsulphonate.

ALUMEN.—U. S.—Alum.

ACIDUM TANNICUM.—U. S.—Tannic acid.

ARGENTI NITRAS.—U. S.—Silver nitrate.

These several astringents, with or without the addition of well-known antiseptics, such as boric acid, phenol, thymol, glycerin, and the several combinations or preparations of them, practically constitute all that is necessary in the treatment of the most persistent of the chronic forms of gonorrhea.

Clinical Note

ACETANILID POISONING FROM BROMO-SELTZER.*

ALBERT ENGLER BLACKBURN, M.D.
PHILADELPHIA.

Patient.—I was asked by his mother to see the patient, a man, aged 22 years, recently married. The patient did not consider himself sick and did not desire medical attention. It was with great difficulty that I obtained a distinct and correct history. I do not believe there was the intent to deceive, but his marked hebétude and obtuseness made it especially difficult to obtain a connected history. When the facts were finally obtained from his wife and he was asked as to their correctness, he acknowledged their truthfulness and promised to desist taking the drug.

* Reported before the West Philadelphia Branch of the County Medical Society, March 16, 1906.

History.—Following an acute illness some six months previous he began taking an occasional dose of bromo-seltzer for headache. The relief desired was temporarily obtained, and as the headaches recurred the bromo-seltzer was taken in larger amounts and at shorter periods. The quantity consumed grew larger and larger until the six weeks previous to my seeing him his wife assured me he consumed as much as \$4.50 per week, which is approximately ten or twelve bottles. The patient would sit for hours if undisturbed, complained of being tired, and was unable to perform muscular effort, there was dyspnea on exertion. He was given to attacks of despondency and again to outbursts of passion, in which he threatened violence, but soon collapsed.

Physical Examination.—His appearance was appalling; his face was ashy white, the lips and lobes of the ears blue, the fingers were deeply cyanosed up to the second joint. His mucous membranes were very pale; the pupils reacted to light and distance. His expression was vacant, seldom speaking until spoken to. The tongue was heavily coated with a white fur, red at tip and margins; breath the foulest I ever noticed. Bowels constipated; the pulse varied from 85 when at rest to 130 on the slightest exertion, regular, fair volume, low tension. The left ventricle was enlarged downward and to the left. The first sound of the heart was very weak; a soft systolic mitral murmur was present. The surface was cold and lifeless. Urine contained indican, but no albumin or casts. General tremulousness and loss of motor powers were marked. No blood examination was made.

To find the constituents of bromo-seltzer I wrote the Emerson Drug Company of Baltimore. Inasmuch as *Collier's Weekly* had lately given them a writeup, a member of the firm called. He told me that bromo-seltzer contained to the 100 grains (a heaping teaspoonful):

Acetanilid	gr. ivss
Potassium bromid	gr. x
Caffein	gr. ii
Soda bicarb.	
Citric acid.	

THE JOURNAL A. M. A., Feb. 10, 1906, states its analysis to be, to the 100 parts:

Potassium bromid	10.53 parts
Acetanilid	4.58 parts
Caffein	1.20 parts

Assuming an average dose to be a teaspoonful—not heaping as above stated—to weigh 76 grains, the dose would contain:

Potassium bromid	7 grains
Acetanilid	3 grains
Caffein	8 grains

Since many people take a tablespoonful repeatedly, the quantity of acetanilid varies between 12 and 15 grains, an amount which is exceedingly dangerous. For "business reasons" the representative requested that I refrain from quoting him.

Course.—Herrick states in his paper that great difficulty is experienced in withdrawing the drug. This was not the case in this patient. He was plainly told the cause of his trouble, and the effect if he continued taking it. He was carefully watched, but made no effort to secure the drug. At no time did he express a desire for it, which was probably due to the relatively short time of taking it. His improvement was steady, but slow. The cyanosis disappeared in about a week; the anemia was the last to disappear.

The treatment consisted merely of withdrawing the drug, administering the cardiac stimulants, strychnia and digitalis, and regulating the bowels.

Dr. Quigley reports in THE JOURNAL A. M. A., Vol. XLVI, No. 6, a case that went into collapse from a tablespoonful of bromo-seltzer, and states he would have died had not his stomach been emptied and stimulants administered.

First Total Gastrectomy in South America.—The *Semana Medica* for April 19 relates the successful removal of a cancerous stomach by B. Vasallo of Rosario, Argentina. The patient was a woman of 39 and was apparently in perfect health at date of writing, four months after the operation, with no digestive disturbances.