that these characters suggest the independence of the phenomena on the nervous system, and their dependence on the mere living undifferentiated protoplasm of the walls of the capillary vessels. But I am not bigoted in the matter of explanations of phenomena, and care more for the accuracy of observations and for the determination of facts.

I want further opportunities for examining the various etiological factors which must concur and the different conditional causes which must be absent for the development of this phenomenon. I make no doubt in my own mind from general grounds of causation, that one chief agent or agency must be at work in the production of the phenomenon, but, nevertheless, all that we yet know of etiology, whether of disease or of other earthly processes, points conclusively to the belief that many other conditions besides the chief one—the exciting cause as we call it—must be operative, and some conditions not operative, in the causation of the process. I have a few other facts of special connexion with the present phenomenon. Frequently, but not always, it will be found that exposure of the skin of the arm and forearm in the same subject presents a contrast with the previously or simultaneously exposed surface of the abdomen, the general tint being white with irregular patches, sometimes streaks, of pink or darker red. When this contrast is evident, and occasionally when there is no such general and spontaneous dissimilarity of appearance, mechanical, and sometimes (? always) electrical and thermic, irritation of the skin of the arm will be followed after a similar, but perhaps generally a trifle shorter, period of latency by the gradual development of the *tache cérébrale* or red streak, having the usual characters of that phenomenon. Sometimes the surface of the arm has a general pink tint on first exposing it to the atmosphere, and this may rapidly change to white, but the colour of the written characters thereon may be deep red. The contrast in colour may be still more remarkable, even in the same case; the writing may be white on the surface of the abdomen, red on the forearm, white on the legs. I do lay considerable stress on the circumstance that the temperature of the skin and of the body is usually high in cases which present these vascular phenomena, and, further, I have most generally observed them in full perfection when the skin has not been perceptibly sweating. But, as proving that neither increased heat nor an especial dryness of the cutaneous tissues is necessary to the production of such phenomena, I may observe that I have seen them very well marked in cases of epilepsy and chorea of uncomplicated and afebrile sort. It may be worth placing on record also in this connexion that the majority of cases in which these observations were made the brain was certainly perturbed in its functions, for delirium was frequently present at some period of the twenty-four hours, mostly at night-time; still I must add that my observations were usually made during the day-time, and when the intellect of the patient, to all ordinary modes of investigation, did not seem to be abnormally affected. I should like to make more investigations at different hours of the day in order to ascertain whether there are any daily or other periodic variations in the phenomena, and also to determine whether the morbid signs present any concomitant variations with other corporeal phenomena, either normal or abnormal. I do not think that a careful clinical study of this nature would be either a waste of time or would not be of some clinical avail.

Harley-street. W.

GLASGOW WESTERN INFIRMARY. — At the annual meeting of this institution, held a few days ago, the report submitted showed that 14,661 out-door patients had been treated during the year, and 3928 in-door. Of these latter 277 died, giving a percentage mortality of 7·7; but of the fatal cases 44 were hopeless when admitted, reducing the death-rate to 6·6 per cent. The average residence of each patient was 37·2 days. The cost of each bed fully occupied during the year was £49 0s. 3½d.; the average cost per patient was £4 19s. 10d. Financially the infirmary was reported to be in a flourishing condition, a balance of £9000 being carried forward to next year.

THE Blair Convalescent Hospital, which has been erected as a result of the munificence of the late Stephen Blair, formerly M.P. for Bolton, at Bromley Cross, near Bolton, is now practically completed, and will shortly be opened for the free admission of convalescents "without limit of domicile."

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

MULTIPLE MINUTE ULCERS OF THE CORNEÆ FOLLOWING EXPOSURE TO ELECTRIC LIGHT.

BY T. PHILLIPS,

CLINICAL ASSISTANT, MOORFIELDS EYE HOSPITAL.

IN this case the causal relations between the exposure to the electric light and the lesions of the corneæ seem to be fairly complete.

On August 31st an electrician, aged twenty-four years, while engaged in making observations of a very powerful electric arc, removed the usual protection in order to make a "better examination," thus exposing his eyes for about four minutes to the action of the light. He suffered no immediate inconvenience; resumed the shade, continued his labours for a short time, went home, had a "good sleep," and had no cause for complaint until he awoke about four hours after the exposure. He then suffered from acute pain in and about the eyes, great intolerance of light, and was unable to sleep. He said, "I was stone blind."

I saw the patient on September 5th. He was still suffering from acute pain and intense photophobia, which prevented examination till the application of a 10 per cent. solution of cocaine. The anterior surfaces of both corneæ were affected. The right was dotted all over with small definite ulcers. The left had six such ulcers arranged in pairs of three rows disposed vertically. The appearance of these ulcers was similar to that presented by the corneal ulceration which occurs in states of perverted innervation of the eye. The ophthalmoscope revealed nothing abnormal.

The eyes were bathed with boric acid lotion, and a solution of cocaine and atropine applied night and morning. In the meantime both eyes were closed with pads smeared with boric acid ointment.

Professor Tweedy kindly saw the case a few days afterwards, and verified the above description and approved the treatment. He also said that he had seen several cases of conjunctivitis with photophobia following exposure to a naked electric light, but had not noticed the corneal ulceration before.

The patient was well in about a fortnight.

A very interesting question suggested by the case is the possible cause of the phenomena—whether (a) due to direct damage of the corneal structure, (b) to direct damage to the trophic nerves, or (c) to indirect damage to the trophic nerves reflexly induced through the optic nerve. It should perhaps be stated that there was no past history of eye disease, except a trifling injury, and the vision was previously normal.

ABSENCE OF THE OCCIPITAL BONE IN NEWLY BORN INFANTS.

BY W. C. BROWN, M.D.

The publication of an instance of this rare deformity in THE LANCET of June 11th, has induced me to give particulars of two cases that have recently occurred in my practice.

On April 26th, 1886, I was called to attend Mrs. N—, aged twenty-four, in her first confinement. She complained of having been suffering from extremely strong foetal movements for a day or two. On examination I found the os dilated, the head well down, and the face presenting. Anticipating a tedious confinement, I was surprised at the ease and rapidity of delivery. On the birth of the infant the reason of this was made apparent; the back of the cranium had remained completely undeveloped. A large soft tumour containing the cerebellum and part of the cerebrum protruded, and on running the finger round its edge the parietal bones could be felt soft and gelatinous. The arches of several of the cervical vertebræ were also wanting, the tumour extending down the back of the neck for some distance. Unlike Dr. Fraser's case, the skin was quite intact over the tumour, and a considerable quantity of hair was growing on it. The child, a healthy

and well-nourished female of full term, drew a few respirations after its birth, but did not live for five minutes. The mother made a good recovery, and I saw nothing more of her until two months ago, when she came to tell me that she was again pregnant, and that she expected to be confined about the beginning of September. I was called to attend her, however, on July 7th, and found her well advanced in labour, with a foot presenting. There was a very large quantity of liquor amnii, but delivery was rapid and easy. On examining the foetus, an eight-months' still-born female, exactly the same deformity was found to exist. The tumour was, however, not so large, and the arches of the cervical vertebrae were in this instance complete. In every other respect the deformity was identical with that of the first child. Notwithstanding the fact that the foetus had not respired, pressure on the tumour caused convulsions, and, if maintained for some time, severe tetanic spasms. In each instance there was a peculiar protrusion of the eyes, which, I observe, is noted also by Dr. Fraser, and the neck was extremely short. In each instance, also, the skin over the tumour was so loose that the finger could easily be pushed between the cerebellum and the medulla, and also for some distance down the spinal canal. The occipital bone—in fact, the entire bony arch behind the foramen magnum—did not exist even in the most rudimentary condition, and the skin was lying directly on the membranes of the brain.

This arrest of development of the posterior arch of the skull is evidently caused by the primitive vertebral discs at the upper extremity of the foetal chorda dorsalis being too few. If the number be even one too few, it will easily be seen how a gap may be left. The double curvature forward of the cerebral extremity, which takes place about the seventh week, would tend to pull what vertebral discs there are forward to the anterior part of the skull. In this way a deficiency in number of the vertebral discs results in a gap being left at the back part of the skull, and the posterior cerebral vesicles are left with no vertebral disc to mould itself over them into a bony covering. From the strong foetal movements that had been felt in both labours I concluded that intra-uterine convulsions had supervened at a certain stage of development, and that this had terminated the pregnancy. The parents are both young and healthy Europeans, the father being of German, the mother of English birth. I have been able to assign no cause for both of their offspring being born with this rare and peculiar malformation.

Penang.

THE TREATMENT OF COMA IN YOUNG CHILDREN BY HYPODERMIC INJECTION.

By F. R. HUMPHREYS, L.R.C.P. LOND., &c.

THE following cases may be of interest, as they seem to indicate a line of treatment applicable even *in extremis*.

CASE 1.—An infant aged ten months, very ill for fourteen days with broncho-pneumonia, complicated by relapses. When first seen (3.40 A.M.) the child was deeply unconscious, unable to swallow, the lips blue, and dying; it had been in this condition for two hours previous to my being sent for. The condition had come on very gradually.—3.40 A.M.: Pulse 176; respiration 96. Three minims of brandy were injected hypodermically.—3.45 A.M.: The brandy injection was repeated. The mouth was noticed to be drawn down to the right.—4.5 A.M.: Pulse 168; respiration 92. Better; coughed a little; nystagmus. Hypodermic injection of $\frac{1}{1200}$ th of a grain of digitalin. The child opened its eyes, looked about, and was able to swallow a little.—4.30 A.M.: Pulse 168; respiration 86. The child had improved so much that I left. It died at 9 A.M. I was told it gradually relapsed into its former state. I was only called in in the absence of the doctor in attendance. On making a hurried physical examination, the right lung appeared nearly solid, and the left was badly affected.

CASE 2.—A boy aged nine months, suffering from measles (fifth day). The rash had gone a few hours before. Hyperpyrexia with consolidation of the left lung having set in, the doctor who was in attendance declared the case hopeless, and said he would not come again. I was therefore asked to see what I could do. When first seen the child had just had a convulsion, but I was told it had been unconscious for some hours. The lips were blue, the pupils extremely contracted, and I could get no reaction to any stimulus.—3.30 P.M.: Respiration 12; temperature 104.8°. It was

wrapped in a wet sheet, an enema of brandy, egg, and milk administered, and a hypodermic injection of two minims of brandy given.—4.50 P.M.: Respiration 22; temperature 102.7°; pulse 140. The child was breathing better, and appeared improving. During this period six minims of brandy and $\frac{1}{1200}$ th of a grain of digitalin were separately injected under the skin.—7.30 P.M.: Temperature normal; much better; able to swallow. The consolidation cleared up; but meningitis set in, and the child died from exhaustion on the seventh day after I first saw him. Each time he flagged brandy or digitalin was administered hypodermically, and each time there was temporary improvement. The child was unhealthy, living under bad hygienic circumstances, in great part nursed by a mother who was absolutely ignorant of such work, and it was only after repeated relapses that it finally succumbed.

In an extremely similar case to the last, broncho-pneumonia passing on to meningitis, with hyperpyrexia (107°), occurred in an elderly but healthy man, and death ensued in a few hours. Hypodermic injection was not made use of, and sponging was of no avail.

Queen's-crescent, N.W.

COMPOUND DISLOCATION OF THE PROXIMAL END OF THE MIDDLE METACARPAL BONE FROM THE OS MAGNUM.

By ERNEST CUSSE, M.R.C.S., &c.

THIS case being an instance of a rare injury may be worthy of note.

J. E—, a gardener, fell from a tree some twelve feet on to the palm of his left hand. On presenting himself at the Bournemouth Dispensary, I found the hand flexed, and the head of the middle metacarpal bone exposed in the palm. The patient complained of very great pain. Having thoroughly cleansed the parts with an antiseptic solution, counter-extension was made from the wrist and extension from the middle finger, pressure at the same time being made on the shaft of the metacarpal bone. The dislocation was easily reduced. The wound was not closed. An antiseptic dressing was used, and a splint applied from the elbow-joint to the ends of the fingers. The case did very well, without suppuration or impairment of the functions of the hand.

Bournemouth.

A VERY RARE PRESENTATION IN MIDWIFERY.

By J. R. BUCHANAN, L.F.P.S. GLAS.

THE very rare presentation in midwifery narrated by Mr. Borham in THE LANCET of Oct. 15th is certainly an unusual one, so far as my experience and reading have extended, but it is by no means unique. On Aug. 20th last I met with a case almost identical with that of Mr. Borham, but with a different result. The messenger who summoned me to assist the midwife, said that a loop of the cord had come down, but that the midwife had been able to replace it. On making an examination, I found a loop of the funis protruding from the vulva. It was quite cold and pulseless. A further investigation showed that the head and left hand were presenting, the hand lying closely pressed against the temple and cheek. I was able quickly and easily—the pelvis being roomy—to push the hand above the brim, and, applying the forceps, delivered in a few minutes, but the child had evidently been dead for some little time. The answer I got, on asking the nurse what she meant by saying that she had replaced the cord, was that she pushed it into the vagina under the impression that that made matters all right.

Glasgow.

DWELLINGS OF THE PEOPLE.—At a meeting of the members of the Mansion House Council on the Dwellings of the People on the 30th ult., it was reported that the inquiry ordered by the Home Secretary into the sanitary condition of Bethnal-green had been concluded, and that the Commissioners' report might shortly be expected. Meantime a very large proportion of the nuisances complained of had been remedied by the vestry. In Mile-end and in Limehouse district a similar condition of sanitary progress was reported.