Otherwise the sight was good, patient being a Rifle Volunteer, and able to shoot at one thousand yards' distance. Health good. No history of syphilis.

On examination of the eyes by the so-called oblique illumination, fine dark lines were seen radiating from about the centre of the pupillary area in the lens of each eye. These lines appeared to be very superficial, and on the surface of the lens, immediately behind the level of the iris. On putting some of the ordinary solution of the sulphate of atropine within the eyes, the pupils dilated fully and regularly, and then the lines could be distinctly traced extending outwards to the circumference of the lens as far as the margin of the iris would allow of them being followed.

An examination of the accompanying figures will show that, although these lines are connected together, they do

Right Eye.

Left Eye.

These lines must be regarded as simply diagrammatic, for as seen in the living subject they do not present the white borders shown in the figures, but appear as black lines on a lighter ground, and this the engraver has been unable to represent.]

not all start from the exact centre of the anterior surface of the lens, nor do they all come from the same point; and that they are more or less irregularly curved in their outward course. Similar lines were seen in several other patients whom I then examined.

This case was shown to several gentlemen, but, at this time, all failed to recognise them; the reason of which was, I now believe, that they did not look in the right direction, nor concentrate the light properly. There is some difficulty in bringing the lines into view; but when this is overcome they are quite distinct. To see them, I use a bi-convex lens of about 2½-in. focus, and throw the light of an argand gas-burner obliquely on the eye of the patient, concentrating the light on the lens. I then look very obliquely (at about an angle of 130° to 140°) on the plane of the anterior surface of the lens, from the side opposite to that on which the light falls. I have never succeeded in seeing them from the front, nor can they be seen with the ophthalmoscopic mirror by transmitted light.

On November 20th this patient was taken to Moorfields Ophthalmic Hospital, where he was seen by seven or eight gentlemen, all of whom, with one exception, said they saw the lines when they took the precaution to examine in the manner I have just described. Mr. Streetfeild suggested that, in Mr. Bowman's work on "The Parts concerned in the Operations on the Eye," 1849, p. 68, there was a drawing of lines in the lens, the plan of which was very much like those I have figured in this case. Mr. Bowman says that the lines represented by him are the central planes of the lens. It will be seen, however, that the lines shown by this gentleman, which I presume are from the prepared lens, do not extend so far outwards as those I have described.

Mr. Wharton Jones was also asked to examine this case at University College Hospital, and he distinctly saw the lines, but said that he had never noticed them before. We then examined several other patients and found similar lines in all; and, in a case of commencing senile cataract, they could be seen intersecting the streaks of opaque lens matter, but apparently on a more superficial plane. In a case of advanced senile cataract, however, they could not be distinctly made out.

As far as I have at present been able to ascertain, these lines are not mentioned in any work as an ophthalmoscopic appearance. They are evidently not pathological, but only physiological; for I have now seen them in thirty or forty cases, of various ages, from two years and a half to seventy years, the majority being healthy boys and young men; in fact, I have never met with a case in which they were not to be seen, although sometimes they could only be brought into view after some considerable time had been spent in looking for them. In the child and younger persons they were less defined and distinct than in those older. As to their real nature, I do not at present venture to give an opinion. They are not light-coloured and generally straight, as cataractous striæ are, but apparently darker than the body of the lens, and irregularly curved. In the adult they are ten in number, more or less, as in the case figured. I have not, so far, been able to make out satisfactorily whether the number differs in younger and older persons.

Stanhope-street, N.W., Nov. 1871.

CASE OF FRAMBÆSIA, OR YAWS.

By J. P. HUGGINS, M.B., C.M., L.R.C.S.E.,
SURGEON TO THE CONVICTS, TRINIDAD.

THE following very remarkable case I had the good fortune to witness in one of my State hospitals. There are some facts which may prove interesting to the profession. It has been stated that the yaws is a disease which is highly contagious; but during the whole time the patient was under medical treatment none of the other patients were troubled with the disease, although we had many open ulcers.

The patient is an East Indian, a lad of fourteen years; and was admitted on the 11th October, 1870. When he came into hospital he complained of fever, headache, pains in the back and loins, and general malaise. Assuming it was a case of ordinary malarious fever, I ordered him a diaphoretic mixture and a warm bath; but on my next visit, I was astonished to see a number of raised pustules resembling those of small-pox. They came out in different parts of the body, more especially on the elbows and knees; and, like the rash of typhoid fever, not at one time, but successively. They were at first about the size of a pea; then they grew larger; and from the top of them exuded a thick grumous fluid. In two or three days afterwards a fungoid excrescence came out, resembling that of a mulberry. I put the patient under a course of iron, in the form of the arseniate, in quarter-grain doses, increasing to half a grain; dressed the ulcers with carbolic oil (which, I must say, is so useful in this country, thanks to Prof. Lister's antiseptic method); and, after the ulcers cicatrised, gave him ten grains of iodide of potash. He was discharged, perfectly recovered, on the 26th January.

I think the yaws is no more than a form of secondary syphilis, and is not a distinct disease.

Queen's Park, Trinidad, 1871.