CASE
OF
CYNANCHE LARYNGEA,

By J. R. FARRE, M.D. *1812*

Read February 18, 1812.

CASE I.—On Monday evening, the 13th August, 1810,—Essex, aged 38, complained of sore throat. On the 14th at noon, his respiration was affected, and in the evening it was loud and hoarse, and he was long in completing each inspiration. He could not sit still for many moments, but walked about the room in the greatest distress, often struggling for breath. Unable to lie down, he passed the ensuing night in the same restless and agitated state, with occasional delirium. On the 15th, the difficulty of his breathing was in no respect diminished. On the 16th, at four in the afternoon, a medical man observed the following symptoms; respiration performed with extreme difficulty; pallid face, protruding eyes; an anxiety difficult to be expressed; the uvula large, œdematous, and speckled, as if about to ulcerate; tonsils not affected;

pulse laborious and frequent. The treatment had been hitherto limited to the use of cathartics, and the application of blisters to the chest and throat. A momentary relief was now obtained by taking twenty-four ounces of blood from his arm. At seven he was under the operation of a preparation of tartarized antimony and squills, which had excited vomiting, and was then acting on his bowels. The blood which had been drawn exhibited the appearances which are usual under active inflammation. At nine o'clock, when sitting up, he expired.

Morbid appearances 24 hours after death.

The velum palati, pharynx, and a portion of the œsophagus with the larynx, trachea and bronchia were carefully detached. The pharynx and œsophagus being slit down posteriorly, the mucous membrane investing the epiglottis, rima glottidis and the posterior part of the larynx, was found to be œdematous to so great a degree, as to make it evident that suffocation had been produced by stricture of the glottis. The tumefaction ceased at the junction of the larynx and trachea. The internal mucous membrane of the larynx was red and thickened. The traces of inflammation were less apparent in the trachea and bronchia, the membrane being irregularly reddened. No coagulable lymph had been effused on any portion of this membrane; and even its mucus was not

changed. There was an appearance of slight congestion in the lungs ; some adhesions of the right pleura existed, but these were not recent ; the heart was natural, and the liquor pericardii slightly increased. Excepting an enlarged spleen, there was no disease of the abdominal viscera.

In this case the inflammation did not pass the stage of congestion in the mucous membrane of the trachea, and of serous effusion under the mucous membrane surrounding the glottis ; but in the following, the inflammation went into the adhesive stage, and closed the glottis by the effusion of coagulable lymph.

CASE II.—A poor, but industrious and temperate man, about 60 years old, on Sunday the 31st of March 1811, was affected with a painful and difficult deglutition. On the following morning, April 1st, alarmed at his inability to swallow fluid, which, on the attempt being made, returned by his nose, he sent for Mr. Weston, who found the tonsils inflamed, and disposed to ulcerate. A brisk purge was ordered. At ten o'clock, on the same evening, his respiration suddenly became difficult, when Mr. Weston immediately drew from his arm thirty-two ounces of blood, which proved to be very sizzly. In the course of an hour I attended, and observed the character of his disease at this stage. Although the tumefaction of the tonsils was inconsiderable, deglutition was extremely painful, and very difficult:

respiration was performed with convulsive and long continued efforts : his voice was nearly inaudible, and he could only whisper. He answered my inquiry respecting the seat of his suffering, by putting his finger on the superior part of the thyroid cartilage. He felt no pain in the chest. All the muscles of respiration were thrown into violent action, and he lay with his mouth widely opened, pupils dilated, face pale and sunken, skin covered with a clammy sweat, and his pulse at 133, and small. His powers were prostrate, and general bleeding could not be repeated. The anterior part of the throat was covered with leeches, but the disease never paused. At midnight the operation of bronchotomy seemed to be the only resource ; and soon after one o'clock, I decided on its employment.

About two o'clock, Mr. Astley Cooper attended, and as suffocation was instantly impending, the operation was immediately performed, by dividing, laterally, the ligament which connects the thyroid with the cricoid cartilage *. The dyspnœa was much relieved by the operation. He now lay pas-

* The following note, made by Mr. Cooper on the morning after the operation, he obligingly sent me with the drawing.

" I was called by Dr. Farre and Mr. Weston, at half past one A. M. on the 2d April 1811, to a person aged more than 60, who had Angina Pharyngea. He was gasping for breath; every muscle that could be brought into action to assist respiration was employed. He was seized with sore throat yesterday, and at ten o'clock to-night with dyspnœa, which has increased so rapidly

sive, breathing by the natural and artificial apertures, and the inordinate action of the muscles of respiration had ceased. He swallowed some nourishment with a painful effort. In this state he passed the night, and the greater part of the following day. In the afternoon, the respiration by the natural passage entirely ceased, but was continued by the artificial aperture. He was now evidently sinking, and expired at six in the evening.

April 3d, eight o'clock A. M. Dissection by Mr. Astley Cooper.

The right tonsil inflamed and vesicated. The epiglottis swollen, its edges meeting behind, excepting just at the upper part. Pharynx inflamed, somewhat vesicated, covered with coagulable lymph about the epiglottis, but free from inflammation near its termination in the œsophagus. The aperture which had been made between the cartilages, appeared to be about half the size of the glottis.

The mucous membrane of the larynx and trachea was pale.

that he appears to be dying, pulse quick, face and lips pale, pupils dilated.

" I began the operation by making a cut between the thyroid and cricoid cartilage, longitudinally, and then transversely into the membrane which united these cartilages. No vessel of the least importance was divided, and he breathed through the opening, so as to be immediately relieved."

There was some accumulation of mucus in the cells of the lungs, and a slight effusion of serum into their reticular texture. The left pleura partially adhered, and the cavities contained rather more fluid than is natural to them. The abdominal viscera were not examined.

I may be allowed to remark that Cullen's character of his 4th species of cynanche will not embrace the cases which I have above described.

“Cynanche (Pharyngea) cum rubore in imis præsertim faucibus; deglutitione maxime difficili, dolentissima, *respiratione satis commoda*, et febre synocha.”

The anterior part of the pharynx was indeed the seat of the inflammation; but that portion of the membrane being also common to the larynx, it may, in a practical sense, be useful to term the disease *laryngea*, because it proves fatal by constricting or actually closing the glottis, and constitutes precisely that case which, in its ultimate degree, imperiously demands the operation of bronchotomy.