A New Integrating Apparatus

This content has been downloaded from IOPscience. Please scroll down to see the full text.

1894 Proc. Phys. Soc. London 13 227

(<http://iopscience.iop.org/1478-7814/13/1/322>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 128.6.218.72

This content was downloaded on 03/10/2015 at 04:59

Please note that [terms and conditions apply](#).

XXI. *A New Integrating Apparatus.* By A. SHARP, B.Sc.*

(Abstract.)

THE paper describes an improved form of harmonic analyser, giving the amplitude and epoch of each constituent term, the mechanism of which is an inversion of that described in a communication made to the Society on April 13, 1894 (*ante*, p. 89). Numerous drawings accompany the paper, showing the various parts in detail. The mechanism is also shown to be applicable for integragraphs, and by suitable modification may be employed for mechanically integrating differential equations of various forms.

XXII. *Magnetic Shielding by a Hollow Iron Cylinder. Simplest Case.* By JOHN PERRY, F.R.S.*

I CONSIDER going and return electric conductors each at a distance a from the axis in a diametral plane of the hollow cylinder, the current in each being C .

As to the inducing magnetic potential, all over one half of the cylindric surface of radius a the potential is constant, πC , and all over the other half it is constant, being $-\pi C$. At a point P whose distance from the axis is r , the angle made by the axial plane through P and the plane through the conductors being θ , it is evident that V , the inducing potential, is

$$V = 4C \left(\frac{a}{r} \sin \theta + \frac{1}{3} \frac{a^3}{r^3} \sin 3\theta + \frac{1}{5} \frac{a^5}{r^5} \sin 5\theta + \&c. \right),$$

as this becomes $+\pi C$ or $-\pi C$ where $r=a$, and also satisfies $\nabla^2 V = 0$.

Let the coefficient of $\sin n\theta$ in V be denoted by $A_n r^{-n}$; let the corresponding coefficients be $A_1 r^n + A_2 r^{-n}$ for the total (induced plus inducing) potential inside the tube; $A_2 r^n + B_2 r^{-n}$ total in the iron and $B_3 r^{-n}$ in outside space. Let the inside

* Read June 22, 1894.