A Mirror OF HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

WEST LONDON HOSPITAL.

OBSTRUCTION OF THE BOWELS; DIVISION OF BAND;
RECOVERY; REMARKS.

(Under the care of Mr. PAGET.)

CONSTRICTION of some part of the small intestine by the pressure of a band is one of the commonest causes of acute intestinal obstruction. The band is usually of inflammatory origin, though the exact way in which it has arisen is not always apparent, especially in the male. There can be no doubt about the necessity for early operation in these cases, and the percentage of recoveries (41 per cent.—Curtis), although favourable now compared with that for operation for other conditions, will be much improved in the future. The closing of the wound is, we think, best performed by sutures passing through all the layers at once. This method is speedy and gives very good results. There should be no likelihood of including intestine, and if the omentum is carefully drawn down towards the pelvis, as in ovariectomy, subsequent adhesion of the bowel to an imperfectly sutured wound is improbable. For the notes of this case we are indebted to Dr. Causton, in whose practice it occurred, and for those taken in the hospital to Mr. E. W. Lewis.

A young man twenty-five years of age was admitted to the West London Hospital on Jan. 8th, with signs of acute obstruction of the bowels. He had always enjoyed good health, save for an attack of typhoid fever in early childhood. His bowels had always acted regularly, and he had never felt pain in the region of the appendix vermiformis. His illness began on New Year's Day. He was in his usual health, his bowels having acted well in the morning. He had not been exposed to any chill and had not taken any unwholesome food. He was seized with pain, which began at the umbilicus and then became more diffused, and he vomited. He came to the hospital, but was not admitted. From Jan. 1st to 7th he was under the care of Dr. Causton. The abdomen at this time was very tender, but soft; no coils of intestine were visible, and no noises were heard save a slight gurgling on the right side. There was at first no distension, or very little. He vomited once or twice daily. On Jan. 4th an enema brought away some large lumps and some undigested food. This gave him relief, and the vomiting stopped for a time. His temperature was always found to be normal, and he would insist on leaving his bed for an armchair. Mr. Paget saw him first, with Dr. Hood, on the afternoon of the 8th. He had walked into the hospital the day before, not seeming to be very ill, but on this day his condition had taken a great change for the worse. An enema had been given without any result. He had just brought up nearly a pint of thick brown, lumpy, faecal vomit. The face was pinched and haggard, and the eyes were sunken. He was in pain, restless, depressed, and weak. There was frequent hiccough; the tongue was moist and fairly clean, with a thin white coat all over it. The pulse was 100, regular and fairly strong, the temperature was normal, and the breathing quiet. The urine was loaded with urates, but there was no albumen. The abdomen was slightly distended, but not more in one part than in another; there were no dilated veins. A high-pitched note was heard everywhere, but it was not tympanitic. There were no visible coils of intestine, no movements, and no noises; there was also no marked tenderness. The legs were not drawn up. Nothing was to be felt in the right iliac fossa, or by the rectum. It seemed to be certain that the patient was not suffering either from inflammation round the appendix, or from intussusception, or from impacted gallstone, or from volvulus. The diagnosis lay between acute inflammation of the bowels, arresting their action at the inflamed part, and internal strangulation. The following

considerations were in favour of internal strangulation: 1. He had not been exposed to cold or wet, he had taken no unwholesome food, he had kept quiet all Sunday, and had been seized with sudden pain on Monday morning. 2. He had felt no previous uneasiness in his bowels; they had acted properly just before he was attacked with pain. They had never given him any trouble in his life, and he had never had attacks of pain in them. 3. He had been ill a whole week. Acute inflammation of the bowels would surely either have proved fatal within the week or would have abated under treatment; and there would have been large, motionless coils of distended bowel plainly visible, lying inert beneath the abdominal wall. With Dr. Hood's approval Mr. Paget operated at once. Two coils of intestine presented through the wound, distended and inflamed, one of them slightly rough with lymph. Mr. Paget first made sure that the appendix felt healthy; then he took the coil that had the lymph on it and followed it downwards and to the right for eight or ten inches. Here it was fixed and could not be drawn forward, and began to bleed a little, and something could be felt holding it down. The wound was enlarged to five inches; the bowel could then be drawn forward a little, and was found to be constricted by a tough, rounded band about an inch long and about the thickness of a No. 7 catheter, attached at either end to healthy coils of small intestine. It was easily divided, and the constricted bowel at once became distended. The wound was quickly closed, the vomiting ceased, and the patient made a good recovery. On the tenth day after operation an enema was given and acted well. It was repeated on the twelfth and fifteenth days, and on the eighteenth day he had some castor oil. The bowels now act regularly without medicine and the patient is in good health.

Remarks by Mr. PAGET.—The diagnosis in this case was helped more by the negative evidence against inflammation than by any positive evidence of strangulation. The operation was happily free from difficulty. The prolonged inaction of the bowels after strangulation is worthy of note. The origin of the constricting band must remain a matter of conjecture; possibly it was due to the attack of typhoid fever in childhood.

THE ENGLISH HOSPITAL, HUELVA, SPAIN.

EXCISION OF THE LARYNX; THE AFTER-TREATMENT OF
THE PATIENT; REMARKS.

(Under the care of Dr. MACKAY.)

THIS case of tumour of the larynx not only shows the result of a case of extirpation of the larynx, but is also a good example of sarcoma affecting that region. As a rule, sarcoma of the larynx is of slow growth; in this instance it was rapid, and death threatened to occur from suffocation. Mr. Butlin¹ writes of these cases: "Partial or complete removal of the larynx appears to be a suitable one for cases of sarcoma. The more extensive the disease and the more extensive the consequent operation the less satisfactory is the result likely to be in every respect, not only with regard to the probability of recurrence of the disease, but on account of the after-condition of the patients." The results of operations for sarcoma are much better than similar ones for carcinoma. Hahn collected a series of sixty-five cases of operation for carcinoma; of these, twenty-eight died within fourteen days and five within six or seven weeks from pneumonia or broncho-pneumonia. He also gives records of six complete and two partial excisions for sarcomatous tumours. Of the complete operations two died with recurrence of the disease at the end of seven and fifteen months respectively, one was quite well six years after operation, another two years afterwards, and the fifth died from phthisis eighteen months afterwards with evidence of recurrence of the disease. Mr. Butlin has pointed out that in the second half of the series the results were better than in the first half, and we have no doubt that the future published results will be even better. It must, however, be recollected that it is almost impossible to arrive at the exact truth, for so very often the unsuccessful cases are not recorded. When each hospital is obliged to publish a yearly statistical account of the patients under treatment, according to an approved standard, we shall perchance be able to give a more definite and trustworthy opinion on many similarly obscure questions.

The patient, a man aged forty-four years, was admitted to

¹ Operative Surgery of Malignant Disease.