Schreiben des Herrn *Valz*, Directors der Sternwarte, an den Herausgeber.

Marseille 1850. October 19.

J'ai pu suivre ici la dernière comète jusqu'au 13 Ct. et je vous en remets ci-dessous les observations, dont je vous avais déjà envoyé les deux premières, établies sur un moindre nombre. Je n'y ai pas tenu compte de la réfraction, qui a du agir surtout sur les dernières. Le 4 Octobre les erreurs des éphémérides étaient peu sensibles, mais le 13 Ct. leurs déclinaisons se trouvaient trop faibles de 15 min. et les AR. de 4 min. Je crains qu'à la fin de Novembre, où la comète pourra reparaitre, elle ne soit trop faible pour être distinguée

du crépuscule. Voici les seconds élémens que j'ai obtenus, et qu'il faudra encore un peu corriger.

Pass. au Pérh. Octobre 19, 288 T. M. à Marseille

Dist. Pérh. 0,5686

Long. Pérh. 88°56'

Long. Ω 206 27

Incl. 39 19

Mouvement Direct.

Observations de la 2e comète de 1850 faites à Marseille par Mr. *Benjamin Valz*.

	tems moy.	Etoiles comp.	Différ. AR.	Diff. en D.	Asc. droit.	Déclinaisons.
Sept. 13	14 ^h 14 ^m	Lal. — Baily 14014	—1° 1' 33"	+19' 12"	105° 35' 54"	46° 54' 0"
14	14 37	L. B. 13712	+4 35 52	+15 8	109 10 39	44 44 29
—	—	— 14282	+ 39 0	—23 31	109 10 30	44 44 37
Oct. 4	17 0	— 19949	+ 18 55	+ 1 0	152 26 30	— 4 20 15
—	—	— 20043	— 33 44	+ 0 38	152 26 35	— 4 20 40
—	—	— 20045	— 34 22	+ 8 14	152 26 48	— 4 20 29
—	—	— 20076	— 51 45	+17 12	152 26 40	— 4 20 44
—	—	— 20086	— 57 30	+19 35	152 26 38	— 4 20 28
7	16 51	— 20140	+2 28 30	— 4 12	156 17 44	— 8 42 46
—	16 36	— 20446	— 11 0	— 4 27	douteux avec nuages.	
8	16 50	— 20521	+ 18 20	— 9 28	157 31 5	— 9 58 3
—	17 1	— 20753	—2 3 35	— 3 56	157 31 42	— 9 59 40
10	17 1	— 20768	+ 11 32	+25 23	159 56 50	—12 15 15
—	16 57	— 20609	+1 52 49	— 1 14	159 56 24	—12 14 24
11	16 40	— 20978	— 22 0	—25 48	161 22 48	—13 23 47
12	16 50	— 20893	+1 24 30	— 4 51	162 19 13	—14 11 4
—	16 56	— 21046	— 3 45	— 4 28	162 19 9	—14 10 37
13	17 2	— 21150	+ 4 15	— 4 36	163 28 53	—15 4 1
—	17 11	— 21178	— 13 15	—	163 28	

Benj. Valz.

Schreiben des Herrn Professor *Challis* an Herrn *Rümker*, Director der Hamburger Sternwarte.

Cambridge Observatory 1850. October 3.

I have received by different conveyances two copies of your Star-Catalogue, the one having arrived, which you had given me reason to expect, through a London Bookseller. I have consigned one of the Copies to the Library of this Observatory and shall esteem it a favour to be allowed to retain the other in my own possession. I delayed writing to acknowledge the receipt of these, till I could at the same time send my final observations of Parthenope reduced. Under the advantage of the fine evenings we often have here in the month of September, and by the aid of Mr. *Gould's* Ephemeris in the *Astronomical Journal*, I have been able to pursue this planet much longer than I anticipated. It was seen to a later date than Sept. 16, but no observation worthy of

confidence could be taken. May I beg of you the favour, after making what use you please of the enclosed paper of Astronomical communications, to forward it to Professor *Schumacher*?

The printing of the Volume containing the Cambridge Meridian observations of 1844 and 1845 is drawing to a close, and when completed, I shall have the pleasure of forwarding copies through the Royal Society (my usual medium) to Hamburg and to Altona.

Allow me to take this opportunity of congratulating you on the honorary distinction you have lately received from the King of Hannover.

Results deduced from Occultations of Stars by the Moon observed at the Cambridge Observatory in the year 1845.

The following are the observations from which the results were calculated.

Reference number.	Greenwich Mean Time of observation 1845.		Star.	Phenomenon.	Moon's Limb.	Remarks.
1	Febr. 14	10 ^h 53 ^m 10 ^s 17	ω^2 Tauri	Disappearance	Dark	Very exact.
2	Aug. 20	9 23 2,20	δ Piscium	Reappearance	Dark	
3	26	12 46 17,20	χ^4 Orionis	Reappearance	Dark	
4	—	16 12 28,72	ϵ' Orionis	Disappearance	Bright	
5	Sept. 13	8 30 2,49	ϵ' Capricorni	Disappearance	Dark	Mean of two observations differing by 0 ^s 17
6	—	8 31 20,01	B. A. C. 7562	Disappearance	Dark	Star not in Naut. Alm. place taken from B. A. C.
7	14	7 35 55,94	α Aquarii	Disappearance	Dark	Very good.
8	—	8 9 59,58	α Aquarii	Reappearance	Bright	Doubtful.
9	15	13 17 42,13	λ Piscium	Disappearance	Bright	Star excessively faint.
10	—	14 26 21,42	λ Piscium	Reappearance	Dark	Uncertain from clouds.
11	Oct. 23	12 15 51,12	$\mathcal{A}^2$ Cancri	Disappearance	Bright	The Moon low and star faint.
12	—	13 11 8,16	$\mathcal{A}^2$ Cancri	Reappearance	Dark	Very exact.
13	Dec. 6	10 24 54,96	λ Piscium	Disappearance	Dark	Mean of two observations differing by 0 ^s 13.
14	—	11 13 6,02	λ Piscium	Reappearance	Bright	Perhaps late.

Let the Longitude of the Cambridge Observatory be $-23^\circ 54' + \tau^s$, the Latitude $52^\circ 12' 51'' 63$, the angle of the vertex $11' 12'' 00$, the Earth-radius [9,9990916]. Also let the true R. A. and N. P. D. of the Moon and the true R. A. and N. P. D. of the Star, be equal to their assumed values, derived from the Nautical-Almanac increased respectively by

x, y, e, f , in seconds of arc, and the Moon's true Horizontal-Equatoreal-Parallax and Semidiameter be equal to the values of the Nautical-Almanac multiplied respectively by $1 + 0,001m$ and $1 + 0,001n$. Then from these data the following equations result:

Reference number.									
1	$+ 9'' 22 = -0,900 \times x + 0,300 \times y + 0,895 \times e - 0,300 \times f - 0,487 \times \tau + 2,407 \times m - 0,894 \times n$								
2	$-13,84 = +0,860 \times x - 0,505 \times y - 0,859 \times e + 0,502 \times f + 0,554 \times \tau + 0,481 \times m - 0,949 \times n$								
3	$+ 1,05 = +0,891 \times x + 0,318 \times y - 0,892 \times e - 0,320 \times f + 0,475 \times \tau + 2,672 \times m - 0,888 \times n$								
4	$+ 0,25 = -0,715 \times x + 0,663 \times y + 0,712 \times e - 0,654 \times f - 0,351 \times \tau + 0,118 \times m - 0,894 \times n$								
5	$+17,27 = -0,846 \times x + 0,531 \times y + 0,837 \times e - 0,528 \times f - 0,591 \times \tau + 0,904 \times m - 0,984 \times n$								
6	$+12,97 = -0,744 \times x + 0,669 \times y + 0,736 \times e - 0,666 \times f - 0,558 \times \tau + 1,437 \times m - 0,984 \times n$								
7	$+ 6,10 = -0,126 \times x + 0,998 \times y + 0,124 \times e - 0,992 \times f - 0,274 \times \tau + 2,708 \times m - 0,977 \times n$								
8	$- 6,95 = +0,823 \times x + 0,576 \times y - 0,817 \times e - 0,572 \times f + 0,346 \times \tau + 2,889 \times m - 0,978 \times n$								
9	$+ 6,33 = -0,982 \times x + 0,234 \times y + 0,973 \times e - 0,232 \times f - 0,587 \times \tau + 1,410 \times m - 0,972 \times n$								
10	$-26,21 = +0,790 \times x - 0,627 \times y - 0,783 \times e + 0,622 \times f + 0,560 \times \tau - 2,768 \times m - 0,970 \times n$								
11	$- 3,02 = -0,941 \times x + 0,265 \times y + 0,942 \times e - 0,261 \times f - 0,427 \times \tau - 1,271 \times m - 0,891 \times n$								
12	$- 7,67 = +0,758 \times x + 0,635 \times y - 0,755 \times e - 0,633 \times f + 0,455 \times \tau + 3,078 \times m - 0,894 \times n$								
13	$+ 4,01 = -0,956 \times x - 0,312 \times y + 0,951 \times e - 0,310 \times f - 0,443 \times \tau + 0,895 \times m - 0,951 \times n$								
14	$- 5,96 = +0,462 \times x - 0,891 \times y - 0,460 \times e + 0,888 \times f + 0,415 \times \tau - 3,344 \times m - 0,949 \times n$								

(Schluss folgt).

J. Challis.