masses on the growing secondary optic vesicle. The macular coloboma remains inexplicable.

4. *Changes Due to Diphtheria Toxemia.*—Dudgeon contributes a lengthy, well-illustrated and detailed morphological study of the tissue changes in acute diphtheritic toxemia, paying particular attention to the changes thought to be responsible for acute cardiac failure. Sixteen cases served as a basis of the examinations in conjunction with extensive experimental work. The changes found in the tissues coincide with those with which pathologists are already familiar. He concludes, however, (a) That the most important lesion in the acute cases is a fatty change of the heart muscle and diaphragm, which is due to a direct action of the toxins on those tissues, (b) That similar fatty changes may be found in certain of the important viscera, especially the adrenal gland and liver, (c) That the expression "cardiac paralysis" in acute diphtheritic toxemia should be abolished and replaced by "acute cardiac failure," (d) That the changes found in the nervous system are secondary factors and not the primary cause of cardiac failure, (e) That the antitoxin, if given in sufficient quantity and within the first forty-eight hours, may prevent or considerably diminish the possibility of death from cardiac failure.

JELLIFFE.

Miscellany

FRACTURES OF BASE OF SKULL. (Annals of Surg., December, 1906.)—Drs. L. R. G. Crandon and Louis T. Wilson analyze 530 cases of fracture of the base of the skull. The mortality was 44 per cent. Alcoholic intoxication was present in 32 per cent., of whom 38 per cent. died. Many of these developed delirium tremens, which could not be wholly differentiated from symptoms of laceration or meningitis. Forty-four per cent. were unconscious at first examination, and fully half the conscious ones were dazed or stupid. Fifty-eight per cent. of the unconscious died, against 33 per cent. of the conscious. Early delirium was usually due to alcoholism, but when late, infection was the cause. Hemorrhage from at least one cranial orifice was present in 74 per cent., the order of frequency in this series being the ear 281, nose 168, subconjunctival 53, two ears 47, mouth 44. In practically all cases of aural hemorrhage the drum was ruptured. Where pharyngeal bleeding occurs the mortality is very high. Respiration as a symptom is unimportant except in primary shock and toward a fatal termination, and the quality of the pulse is of greater significance than its rate. In 117 cases presenting stertorous breathing 106 died. A subnormal temperature seems to indicate shock or alcoholism. Nearly all those with high initial temperature died. The escape of cerebrospinal fluid was noted in only 27 cases of this series. The most striking result of the study of pupillary change was that in 142 cases without reaction 131 died. Of these 530 cases 59 were operated, with a mortality of 53 per cent. The writers conclude that only those cases should be trephined where either hemorrhage seems the most important part of the clinical picture or the course is considered one of continuous and progressive intracranial compression. Patients with non-reacting pupils, or unconscious with stertorous respiration are nearly always bleeding deep in the brain substance, and are hopeless. It is urged that in suspected basal fracture that rest in bed for three full weeks be enjoined.

COWLES (New York).

CEREBROSPINAL MENINGITIS. By J. S. Billings, Jr. (Journal A. M. A., June 2).

Dr. Billings gives a history of the epidemic of cerebrospinal meningitis in New York City in 1904-05, with the methods employed to meet it. It