THE VAPOUR BATH IN THE TREATMENT OF IDIOPATHIC TETANUS.

To the Editor of THE LANCET.

SIR,—In answer to inquiries of private correspondents as to the construction of the Vapour Bath mentioned in my letter on the "Treatment of Idiopathic Tetanus," which you did me the favour to publish on the 10th ultimo, p. 465, I take the liberty of sending you a sketch and description of the kind of bath which I have found to answer the purpose admirably. It differs from the ordinary vapour bath by including the head, so that the patient is respiring air of the same temperature as surrounds his body; and by the radiated heat from the cover of the reservoir, the upper part of the bath level with the head is from 10° to 20° cooler than it is at a level with the feet. It differs also from the hot-air or Turkish bath in having the atmosphere perfectly saturated with moisture without the bath being filled with visible vapour.

- a, b, c, d. Steam reservoir of wood, lined with copper, tinned, with a top or lid perforated, so as to admit only steam enough to saturate the air in the bath with moisture.
 e, f, g, h. Canopy, formed by four pillars, covered with twilled or swan's-down cotton, and drawn close at the bottom.
 i. Steam service-tube, with two way cocks—one to heat the bath, the other to pass through the herb-pot (k) to medicate the bath.
 l. The chair, cane-bottomed.
 m. Foot-stool of wood, perforated.
 Steam may be supplied from a common kitchen boiler, or by a suitable boiler for the purpose. As the bath is heated partly by radiation from the lid, and partly by steam admitted through the perforations, a considerable force of steam is required.

The advantages of this kind of bath are:

1. That free perspiration is elicited at a temperature not exceeding 100°.
2. That the canopy permits perfect ventilation, whilst it retains sufficient heat to produce the effect.
3. That it admits of agreeable medication with aromatic herbs, without which steam is sickening.
4. That there is no limit to the time during which the bath may be used; by drinking freely whilst in it, to supply the waste by perspiration, the bath may be continued for several hours, or until the spasms be overcome, without occasioning exhaustion—a point of great importance in the treatment of so formidable a disease as tetanus.

Lastly, that the reservoir receives the condensed steam and water—warm, tepid, or cold—poured over the patient to wash off the perspiration before leaving the bath.

I am, Sir, your obedient servant,

Montague-street, Bloomsbury, Dec. 1860. J. PIDDUCK, M.D. Edin.

VACCINATION.

To the Editor of THE LANCET.

SIR,—I observe in your impression of the 10th ult. a letter from Dr. Hogg, in which he mentions a case of vaccination, the pock of which was not fully developed till the *tenth* day after the insertion of the vaccine virus. Since that time I have had a case of vaccination in which the pock was not fully developed till the *twenty-ninth* day after the insertion of the vaccine virus. I have not, during twenty-four years' practice, met with another case where the action of the inserted virus has been retarded for so long a time.

Ipswich, Dec. 1860.

WEBSTER ADAMS, M.R.C.S.

594

NEW METHOD OF REDUCING STRANGULATED HERNIA.

To the Editor of THE LANCET.

SIR,—The subject connected with the reduction of strangulated hernia is of such importance that I trust, as the originator of the present discussion, I may be permitted again to trespass upon your columns.

Apart from all question as to the novelty of the treatment alluded to, I would venture to express my surprise that, with the references lately pointed out to the notice of the profession by your various correspondents, the plan has not been more frequently and legitimately carried into *practice*, and the results made known to the profession.

In further illustration of the proceeding, my friend, Dr. Wilson, of this town, has drawn my attention to the following very curious extract from the lately published Life of Lord Dundonald (vol. i., p. 230):—"Another man was refused admission (to hospital) because everything possible had not been done to reduce a rupture, as he had not been hung up by the heels, in a rolling sea, which might have caused his death."

Dr. Evans, of Gloucester, informed me that many years ago, on entering a cottage, he found a young man with his heels high up against the wall, and with his hands firmly applied to his groin. Upon inquiry, he found that the young man was putting back a rupture, and that it was the only way in which he could effect his purpose with certainty and comfort. Common sense had served him well in his emergency.

On referring to one of our best standard text-books on Surgery, I find the following directions given previously to the application of the taxis:—"The bladder having been emptied, the patient should lie down, with his shoulders raised, and both his thighs should be bent towards the belly, and be placed close to each other, so that every muscle and ligament connected with the abdomen may be relaxed." Further on, I read:—"It sometimes happens that the taxis succeeds better when the abdominal parietes are not so much relaxed; at all events, this plan might be tried if the ordinary one fails."

What the muscles and ligaments of the abdomen, excepting those in immediate connexion with the seat of stricture, can have to do with the reduction of strangulated hernia, I cannot conceive; but I can well understand that the flexed condition of the thighs and the raised position of the shoulders, while relaxing the abdominal parietes, must necessarily tend to huddle and throw the weight and the pressure of the intestines towards the seat of mischief, complicating the difficulties already existing. The reduction of a strangulated hernia under such circumstances I believe to be more the result of a happy accident than of any hitherto well-explained rule or law of treatment. On the other hand, let the pelvis be sufficiently well raised, and the intestines thrown back upon the diaphragm, and I venture to say that, where no serious adhesion has taken place, the hernia may in a large majority of instances be readily reduced by a fair trial of the means referred to in this discussion.

I trust those gentlemen who take an interest in the subject, and who may have opportunities of testing the practice, will, through your columns, favour the profession with the results of their treatment. In this manner, we may be enabled to collate valuable data for future consideration.

I am, Sir, your obedient servant,

WALTER JESSOP, M.R.C.S. Eng.

Royal-crescent, Cheltenham, Dec. 1860.

TREATMENT OF STRANGULATED HERNIA BY GRAVITATION (?)

To the Editor of THE LANCET.

SIR,—Perhaps you will afford space in your standard serial for an additional remark or two on the above topic.

In a "System of Surgery," published upwards of seventy-two years ago by an accomplished practitioner of Edinburgh,* a case of the above nature is detailed, where, along with various usual items of treatment, the plan alluded to by your several correspondents is described as having been had recourse to, and this without intimation of any sort to indicate that such proceeding was a discovery or novelty even at that date.

It seems highly probable that the idea of reducing hernia by *inverting* the patient, and thus aiding the effect of gravitation on the abdominal viscera, has been of *popular* origin ages ago; just as we know that the same *force* was in former times sometimes (most absurdly) made available towards another purpose,

Practical System of Surgery. By Jas. Latta, Surgeon, Edinburgh.