away from the renal origin of eclampsia, and made them attribute the whole phenomenon to some poison of unknown nature, but, according to Edgar, not uremic. Why, then, Dr. Woods asked, should ophthalmologists persist in picking out one symptom of this phenomenon as certainly uremic, especially when the obstetricians had taught that albuminuria could also be a secondary result of toxemia?

THE RELATION OF OCULAR AND CARDIO-VASCULAR DISEASE.*

MELVILLE BLACK, M.D.

Professor of Ophthalmology, Denver and Gross School of Medicine.
DENVER.

This subject, I am sure, is not new to you, but since we are learning more about it every day, a few of my personal ideas and findings may prove of interest. The estimation of blood pressure has brought about a more comprehensive knowledge of cardiovascular disease. The remote disturbances which follow this important condition, and the etiologic factors which gradually lead up to it, are now receiving much attention. It is a well-recognized fact that a man may be on "his last legs," so to speak, from cardiovascular changes, and yet feel no serious bodily discomfort. He may have, therefore, no urgent reason for consulting his physician, but may have reason to consult his oculist, and we should constantly be on the lookout for ocular evidences of these vascular changes.

The recognition of these early changes in the retinal circulation may enable us to give many years of life to a considerable number of our patients. The "optometrist" will always be incompetent to recognize these danger signals in the eye, and it is one more urgent reason why he should not be licensed to practice his would-be profession. While the ophthalmologist keeps pace with the body of the medical profession in its new discoveries, the technical knowledge and the equipment required to make a thorough examination of the urine, the blood, and an all-round physical examination of the individual goes beyond his province. He has served a most useful purpose, however, in having recognized the ocular signs which warrant such a physical examination, and has done his duty by seeing that the patient had such an examination made.

It is no disparagement to the general practitioner to say that he also is incompetent to make this examination. It has been my policy to refer such a patient to some one capable of making this all-round examination, and then send him to his physician with the full report of our findings. I must confess that my efforts have been several times thwarted by the family physician making light of the entire matter. He thumps the patient a little, listens to his heart through a few intervening thicknesses of clothing and directs him to "go about his business." A dilated heart can only be mapped out by the most painstaking deep percussion. The presence of a low blood pressure, polycythemia, and dilated, tortuous retinal veins, uricacidemia and indicanuria all point to a clear diagnosis of general venous stasis, which tends not only to shorten life, but also opens the way to many physical mishaps before the end comes. The cases I am about to report show how these mishaps may affect the eyes.

I believe it is generally accepted that high arterial

pressure is a common cause of renal and arterial degeneration. It has been my observation that in chronic interstitial nephritis, with high blood pressure, where the retina is involved, the reduction of the blood pressure is all important, and that thereby several years can be added to the life of the individual (Cases 11, 12 and 13). Case 11 shows not only how life was prolonged when the patient was on the verge of uremic convulsions, but how useful vision was preserved until her death.

I have been especially impressed with the frequency with which general venous stasis was found present in many cases of degenerative ocular changes (Cases 1, 2, 3, 4, 5, 6, 7 and 9). Several of these patients I never saw again; consequently I am unable to report on the results of treatment, but they serve to show the relationship, as mentioned. If the vascular changes were not instrumental in causation, then the etiology was certainly indeterminable. At our altitude of one mile above sea level we undoubtedly see more cases of heart dilatation and venous stasis than do those living at lower levels. How much greater the relative proportion is I do not know.

It is now a recognized fact that four-fifths of the heart lesions are muscular and that one-fifth are valvular. The influence which impaired circulation, caused from a weakened heart, has on the economy is now being studied extensively. The ophthalmologist is in a position to see some of the very direct results of muscular heart lesions; he should, therefore, be thoroughly alive to the importance of this subject.

CASE 1.—Retinal Hemorrhage, Due to Dilated Heart, General Venous Stasis and Polycythemia.—Mrs. J. G. M., age 37, Aug. 8, 1907. Has been feeling well physically, with the exception of some shortness of breath on exertion. Two months ago she noticed a blurring of vision in the left eye, which has undergone no improvement.

Status Præsens.—The retina of O. S. is highly engorged. The veins are tortuous and greatly distended. Flame-like hemorrhages are present everywhere along the veins. The nerve head is red, outlines indistinct and somewhat swollen above.

The appearance of the fundus of O. D. is practically normal. The veins are somewhat larger than they should be. O. D. V. = 20/20; O. S. V. = 2/200. She was referred to Dr. E. C. Hill for physical examination, who reports as follows: Pulse, 92, sitting; 110, standing. Heart, one-half inch to left of nipple and one-half inch to right of sternum. Blood pressure, 105; 107 after exertion. Hemoglobin, 95 per cent. Red blood cells, 7,800,000. Urine, 1,027 c.c. in twenty-four hours. With the exception of a slight excess of indican and uric acid it is normal.

Treatment.—Fifteen drops tincture of strophanthus before meals, and 1/30 gr. strychnia after meals. Diet: Not more than two and one-half pints of fluid daily. Meat once daily. No coarse vegetables, such as cabbage, corn, etc. Potatoes and bread restricted. Fruit and green vegetables freely. Generally speaking, her diet is to be very simple and she must eat sparingly. Results two months later: O. S. V. = 20/30. Retinal hemorrhages all absorbed. Veins more normal in size and appearance; nerve head a little muddy. Dr. Hill finds her heart normal in size. Blood pressure, 108. This lady lives in Ohio and was seen by me while on a visit to Denver, consequently our altitude had nothing to do with the production of her venous stasis.

CASE 2.—Optic Neuritis, Associated with Exophthalmic Goiter, Dilated Heart, General Venous Stasis and Autoinfection.—Mrs. C. H., age 31, November, 1906. History of enlargement of the thyroid gland with some exophthalmic symptoms. The enlargement has about disappeared under electropuncture. Eyes have been paining her for two years. Has had a number of transient blind spells. Vision became per-

* Read in the Section on Ophthalmology of the American Medical Association, at the Fifty-ninth Annual Session, held at Chicago, June, 1908.

manently affected eight months ago. It has gradually been growing worse. Has not been able to read for three months.

Status Præsens.—O. D. V.=fingers at four feet. O. S. V.=fingers at six inches. Left pupil larger than right, reactions for light and accommodation slow. Fields: Not enough vision to take left field; the right is contracted equally in all directions to the 5 degree line. With the ophthalmoscope, the nerve head of O. D. is swollen 5 D. No retinitis. The left nerve is almost completely atrophic. She was sent to Dr. Hill for physical examination, who reports as follows: Heart dilated to one and one-eighth inches beyond sternum and two inches beyond nipple. Blood pressure, 129. Pulse, 108. Urine, great excess of uric acid and indican. Red blood cells, 4,480,000. Patient disappeared.

CASE 3.—Retinal Hemorrhage, Associated with Polycythemia and Autointoxication.—Mr. B. D. S., age 44, Aug. 13, 1903. Very full-blooded, florid-looking man, who is apparently somewhat over weight. Claims to have always enjoyed good health. Is a heavy eater, but has no other dissipations. About six weeks ago noticed that vision of O. D. was poor; no improvement since.

Status Præsens.—O. D. V.=3/40. O. S. V.=5/6. The entire macular region of the right eye presents the appearance of numerous hemorrhages, with chorioiditis. The veins are very large and tortuous, and the nerve head is red. Sent him to Dr. E. C. Hill for physical examination, who reports that his heart is about normal, but that his red blood cell count is 8,200,000; that he is passing only about one-half the normal amount of urine in twenty-four hours, which contains a considerable excess of indican, and three times the normal amount of earthy phosphates. He advises that he drink twice as much water as he is now drinking, and that his diet be of the simplest form; that it be largely vegetable and that he be allowed only one full meal a day. He was given 10 gr. iodid of potassium, t. i. d., 1/32 gr. biniodid of mercury, t. i. d. In three months time his vision had improved to 5/15. The fundus shows a few scars. No new hemorrhages. I saw him last in February, 1906. He had O. D. V., 20/30. Almost no retinal signs of his former trouble. I had Dr. James Rae Arneill examine him physically for me. He found his blood pressure to be 135; hemoglobin, 105 per cent.; urine normal. A letter received from him a few days ago says the eye still remains well.

CASE 4.—Retinal Hemorrhage, Associated with a Weak Heart and General Venous Stasis.—Dr. J. M., age 53, Aug. 21, 1907. Has noticed a dimness of vision in lower field of right eye for ten days. He has been actively engaged in country practice for many years. Has no dissipations, unless it be in his eating. He is a hearty, rugged looking man.

Status Præsens.—The upper temporal quadrant of retina of O. D. is covered with flame-shaped hemorrhages. The veins are very full and tortuous. Nerve head congested. He was referred to Dr. E. C. Hill for physical examination, who reports as follows: Blood pressure, 99 sitting, and 107 after exertion; hemoglobin 90 per cent., red blood cell count, 5,350,000; heart is not dilated but degenerated and weak; urine 26 ounces in twenty-four hours; uric acid, one-third above normal daily maximum; no excess of indican.

Diagnosis.—Passive congestion.

Treatment.—Strychnia and hydrastinin hydrochlorid, and correction of his diet. Dec. 24, 1907, he reported that the vision in the lower nasal field is much improved. Is feeling well. Thinks blood pressure is higher than it was.

CASE 5.—Recurring Retinal Hemorrhage, Associated with Dilated Heart, General Venous Stasis and Autointoxication.—F. S. T., age 26, Sept. 26, 1905. Occupation, farm hand. Has always enjoyed good health. For the last three weeks he has seen a curtain hanging in front of left eye.

Status Præsens.—O. D. V.=5/5. O. S. V.=5/27. With the ophthalmoscope a membrane is seen lying in the anterior portion of the vitreous, which is fixed; also large floating pieces of membrane. The appearances are of a previous vitreous hemorrhage which is being absorbed. The fundus can not be seen. Sent him to Dr. E. C. Hill for physical examination, who reports: Blood tension, 165; heart normal; pulse, 88 sitting, 100 standing; hemoglobin, 90 per cent. The urine

shows considerable excess of indican, uric acid and earthy phosphates.

It was decided to reduce tension somewhat by the use of a very simple diet, also to correct intestinal fermentation by giving him 10 grains of sulphocarbonate of sodium, t. i. d., and 3 grains of calomel once a week. A month later the vitreous was found almost clear and V.=5/9. No fundus lesion could be found. He was told to continue treatment for another month and report. He was not seen again for nine months, when he came because of another vitreous hemorrhage, which had occurred six weeks previously. I found his vitreous full of floating membrane, and a retinal spot could dimly be seen below the macula. Dr. Hill went over him again and found his blood tension 115; heart, much dilated; no indicanuria. He was given 10 drops each of tincture of strophanthus and nux vomica, t. i. d.

Two months later Dr. Hill found his heart still somewhat dilated and had him continue the heart tonics. I found a little floating membrane in the vitreous, and that the spot below the macula was becoming atrophic. I did not see him again until two months later, when he came in with a corneal ulcer caused by the diplobacillus, which quickly healed under zinc. The vitreous is now perfectly clear and O. S. V. is 20/20. There are three spots of atrophic retina to be seen, a small spot just to temporal side of disc, a large one below the macula and a third some distance below the disc.

Four months later he came in saying that the right eye had been blurring for ten days. O. D. V.=20/70; O. S. V.=20/20. The vitreous of right eye contained floating membrane and no fundus details could be made out. Dr. Hill found the heart so dilated that the heart border was one and one-quarter inches to the right of the sternum and one-half inch to the left of the nipple in the sixth interspace. Pulse, 92 sitting, 105 standing. Blood pressure, 132; hemoglobin, 100 per cent. Marked indicanuria. In four days the vitreous had so cleared that a retinal spot could be seen on the nasal side of the disc. He was seen again three months later, when vision in each eye was found to be normal. The blood vessels of the retina have not the appearance of sclerosis. Dr. Hill found the blood pressure 145 mm., and heart almost normal in size.

Three months later he had another attack in the left eye. The vitreous was muddy, but there were no large pieces of floating membrane. Dr. Hill found his heart again badly dilated with a blood pressure of 115. This case illustrates the relation in a young man between vitreous hemorrhage and venous stasis, and how between attacks of the latter his vascular condition was comparatively normal.

CASE 6.—Asthenopia, Associated with Retinal Venous Congestion, Heart Dilatation and General Venous Stasis.—Mrs. M. A. H., age 35, Jan. 9, 1906. Complained of eyes watering and tiring for near work. Thinks vision is poor.

Status Præsens.—The vision of each eye is 5/7.5, with —.50 cyl., ax. 180°. Both nerve heads are congested, and the medium-sized veins are tortuous. I sent her to Dr. Hill for physical examination, who reports as follows: Amount of urine passed in twenty-four hours is 8 ounces, which contains a trace of albumin. No excess of indican nor uric acid. No casts. Blood pressure, 121. Heart very much dilated and leaking all around. The right border one inch to right of sternum. Hemoglobin, 95 per cent. Diagnosis, general venous stasis.

She was referred to her family physician for treatment. Results, unknown.

CASE 7.—Conjunctival Edema, Associated with Heart Dilatation, General Venous Stasis and Autointoxication.—Mrs. F. H. B., age 52, February, 1906. Eyes have been inflamed and in about the present condition for three months. Has worn glasses for three years. Has always enjoyed good health.

Status Præsens.—The eyes look very red and watery. The ocular conjunctiva is edematous and rolls up in front of the lid margins when winking. The palpebral conjunctiva does not appear to be affected. Sent her to Dr. Hill for physical examination and for bacteriologic examination of the ocular secretions, who reports as follows: A smear of the ocular secretions shows fatty degeneration of the epithelial cells, some ordinary staphylococci and proteus bacilli, but no diplobacilli. Pulse, 87 sitting, 90 standing. Heart, one and three-eighths

inches beyond left nipple and one-half inch to the right of the sternum; the apical sounds were somewhat weak. Blood pressure, 121. Red cells numbered 6,200,000; hemoglobin about 95 per cent.; no excess of leucocytes. The urine shows an excess of uric acid and indican. He believes her eye trouble is partly due to uricacidemia, but mainly to stagnant circulation at this altitude. The treatment prescribed was strophanthus and nux vomica. Jan. 4, 1908, a letter from her stated that her eyes are better.

CASE 8.—Asthenopia, Associated with Heart Dilatation, General Venous Stasis.—Mr. G. M., age 42, April, 1907. Has been having so much pain in eyes and back of head that he has been compelled to stop work. His refraction was estimated to be +1.50 W—50 cyl., in each eye, with 20/20 vision. These glasses were ordered. In the macular region of each eye there were seen shimmering lines and irregular splotches of a yellowish color. The veins were greatly dilated and the arteries small. He obtained no relief from wearing the glasses. He was referred to Dr. E. C. Hill for physical examination, who found: Blood pressure, 111; right side of heart greatly dilated. Hemoglobin over 100 per cent. He was placed on appropriate heart tonics, and in the course of two weeks time his headache was relieved and he was able to return to work.

CASE 9.—Episcleritis, Associated with Dilated Heart, General Venous Stasis and Autointoxication.—Mrs. R. C. D., age 38, March, 1907. Her left eye has been subject to attacks of inflammation on the temporal side of the cornea. The eye becomes intensely injected over this portion of the globe, and is very tender and painful. The attack subsides in a few days and the eye becomes almost white; another attack follows in about a week or ten days. She was referred to Dr. Hill for physical examination who reports that she has a dilated heart, which is extended three-eighths of an inch to left of nipple and one-half of an inch to the right of the sternum. Pulse, 95 sitting, 106 standing. Blood pressure, 102. The urine shows a large excess of indican and uric acid. She was placed on strophanthus and sulphocarbolate of sodium and simple diet.

CASE 10.—Retinal Hemorrhage, Associated with High Blood Pressure.—Mrs. Henry W., age 72, Aug. 19, 1907. General habit, plethoric. Has no complaint to make physically. For three months she had had two spots before left eye and has been aware of a gradual failure of vision.

Status Præsens.—O. D. V. = 20/20 W + 2.00. O. S. V. = 20/70 W + 1.75. With ophthalmoscope numerous hemorrhages are seen scattered throughout the retina of left eye. There are no hemorrhages in the right eye, but the blood vessels show strong evidences of sclerosis. She was referred to Dr. F. W. Kenney for physical examination, who reports as follows: Blood pressure, 240; heart normal; no kidney disease. This patient is still under observation. Her blood tension is still high.

CASE 11.—Neuroretinitis, Associated with Chronic Interstitial Nephritis and High Arterial Tension.—Mrs. M. B., age 45, Jan. 19, 1907. Has been suffering of late with severe diffused pain in head. Eyes so sore she can scarcely move them. Has been aware of a central blind spot in right eye for several days.

Status Præsens.—O. D. V. = 20/40; not improved. O. S. V. = 20/20; not improved. With the ophthalmoscope a typical picture of albuminuric neuroretinitis, with hemorrhage, is seen in each eye. The condition was more advanced in her right eye. She was referred to Dr. E. C. Hill for physical examination, who found she was on the verge of uremic convulsions from chronic interstitial nephritis. Her blood pressure was 240. Active measures were taken to reduce the high blood pressure. When it came down to 200, her headache stopped. It was gradually brought down to 185 and 190. The neuroretinitis gradually improved. In two months time the retinal hemorrhages were all absorbed, but the striæ and spots of retinal degeneration, and the opaque sclerosed blood vessels were more marked than ever. Four months later there was scarcely a trace remaining of the retinal degeneration. Vision was 20/20 in each eye, with full fields. She died just a year after my first examination, and enjoyed a fair degree of comfort with good vision until the end.

CASE 12.—Neuroretinitis, Associated with Chronic Interstitial Nephritis and High Arterial Tension.—Mrs. J. I. G., age 50, March 27, 1905. Referred to me by an optician to have her eyes examined for glasses. Has little complaint to make about eyes or her physical condition.

Status Præsens.—O. D. V. = 5/9:5/6 W + 75. O. S. = 1/200; not improved. With the ophthalmoscope a typical picture of albuminuric neuroretinitis was seen in each eye, which was more advanced in the left eye. Numerous retinal hemorrhages in both. She was referred to Dr. James Rae Arneill for physical examination, who reports that she has a dilated leaking heart, blood tension 260. Urine shows albumin and hyaline casts. I never saw this patient again. She felt well and could not believe that she had chronic Bright's disease. Much could have been done to prolong her life had she been willing to follow directions. As it was, I learned that she died after a few months, and that she became almost blind before her death.

CASE 13.—Retinal Hemorrhage, Hemorrhagic Glaucoma and Optic Nerve Atrophy, Associated with Chronic Interstitial Nephritis and High Arterial Tension.—Mr. H. L., age 60, Nov. 16, 1903. Vision has been blurred for ten days. Has always enjoyed good health. Has been a dissipated man as regards alcohol and tobacco. Has not drunk any during recent years.

Status Præsens.—O. D. V. = 5/15. O. S. V. = 5/15. The field of O. D. is contracted temporally to the 5 degree line, the left is normal. The ophthalmoscope shows marked evidences of sclerosis of the retinal vessels in each eye. The disc of O. D. is cupped. In O. S. the nerve head is congested, the adjacent retina edematous, the veins engorged and tortuous, and numerous retinal hemorrhages occupy the inferior nasal quadrant. He was referred to Dr. Hill for physical examination, who reported as follows: Albumin, 1 2/5 per cent. by volume; hyaline, coarsely granular and disintegrated blood casts. Blood tension, 240. He was placed on a strict diet, and was given 3 grains sodium nitrite, t. i. d., and a pill composed of euonymus, aloes and belladonna.

Jan. 18, 1904.—Great loss of vision. O. D. = Fingers at two feet. O. S. = Hand movements. O. S. was inflamed. T + 1. Vitreous so hazy that fundus could not be seen. Iris sluggish and off color. Dr. Hill found the albumin in urine reduced to 1 1/2 per cent. by volume, and his blood tension was 215. The left eye went from bad to worse.

Diagnosis.—Hemorrhagic glaucoma. Enucleation. The vision of his right eye was now so poor that he could not see to get about alone. This was due to a scotoma which encroached on central vision and extended below temporally and nasally, about 20 degrees in each direction. He has been using eserine in oil in this eye, but I can not see that it has been of any service. It is doubtful if there has been, at any time, any increase of tension. For some months this eye remained about the same, and then the central scotoma increased until it occupied the entire temporal field. He has had numerous hemorrhages, but they have not caused hemorrhagic glaucoma as they did with his left eye. I saw him last in July, 1907. He had perception of light only, and numerous retinal hemorrhages were present. Dr. Hill estimated his blood tension several times, and found it to range from 135 to 145. He had a small amount of albumin in his urine. This patient is alive to-day because of the reduction of his blood pressure. His ocular mishaps were most unfortunate, and had he been seen a year or so sooner they might have been prevented.

The above cases serve to show how important are the ocular changes in pointing to the existence of cardiovascular disease. A greater number would needlessly prolong this paper.

DISCUSSION.

DR. LEWIS H. TAYLOR, Wilkesbarre, Pa., said that the profession was coming to recognize more and more the important fact that in certain cases treatment must be directed not to the eye but to the general system. It was hard for the oculist in the face of such conditions to say frankly: "This is a serious eye trouble but I am not competent to treat it." He need not say this if he were physician as well as oculist. Moreover, Dr. Taylor said it was not always convenient and

sometimes not possible to refer patients to expert heart specialists.

Dr. Taylor said that he had had more cases of vascular trouble ending in retinal hemorrhage in the past two years, than ever before in a corresponding period in his whole practice. He related part of the history of a case of retinal hemorrhage. The patient was a strong, hearty man, aged 44 in 1884. He was then fitted with glasses and again twelve years later. In this year he had his first hemorrhage. Two years later the vision rose to 20/XXX some in XX in the patient's right eye, and this was now his better eye. Two years later another attack came and vision of the right eye fell to mere perception of waving hand. Subsequent hemorrhages occurred, ending in hemorrhagic glaucoma for which condition Dr. Taylor enucleated the right eye. The patient's remaining eye had had retinal hemorrhages some years before and was for some time the poorer eye. The patient had always had a full pulse, but soft. His diet was regulated. Fresh hemorrhages, however, occurred in the remaining eye at intervals of two and one years and a severe one in 1907. The patient had been treated by local and general blood-letting, regulation of life, and various medicines. Following the hemorrhage in the spring of 1907, he was practically blind through the greater part of the summer, but rallied and recovered a considerable amount of vision so that he was able to go about alone, but was never able to read. In February, 1908, a day of considerable excitement brought on dimness of the eye, but no hemorrhage was visible. The patient was at once put to bed and placed on treatment to lower arterial tension and on March 2 twenty-four ounces of blood were taken from the arm. The hemorrhages increased with great intensity, causing great pain and high tension which eserine, pilocarpin and dionin could not relieve. The anterior chamber was filled with blood and an iridectomy or sclerotomy was out of the question. Even the skin around the eye was affected with hemorrhages, extending down to the neck. The patient suffered along for a month longer being kept part of the time under the influence of morphia, having at times some improvement to be followed by subsequent hemorrhages. He finally agreed to an enucleation, but in the morning of the day fixed for operation the ball ruptured. After enucleation the hemorrhages continued and were not forcibly checked because of the pain, and because Dr. Taylor considered that it would be better for the patient's head to allow the hemorrhage from the orbit to continue. Oozing of blood, at times free, continued for nearly three weeks, checked finally by adrenalin externally and internally. Dr. Taylor said that the patient had now fully recovered his general health and rode out daily. This patient had been seen by Drs. Knapp, Bull, Risley and de Schweinitz and all had predicted apoplexy as a possibility. Dr. Taylor thought it surprising that the patient had not had a rupture of a blood vessel in the brain before this.

DR. GEORGE E. DE SCHWEINITZ, Philadelphia, said that Dr. Black's insistence on the point that early recognition of angiosclerotic changes might enable physicians to give many years of life to a considerable number of patients deserved to be emphasized. He said that it should also be remembered that even more important than the grosser changes were the early signs of cardiovascular disease, particularly the early signs of general arteriosclerosis, such as lesions of the arterial walls, alternate contractions and dilatations of the veins, indentation of veins by stiffened arteries, and later hemorrhagic extravasations, perivasculitis and silver-wire arteries. Dr. de Schweinitz called attention to still earlier indications, the curious corkscrew appearance of certain arterial twigs, notably those skirting the macula, the crinkled retinal vessels of Alleman's phraseology, the flattening of a vein in contact with an artery, resembling somewhat the appearance of a diminutive strap laid across a solid tube, most frequently evident in the inferior temporal retinal vessels, and the peculiar appearance of the nerve head; with its slightly dull red appearance, differing from a frank congestion precisely as the unhealthy flush of a cheek differs from the brighter color of the normal blush.

Dr. de Schweinitz said that the signs of angiosclerosis looked for by the general clinician—the condition of the pulse,

the character of the heart sounds, particularly that of the first sound at the apex and of the second sound at the aortic area, the increased tidal wave on the sphygmogram, and the elevation of tension recorded by the sphygmomanometer—might result from numerous and varied conditions of the system, organic and nervous in origin, and not necessarily connected with arteriosclerosis. The ophthalmoscopic signs, suggestive or pathognomonic—in their earliest or in their latest stages—he said were certain evidences of local angiosclerosis and frequently of persisting high arterial tension, either of general arteriosclerosis or of its manifestations in special organs, and therefore ophthalmoscopic examination was one of the most ready clinical means for the early detection of important arterial changes. Dr. de Schweinitz believed the ophthalmoscope equal in value if not superior to any of the well-recognized methods of the detection of such a condition; and he thought its findings, unlike those of the sphygmogram and even the sphygmomanometer, diagnostic, if positive. No one, he added, was prepared to say that in the absence of ophthalmoscopic changes there must be absence also of arterial degenerative changes, but in the presence of the former, other things being equal, and local changes in the eye being excluded, the findings, if positive, were diagnostic.

DR. CHARLES H. WILLIAMS, Boston, wished to call attention to the necessity of a very careful ophthalmoscopic examination of these cases. He said that he had a case recently in which the pupils were small and there was general haziness of the vitreous so that it was very difficult to get a good view of the fundus. Under cocaine dilatation, however, he found abundant hemorrhages in the retina and a large number of changes in the retinal vessels, some being obliterated and some reduced to small white lines. This patient was about to sail for Europe. Dr. Williams advised strongly against it and wrote a letter to the patient's family physician. He received an answer taking him to task for having frightened the patient.

DR. EDWARD JACKSON, Denver, illustrated the breadth of this subject and its importance to the ophthalmologist by citing the case of a man, aged 58, whose sight had been growing worse for three months, and who came with vision reduced to 4/22 and 4/12. Ophthalmoscopy showed dilatation of the veins, which were narrowed at the crossings of the arteries. Color-perception was markedly interfered with, but there was still fair perception for yellow and blue near the center of the field of vision. This patient had a blood pressure of 140 mm. Considering the condition of his veins Dr. Jackson inferred that the pressure though not absolutely low, was temporarily low for that particular individual. The patient was referred to Dr. Hill, who found marked dilatation of the heart. Here, Dr. Jackson said, was the practical point: that man was put on restricted diet and proper medicinal treatment, and in a week had improved decidedly. At the end of three months his vision was 4/5. His condition continued good until a year later when he died of some cerebral vascular lesion. Dr. Jackson said that he had seen several cases in which good vision was restored by paying attention to the temporary failure of circulation.

DR. A. R. BAKER, Cleveland, spoke of the importance to the ophthalmologist of the sphygmomanometer. Dr. Baker said that for a great many years he was able to recognize the condition of the fundus in which he had a suspicion of high blood tension. Of course, when there were hemorrhages he could speak positively, but when he could simply say that there was a little tortuosity of the vessels and would urge on the patient the importance of changing his method of living, his advice was usually unheeded. Such patients often congratulated themselves on being particularly well; one, a doctor, slapped himself, and said, "There is not a healthier doctor in Cleveland." He died within the year. When this apparatus was applied to the patient's arm, and the patient was shown his definite high tension and it was explained to him that the life insurance companies would not take him as a risk, Dr. Baker said this offered a definite fact to emphasize the therapeutic measures. Dr. Baker used to try to refer these patients back to the general practitioner, but found that most of them had not such apparatus and didn't know how to use it when loaned to them. Dr. Baker

said that the instrument should be used every day, and that it was not necessary to refer patients to the general practitioner for its application.

DR. G. F. KEIPER, Lafayette, Ind., said that he had found a great deal of angiosclerosis in the young as well as the old and had come to believe that the instrument for measuring blood pressure was just as essential as the ophthalmoscope. Dr. Keiper thought the contention of Janeway correct, namely, that instead of the narrow cuff the broad cuff should be used, and that observations should be based on the use of the broad cuff with the mercury manometer. He added that the kind of instrument used should be recorded because there was bound to be variation of any reading of the Riva-Rocci instrument in comparison with the others.

DR. MELVILLE BLACK wished to emphasize the importance of determining the blood pressure by actual tests. He said that it was impossible to determine it by the appearance of the retinal circulation. High blood pressure or low blood pressure might cause the same appearance in the retina and one should not be positive in the belief that it was always high blood pressure; it might be as Dr. Jackson had said, low for that individual.

HEMORRHAGE INTO THE VENTRICLES.

ITS RELATION TO CONVULSIONS AND RIGIDITY IN APOPLECTIFORM HEMIPLEGIA.*

ALFRED REGINALD ALLEN, M.D.

Instructor in Neurology and in Neuropathology in the University of Pennsylvania.
PHILADELPHIA.

The question to what extent the upper motor neurons are initiative in their function must of necessity be speculative. But since the advent of the theory of definite cerebral localization for well-ascertained function, since the increase of our knowledge of the association tracts and their importance psychologically, the concept of the upper motor neuron as a free agent in initiating its own stimulus becomes more and more untenable. To-day it would seem that the normal motor impulses from the Rolandic region are set in motion from impulses from other parts of the brain transmitted through association tracts.

Now muscular movement—for example, of the right side of the body, face, arm, trunk and leg—may be volitional, in which case it can be considered reflex to psychic stimulation. It may be due to the local action of some circulating poison, either stimulating the perikaryon or its neurons directly, or rendering them so hypersensitive to afferent impulses that, whereas normally there would be no reflex action, in the hyperexcitable condition there is a discharge of motor impulse; this is probably the case in uremia. Intracranial pressure, unless very gradual, is able by direct irritation to cause a motor discharge; and intracranial pressure as well as irritative lesions acting on the right cerebellar lobe can cause stimulation of the motor neurons of the left motor cortex, as in Weber's cases.¹ Electricity applied to the motor cortex or the internal capsule is also capable of causing motor functional activity. Lastly, and from the standpoint of this work of most importance, stimulation or irritation of the high sensory system, either of their perikaryons in the optic thalamus or other basal ganglia, or of their axons in the white mat-

ter of the centrum, is in all probability capable of causing motor impulses—the pure cerebral reflex.

The subject of convulsions and rigidity in apoplectic form hemiplegia is mentioned in a casual unsatisfactory way by Déjerine,² Strümpell,³ Brissaud⁴ and Leube,⁵ all of whom state that these symptoms point to the rupture of the hemorrhage into the ventricles. Nowhere have I been able to find any attempt at a physiologic explanation on this hypothesis.

Before considering my cases it is necessary to deal with the question of the possibility of stimulating the axons of the upper motor neurons when they are separated from their perikaryons. Weber¹ denies this possibility in no uncertain language:⁶

That a direct stimulation of the pyramidal fiber can cause convulsions or rigidity is against all experimental and clinical experiences.

Against this *ex cathedra* dictum of Weber's I would call attention to the work of Beevor and Horsley,⁷ who succeeded in mapping out the internal capsule of the Bonnet monkey and the orang-outang by means of the secondary faradic current.

Hoche, in three articles,⁸ discusses experimental work that he did on the cut end of the spinal cord in criminals who had just been beheaded. He used the faradic current, and although much of his work was negative and seemingly corroborative of Weber's statement, yet in the case where he made his test with least delay after decapitation his result speaks as positively in the affirmative as Weber's in the negative. Hoch's first article⁹ says:

According to the previous accounts I had expected little action; therefore, so much the more surprising was the effect. The corpse, which was lying flat, raised both its arms, bent at the elbowjoint, and with clenched fists, up in the air, the thorax raised itself in inspiration (so that with repeated stimulation the stump of the neck began again to bleed, due to the pump-like action of the thorax) and both legs were in tonic extension.

Hitzig¹⁰ called attention to the fact that the excitability of the central nervous system rapidly failed after exsanguination. This accounts, it seems to me, for many of Hoche's negative results. I have found the lateral columns of the spinal cord in exsanguinated dogs incapable of stimulation. Morat¹¹ says:

2. Déjerine, J.: *Sémiologie du système nerveux*, p. 476.

3. Strümpell: *Lehrbuch der speciellen Pathologie und Therapie*, ed. 13, 1900, III, 477.

4. Brissaud: *Traité de médecine*, 1894, vi, 53.

5. Leube, W.: *Specielle Diagnose der inneren Krankheiten*, ed. 6, 1901, II, 225.

6. "Dass eine direkte Reizung der Pyramidenbahn Zuckungen oder Krämpfe hervorrufen kann, ist nach allen Experimentellen und klinischen Erfahrungen nicht anzunehmen."

7. Beevor, Charles E., and Horsley, Victor: *An Experimental Investigation into the Arrangement of the Excitable Fibers of the Internal Capsule of the Bonnet Monkey (Macacus Sincus)*, Phil. Tr. Roy. Soc. London, clxxxi, B, 49-88; A Record of the Results Obtained by Electrical Excitation of the So-called Motor Cortex and Internal Capsule in an Orang-Outang (*Simia satyrus*), Phil. Tr. Roy. Soc. London, 1890, clxxxi, B, 129-158.

8. Hoche, A.: *Zur Frage der elektrischen Erregbarkeit des menschlichen Rückenmarkes*, Neurol. Centralbl., 1895, No. 14, p. 754; Neurol. Centralbl., 1900, p. 994; *Ueber Reizungsversuche am Rückenmarke von Enthaupteten*, Berl. klin. Wchnschr., 1900, No. 22, p. 479.

9. "Nach den bisher vorliegenden Angaben hatte ich mir wenig Wirkung versprochen: um so überraschender war der eintretende Effect: der flach liegende Leichnam hob beide Arme mit gebeugtem Ellenbogengelenk und geballten Fausten in die Höhe, der Brustkorb hob sich inspiratorisch (so dass bei wiederholter Reizung, Dank der Pumpwirkung des Thorax, der Halsstummel wieder anfang, zu bluten) und beide Beine geriethen in Strecktonus."

10. Hitzig, Edward: *Physiologische und klinische Untersuchungen über das Gehirn*, Berlin, 1904, part 1, p. 23.

11. Morat, J. P.: *Physiology of the Nervous System*, 1906, p. 279.

* Read in the Section on Nervous and Mental Diseases of the American Medical Association, at the Fifty-ninth Annual Session, held at Chicago, June, 1908.

* From the Department of Neurology and the Laboratory of Neuropathology of the University of Pennsylvania.

1. Weber, L. W.: *Gleichseitige Krämpfe bei Erkrankung einer Kleinhirnhemisphäre*, Monatschr. f. Psychiat. u. Neurol., 1906, xix, 478.