the tonsil, settle there and cause its inflammatory enlargement. This would also explain the peculiar position of the red band of oral sepsis on the soft palate mentioned above.

3. *Physiological Function of the Tonsil and other Adenoid Tissue in the Pharynx.*

In children the lymphadenoid tissue is more extensive than in the adult and is spread out on the wall of the pharynx, forming a complete ring continuous with the adenoid tissue in the naso-pharynx. The lower limit of adenoid tissue is very irregularly defined and may in cases reach the glottis. As age increases this adenoid tissue disappears and remains chiefly in the tonsil and the so-called lateral bands of the pharynx. It is possible that this disappearance is the result of the frequent attacks of catarrhal conditions in children, such as measles, scarlet fever, sore-throats and common colds, &c. Each attack would cause a degeneration of certain portions of adenoid tissue which would not be replaced. I would look upon the adenoid tissue in the pharynx as a protective barrier in which the attacking micro-organisms are dealt with actively, in which leucocytosis is most active and where immunisation of the body can be developed. Every attack of the invading micro-organisms would cause an inflammatory enlargement of the adenoid tissue, and in the fight certain portions of the adenoid tissue would be killed, so that the power of the adenoid tissue to resist invasion would be diminished by each attack. In addition, however, the numerous crypts existing in adenoid tissue act as good covering for the attacking organisms where they mass together for a further onslaught, which is rendered the more easy on account of the lessened resistance already produced, and once the adenoid tissue is severely degenerated it is not only useless as a protecting barrier but provides a point of vantage for further invasion.

The tonsil itself I would look upon as the special adenoid (lymphatic) gland of the mouth, lying openly in it, specialised it is true, but just in the same way that a Peyer's patch in the intestine is specialised. A tonsillitis would thus be an adenitis, and a peritonsillar abscess a periadenitis. As a rule the tonsil and other adenoid tissue in the pharynx act as sufficient scavengers for the organisms in the mouth, and at the most there is simply inflammatory enlargement caused. But occasionally the tonsil and adenoid tissue are overwhelmed in the attack and the next link in the lymphatic chain becomes involved—viz., the glands at the angle of the jaw—and these take up the fight, undergoing enlargement in the process. If the attack is still more severe then the organisms no longer confine themselves to the mouth, but travel either upwards to the naso-pharynx or downwards to the stomach and lungs. Naturally when the tonsils and other adenoid tissue are already badly degenerated and act as harbourages for micro-organisms, the barrier power is extremely diminished and attacks of naso-pharyngitis and bronchitis become more and more frequent. I believe that the spread of inflammatory conditions from the oro-pharynx to the naso-pharynx by continuity is more frequent than is supposed. It must be remembered that Greville MacDonald long ago pronounced the nasal cavity to be normally free from organisms, while the mouth is normally crowded with them.

4. *Treatment and Indications for Such Based upon the Above.*

When inflammatory enlargement of the tonsils is recognised the treatment is that of enlarged glands in any part of the body: 1. Remove the cause. 2. If that is not sufficient, then the enlarged degenerated tonsil should be removed by surgical interference. Removal of the cause is the prevention or cure of oral sepsis, the extraction of septic molars, and careful mechanical cleansing of the teeth by brushing. This must precede surgical removal of the tonsil. In fact, the reckless way in which we all have removed tonsils without attending to the causal oral sepsis cannot be too strongly condemned. It is to my mind the explanation of why so many cases show unpleasant complications and bad results. Removal of the tonsils would be less indicated by their actual increase in size than by the evidence of their inefficiency, as shown by swelling of the glands at the angle of the jaw and by frequent attacks of naso-pharyngitis and of bronchitis. The use of the tooth-brush is essential before, during, and after tonsillotomy, and as a means of preventing attacks upon, and preserving the integrity of, the adenoid protecting ring cannot be too strongly advocated.

Ipswich.

A CASE OF PRIMARY SARCOMA OF THE DUODENUM.

BY H. M. MACKENZIE, M.B. EDIN., D.P.H. CANTAB.,
CAPTAIN, I.M.S.; PROFESSOR OF PATHOLOGY, MEDICAL COLLEGE,
LAHORE, INDIA.

THE comparatively rare occurrence of primary sarcomatous growths originating in the small intestine, and more especially in the duodenum, lends some degree of interest to the present case.

The patient, a Hindu labourer, aged 18 years, was admitted to the Mayo Hospital, Lahore, on Dec. 21st, 1908, under the care of Major H. G. Melville, I.M.S., complaining of pain in the abdomen and back and swelling of the feet. His family and personal history were good and up to the date of the present illness his health had always been fair. The present illness began some 12 days before admission, when he first felt pain in the abdomen and back, and some days later he noticed swelling of his feet. No treatment was adopted, but as the symptoms became more severe he came to hospital. On admission the patient, an anæmic, poorly developed youth, was suffering from pain in the abdomen, especially marked in the right hypochondriac and right lumbar regions and also from pain in the back in the middle line, the pain in both places being increased on pressure. He was markedly jaundiced and passed clay-coloured stools, but there was no diarrhoea or melæna. The urine was normal. The appetite was diminished and thirst was increased and the temperature was subnormal. On examination of the abdomen a large, moderately firm mass, moving with respiration and mobile on palpation, was found to occupy the right hypochondriac, right lumbar, and part of the epigastric and umbilical regions, as shown in the illustration, the inner margin being

distinctly nodular. The lower edge of the liver was palpable in parts, other parts being somewhat obscured, and light percussion gave a tympanitic note between the upper margin of the swelling and the edge of the liver, except on the right side where the dullness was continuous. There was some fluctuation of the swelling towards the upper and inner margin. The mass was diagnosed as a malignant growth of the right kidney, probably sarcomatous, with dilatation of the gall-bladder from obstruction of the common bile duct. General symptomatic treatment was adopted, but the condition of the patient steadily grew worse and he died on Jan. 12th, 1909.

The post-mortem examination was conducted 24 hours after death. The body was that of an emaciated, poorly developed youth and showed very marked jaundice. Post-mortem lividity was slight and rigor mortis was present in the lower limbs only. The abdomen, which was retracted, showed slight fulness on the right side but no other abnormality externally. On opening the

abdomen a large, fairly firm, but very friable tumour was found occupying the upper two-thirds of the right half of the abdominal cavity, extending downwards for about 5½ inches from the lower margin of the liver and from the right flank inwards nearly to the umbilicus. The duodenum disappeared into the centre of the mass, and several coils of small intestine were firmly adherent to its antero-internal surface, as was also the cæcum in the region of the appendix. The right kidney was considerably enlarged and partially imbedded in the growth from it could not be separated, but there was a sharp line of demarcation between the kidney substance and that of the tumour, and sections examined microscopically showed vestiges of the kidney capsule and no infiltration of the kidney. The right suprarenal body was enlarged to twice the normal size but was not adherent, while the ureter was patent and not affected. The pancreas was somewhat enlarged and adherent to the posterior surface of the growth. There was no connexion between the tumour and the liver, but the gall-bladder, which projected on to the anterior surface of the tumour, was dilated, containing over 4 ounces of thick bile, due to occlusion of the common bile-duct from pressure exerted by the tumour. The distended gall-bladder accounted for the fluctuation which was made out on physical examination. On following the duodenum into the tumour and laying it open, it was found to dilate into a cyst-like cavity some 3 inches in diameter, with rough, irregular walls, from the lower end of which it emerged again as a recognisable structure. Microscopical examination showed the tumour to have no capsule and to be composed of a stroma of fibrous tissue containing poorly developed blood-vessels and having the spaces filled with small cells. Some of these were irregular in outline, but the majority were of the round-cell type, with large deeply staining nucleus and faintly staining protoplasm. A delicate fibrillated intercellular substance could also be distinguished. The right kidney and the liver presented no specially noteworthy appearances, the kidney showing slight breaking down of tubular epithelium in places and the liver showing commencing monolobular cirrhosis with dilatation of bile capillaries. The pancreas, although adherent, showed no evidence of cellular infiltration. Successive sections of the duodenal wall from the point where it entered the tumour to the point of maximum dilatation of the cavity showed gradual disappearance of the structure of the glandular epithelium, followed by that of the muscular wall, until at the widest portion of the cavity there was no trace of intestinal wall, the wall of the cavity being composed wholly of tumour tissue of the nature described. Smears made from the contents of the cavity showed the ordinary intestinal bacterial flora and also many loose cells resembling those of the tumour in character. Some of the mesenteric glands showed enlargement, but microscopically no metastases were found, the enlargement being of the nature of a simple lymphatic hyperplasia. No secondary growths were found in any of the organs. Death appeared to be due to general cachexia and exhaustion, hastened probably by the toxæmic condition caused by the obstruction of the common bile duct. The rapid development of the primary tumour and the absence of any metastatic growths are noteworthy features in the case.

I am indebted to Major Melville for the report of the clinical symptoms and physical signs.

A CASE OF INHERITED TACHYCARDIA.

By JAMES KIRKLAND, M.B., C.M. GLASG.

THE following case may be of interest to readers of THE LANCET. It clearly demonstrates that heredity plays a very active part in the etiology of tachycardia. I cannot find any other case reported having such a remarkable family history. Dr. A. W. Falconer in the *Practitioner* for February, 1909, records a case occurring in mother and daughter, and several other cases have from time to time been reported where the disease has been passed from a parent, usually the mother, to one or two children. But this case is remarkable in that the inheritance can be traced from both parental branches, and descends to almost all the issue to the fourth generation.

The particular member of the family now under my care, and whose case is here reported, is a male, 56 years of age, single, of no regular occupation, of independent means, of

average build, but whose weight varies considerably (e.g., in February, 1906, he weighed only 6 stones 9½ pounds, in October, 1907, he weighed 12 stones 3½ pounds, and his present weight is 10½ stones) and who is 5 feet 10½ inches in height. He consulted me for nervous depression and tachycardia early in the present year. He has never been strong, having had pneumonia when six months old and peritonitis at 11 years of age. In 1897 he had a very bad attack of heart trouble—tachycardia, with hæmorrhage, supposed to have been from the lungs, but there was no trace of tubercle bacilli in the sputum, nor is there any now. This recurred in January, 1898. After several minor attacks he had a very serious one in October, 1900, followed by collapse. No record has been kept as to the rate of the heart beat, nor the length of time the attacks lasted. In 1901 he had nine attacks of a more or less serious nature, lasting from 18 to 21 hours. Bromide of potassium and morphine were used, evidently with success. In 1902 he had six bad attacks preceded and accompanied by "green sickness" and dyspepsia (flatulent). These lasted from five and a half to 23 hours. Bromides and morphine were pushed more strongly than in 1901. In February, 1903, there was a very bad attack and for the first time dyspnoea was recorded. In March, 1903, the patient had two attacks. On the first occasion irregularity was noticed, and this has remained a feature of the attacks ever since. On the latter occasion it is said that a gall-stone was passed. The "dark green sickness" was excessive and nothing seemed to give any relief until this stone passed. In April, 1903, "congestion of the base of the brain" was noted and strychnine was used for the first time. The attacks now became more frequent; all seem to have been accompanied by more or less dyspnoea, flatulent dyspepsia, and sleeplessness. There has also been a cough, with profuse frothy expectoration free from tubercle bacilli or blood. The irregularity at first would pass off with the attacks, but it is now (1909) constant. There was a rise of temperature on several occasions, but it was due, I think, to other causes. The patient could not eat anything but the lightest food on account of dyspepsia and consequent "heart hurry." One peculiarity that was noticed was the singular effect which hair-cutting or beard-trimming had upon the patient. It seemed to make him feel sick and faint, and he looked pale and haggard. So marked was this that his friends dreaded the ordeal and he himself postponed it as long as possible.

When the patient came to place himself under my care his heart beats varied from about 120 to 240 (the latter when an attack came on). There is a systolic mitral murmur and strong pulsation in the right side of the neck; on the left it is much less marked. The right lobe of the thyroid gland is enlarged and evidently acts as an irritant to the pneumogastric nerve; pressure upon this gland seems to excite the patient and to increase the frequency of the heart beat. The lungs are free from any defect in the apices, but there are moist râles in both bases. There is a varying amount of flatulent dyspepsia and a troublesome cough with profuse expectoration (free from tubercle bacilli). The urine is normal (specific gravity 1022) and the bowels are regular. The nervous system is far from satisfactory and the patient is melancholy and depressed. There is marked irregularity of heart action, which is greatly increased on any slight exertion.

Since I have had the patient under my care he has had only one attack of a serious nature. The heart beats were between 220 and 240. It was difficult to count owing to the irregularity. Dyspnoea was most distressing, increased by the cough and any slight exertion, even eating or drinking. The voice was much affected, amounting at times to aphonia. There were great abdominal discomfort and flatulent dyspepsia. The patient had to be propped up in bed, and was most comfortable and easy in this position and the dyspnoea was less severe. The attack came on suddenly after a walk that was beyond his strength, and lasted 28 hours, passing off as quickly. I tried many remedies without success, but finally a hypodermic injection of strychnine together with a mixture containing apomorphine hydrochlor. (1/16 grain), caffeine hydrobromide, &c., seemed to cut the attack short. I am now using 15 drops of digalen (a continental remedy and an extract of digitalis) thrice daily. It seems to have a very good effect upon the heart, which has greatly improved in every way since I began using it.

Family history.—This is most interesting, and I can find no case recorded with so marked a family idiosyncrasy towards tachycardia or so clear a history of heredity in