cated, the spasm takes place at the glottis; and not unfrequently the stammering, whether occurring in connexion with vowels or consonants, is due to a sudden inspiration. But whatever part the spasm may affect primarily, the arrest may consist either in a sudden, simple, more or less prolonged spasm, or in a series of such spasms in rapid sequence, during which the implicated literal sound undergoes more or less frequent repetition. And, further, there is generally a tendency, especially if the spasm be prolonged, for a greater or less number of other groups of muscles to be involved. So that if, for example, the spasm commences at the base of the tongue, we may find in some cases that the mouth opens widely and remains in that position, the muscles of expression work convulsively, the glottis contracts, respiration becomes arrested, and finally spasmodic movements of remote parts become also excited. It would be easy to write a detailed description of some of these cases, and to show how individual examples differ from one another; but the final result would be, I think, to prove that their resemblances are far more important than their differences, and that the pathological explanation is the same in all. Speech, as I have shown, is, even in its mechanical details, a very complicated art; it requires, for its perfect performance, not only that the respiratory apparatus, the laryngeal instrument, and the organs of articulation shall uniformly co-operate with the greatest nicety under conditions which are ever varying, but, as regards the last organs especially, that they shall pass from one arrangement to another arrangement with unflinching ease, certainty, and rapidity. It is generally in connexion with complicated, rapidly varying muscular combinations, which are rendered easy of performance and more or less automatic by long practice, that momentary spasms or hitches are liable to occur and mar the performance. It is thus that occasionally the swordsman's arm is apt to become arrested in some particular attitude, the pianist has to give up his playing because of some recurring sudden inco-ordination, and the scribe especially is seized during his work with momentary spasm or cramp which checks or spoils his writing. And it is, doubtless, thus that spasm in the utterance of articulate language takes place, and that this spasm (although it may commence in any of the three mechanisms concerned in speech) almost always begins either in the organs of articulation themselves or in connexion with the mutual co-ordination of the several factors of speech—respiration, phonation, and articulation—and at the commencement of words or syllables. It has been asserted that stammering does not occur when the sufferer whispers or when he sings or intones. Neither of these statements, however, is accurate. Whispering is frequently attended with stammering, and, although the singing voice is usually free from it, even in aggravated cases, it undoubtedly sometimes suffers. In concluding this subject, I may remark that I do not pretend to throw any light on the treatment of this affection. Inasmuch, however, as it does not depend on any structural defect of the organs concerned in speech, local remedies, with the possible exception of the continuous galvanic current, are not likely to be of any service. It may, however, be improved, if not cured, by close attention to the management of the breath in speaking, by carefully practising those combinations of letters which are specially liable to give rise to stammering, and by the adoption of a deliberate and carefully balanced method of enunciation.

FOREIGN BODY IN THE HEART.

By SIR JOSEPH FAYRER, K.C.S.I., M.D., F.R.S.

THE following is a more detailed account of the remarkable case of a bullet found in the heart to which I briefly referred at a recent meeting of the Pathological Society. I am indebted to Dr. Ewart's excellent descriptive catalogue of the pathological museum of the Medical College in Calcutta for such particulars as had escaped my memory. The man came under my observation and care at the Field Hospital, Rangoon, during the operations there in 1852.

Israel H—, a private in H.M.'s 80th Regiment, was wounded at the storming of the Great Pagoda at Rangoon, on April 14th, 1852, and was brought to the field hospital. The ball had entered the left shoulder, passing down into

the chest. Blood and air were issuing from the wound, and did so for some days afterwards; there were all the symptoms of a wounded lung. He remained in the field hospital at Rangoon for a fortnight, and was then transferred to the convalescent dépôt at Amherst, where he came under the care of Assistant-Surgeon White. His cough, expectoration, and emphysema had diminished. On his arrival at Amherst he was much better, and very hopeful of recovery. He is described thus:—"His cough was slight; there was no blood in the expectoration; there was great emaciation; small, quick pulse; clean tongue; the skin was cool; the bowels regular; the wound looking healthy; spirits good; there was dulness over the left side; slight respiratory murmur only below clavicle." On May 12th, the wound is reported as closed. Febrile symptoms then came on. On the 13th, he had a restless night from cough and dyspnoea; there was some bloody expectoration. In this state he continued till May 24th, after which he enjoyed comparative ease until June 3rd, when the dyspnoea and bloody expectoration returned. On June 11th emphysema extended from the situation of the original wound into the axilla and over the left side of the thorax. On the 14th he had profuse sweating; no respiratory murmur on the left side. He had become very weak and emaciated, and died on the 24th, having survived seventy-two days after the wound.

Post-mortem examination made by Mr. WHITE six hours after death.—Body much emaciated. The bullet had passed into the thorax between the third and fourth ribs, proceeding in its course downwards and inwards. The pleura was firmly adherent, and formed a cavity which extended from the first to the seventh ribs, and from the spinal column to the cartilages of the ribs. This contained about a pint of pus. The left lung was impervious to air throughout. A small portion of the cloth of the jacket was lying at the orifice of a canal situated about the middle of the lung. This passage stopped short near the course of the pulmonary veins, when all further traces of its passage were lost. On raising the heart a hard substance was felt at its apex, which, on examination, proved to be the bullet in the left ventricle, at its apex, crossed and recrossed by chordæ tendineæ and carneæ columnæ, which secured it firmly in position. The heart was perfect in every respect, and the only conclusion that could be arrived at was that the bullet must have perforated (perhaps by ulceration) one of the pulmonary vessels, and thus passed into the auricle, and so by its weight had dropped into the ventricle.

The above are nearly the words of the report in the catalogue, and no doubt give an accurate description of the case from the time the man left the field hospital at Rangoon, then under my charge, until he died at Amherst.

I was not present at the autopsy, but have often seen the heart in the museum of the Medical College in Calcutta, where it is one of many interesting objects.

I made some experiments later at the Medical College, to ascertain whether it was possible for a bullet to find its way into the ventricle in this fashion, and found that if the bullet were introduced into the larger vessels, it might be made to drop into the heart, thus confirming the view that it had been so in the case of poor Israel H—.

Granville-place.

THE SULPHUR-WATERS OF THE PYRENEES.

By G. V. POORE, M.D., F.R.C.P.,

ASSISTANT-PHYSICIAN TO UNIVERSITY COLLEGE HOSPITAL, ETC.

(Concluded from p. 625.)

THE power which the sulphides possess of stimulating nutrition in scrofulous subjects whose nutrition is sluggish must have been often observed by anyone who has made a patient trial of their therapeutic value. At the Royal Infirmary for Children I have repeatedly ordered them for patients with enlarged glands; and although I have frequently failed to obtain any result, I have, at least as frequently, seen, as an immediate consequence of their administration, activity replace torpor and absorption or suppuration be quickly produced; these effects being usually accompanied by cutaneous hyperæmia and sometimes by a slight tingling of the skin.