PROCEEDINGS OF THE TWENTY-EIGHTH ANNUAL MEETING OF THE GEOLOGICAL SOCIETY OF AMERICA, HELD AT WASHINGTON, DISTRICT OF COLUMBIA, DECEMBER 28, 29, AND 30, 1915.

CHARLES P. BERKEY, *Secretary pro tem.*

CONTENTS

	Page
Session of Tuesday, December 28.....	5
Report of the Council.....	5
Secretary's report.....	5
Treasurer's report.....	7
Editor's report.....	9
Election of Auditing Committee.....	11
Election of officers.....	11
Election of Fellows.....	12
Memorial of Theodore B. Comstock (with bibliography); by Heinrich Ries.....	12
Memorial of Orville A. Derby (with bibliography); by J. C. Branner.	15
Memorial of Joseph Austin Holmes (with bibliography); by J. H. Pratt.....	22
Memorial of William John Sutton; by W. F. Robertson.....	35
Memorial of A. B. Willmott (with bibliography); by A. P. Coleman..	37
Titles and abstracts of papers presented before the morning session and discussions thereon.....	38
Geographic history of the San Juan Mountains since the close of the Mesozoic era [abstract]; by Wallace W. Atwood and Kirtley W. Mather.....	38
Dominantly fluvial origin, under seasonal rainfall, of the old red sandstone [abstract]; by Joseph Barrell.....	39
Influence of Silurian-Devonian climates on the rise of air-breathing vertebrates [abstract]; by Joseph Barrell.....	40
Some littoral and sublittoral physiographic features of the Virgin and northern Leeward Islands and their bearing on the coral-reef problem [abstract]; by Thomas Wayland Vaughan.....	41
Coral-reef problem [abstract]; by W. M. Davis.....	46
Tertiary-Quaternary orogenic history of the Sierra Nevada in the light of recent studies in the Yosemite region [abstract]; by F. E. Matthes.....	46
Titles and abstracts of papers presented before the afternoon session and discussions thereon.....	47

	Page
Geological transformations of phosphorus [abstract]; by Elliot Blackwelder.....	47
Diffusion in silicate melts [abstract and discussion]; by N. L. Bowen.....	48
Petrography of the Pacific islands [abstract and discussion]; by R. A. Daly.....	48
Some factors which affect the deposition of calcium carbonate [abstract]; by John Johnson.....	49
Specific weight of drill cores [abstract]; by Alfred C. Lane.....	49
Chemical and mineralogical composition of meteorites [abstract and discussion]; by George P. Merrill.....	50
Importance of water as a magmatic constituent [abstract and discussion]; by George W. Morey.....	50
"Pele's tears" and their bearing on the origin of australites [complete paper and discussion]; by Elwood S. Moore.....	51
Triassic igneous rocks in the vicinity of Gettysburg, Pennsylvania [abstract]; by George W. Stose and J. Volney Lewis.....	55
Desert regolith and its genetic relations to maximum epirotic deposition [abstract]; by Charles Keyes.....	57
Origin of foliation in the Precambrian rocks of northern New York [abstract and discussion]; by William J. Miller.....	57
Landslides in unconsolidated sediments [abstract]; by David H. Newland.....	58
Annual dinner.....	60
Session of Wednesday, December 29.....	60
Report of Auditing Committee.....	60
Titles and abstracts of papers presented before the morning session and discussions thereon.....	60
Ferrous iron content and magnetic properties of the natural oxides of iron as an index to their origin and history [abstract]; by R. B. Sosman and J. C. Hostetter.....	60
Variable composition of melanochalcite [abstract]; by W. F. Hunt and E. H. Kraus.....	61
Definition and determination of the mineral hydroxides of iron [abstract]; by H. E. Merwin and Eugen Posnjak.....	61
Saline fumarole deposits of the south Italian volcanoes [abstract]; by Henry S. Washington.....	61
Crystals and crystal forces [abstract]; by F. E. Wright.....	62
Extension of the Montana phosphate deposits northward into Canada [abstract and discussion]; by Frank D. Adams and William J. Dick.....	62
Emerald deposits of Muzo, Colombia [abstract]; by Joseph E. Pogue.....	63
Crystalline marbles of Alabama [abstract]; by William F. Prouty.....	63
Oriskany iron ore [abstract]; by R. J. Holden.....	64
Geologic map of the Fort Hall Indian reservation [abstract]; by George R. Mansfield.....	64
Preliminary geologic map of the Wayan quadrangle, Idaho-Wyoming [abstract]; by George R. Mansfield.....	65

	Page
Glacial lakes and other glacial features of the central Adirondacks [abstract]; by Harold L. Alling.....	65
Pleistocene features in the Schenectady-Saratoga Glen Falls section of the Hudson Valley [abstract]; by Herman L. Fairchild.....	65
Pleistocene uplift of New York and adjacent territory [abstract and discussion]; by Herman L. Fairchild.....	66
Studies of glaciation in the White Mountains of New Hampshire [abstract]; by James Walter Goldthwait.....	67
Glaciation and stormy period of the fourteenth century [abstract]; by Ellsworth Huntington.....	67
Pleistocene deposits of Minnesota and adjacent districts [abstract]; by Frank Leverett.....	68
Resolution regarding the taking of expert testimony.....	69
Titles and abstracts of papers presented before the afternoon session and discussions thereon.....	70
Pennsylvanian of Tennessee [abstract]; by L. C. Glenn.....	70
Subdivisions of the Thaynes limestone and Nugget sandstone, Mesozoic, in the Fort Hall Indian reservation, Idaho [abstract]; by George R. Mansfield.....	70
Stratigraphy of some formations hitherto called Beckwith and Bear River, in southeastern Idaho [abstract]; by George R. Mansfield and P. V. Roundy.....	70
Sedimentation along the Gulf coast of the United States [abstract]; by E. W. Shaw.....	71
Relative age of the Detroit River series [complete paper and discussion]; by Clinton R. Stauffer.....	72
Recession of Niagara Falls remeasured in 1914 [abstract]; by J. W. Spencer.....	78
Terrestrial stability of the Great Lake region [abstract]; by J. W. Spencer.....	79
Scour of the Saint Lawrence River and lowering of Lake Ontario [abstract]; by J. W. Spencer.....	79
Pleistocene drainage changes in western North Dakota [abstract]; by Arthur G. Leonard.....	80
Landslips and laminated lake clays in the basin of Lake Bascom [abstract and discussion]; by Frank B. Taylor.....	81
Types of loess in the Mississippi Valley [abstract]; by B. Shimek.....	82
Dry land in geology; presidential address by Arthur P. Coleman.....	82
Session of Thursday, December 30.....	83
Titles and abstracts of papers presented before the morning session and discussions thereon.....	83
Stages in the geologic history of Porto Rico [abstract and discussion]; by Chester A. Reeds.....	83
Cretaceous of Alberta, Canada [abstract]; by Joseph H. Sinclair.....	85
Sedimentary succession in southern New Mexico [abstract]; by N. H. Darton.....	86
Divisions and correlations of the Dunkard series of Ohio [complete paper]; by Clinton R. Stauffer.....	86
Silurian system of Maryland [abstract and discussion]; by C. K. Swartz and W. F. Prouty.....	89

	Page
Homocline and monocline [complete paper and discussion]; by Reginald A. Daly.....	89
Occurrence of intraformational conglomerate and breccia [ab- stract]; by F. V. Emerson.....	93
Keweenaw fault [complete paper]; by Alfred C. Lane.....	93
Some structural features in the Green Mountain belt of rocks [abstract]; by C. E. Gordon.....	101
Faulting in north-central Kentucky [complete paper and discus- sion]; by Arthur M. Miller.....	101
Mechanics of intrusion of the Black Hills Precambrian granite [abstract and discussion]; by Sidney Paige.....	104
Precambrian structure of the Black Hills, South Dakota [ab- stract]; by Sidney Paige.....	106
Titles and abstracts of papers presented before the afternoon session and discussions thereon.....	106
Rectilinear features in the eastern Catskills [abstract and dis- cussion]; by George H. Chadwick.....	107
Physiographic evidence of recent subsidence on the coast of Maine [abstract]; by Charles A. Davis.....	108
Physiographic notes on the White Mountains [abstract]; by Douglas W. Johnson.....	108
Position of the New England upland in the White Mountains [abstract]; by Armin K. Lobeck.....	108
Study of ripple-marks [abstract]; by Walter A. Bucher.....	109
Dead Lake of the Chipola River, Florida [abstract]; by E. H. Sellards.....	109
Tectonic lines in the Hawaiian Islands [abstract]; by Sidney Powers.....	109
Banded glacial slates of Permocarboniferous age, showing possi- ble seasonal variations in deposition [abstract and discussion]; by Robert W. Sayles.....	110
Geology of the Lake Iditarod region, Alaska [abstract]; by Philip S. Smith.....	114
Characteristics of the soil and its relation to geology [abstract]; by C. F. Marbut.....	114
Geologic significance and genetic classification of arkose deposits [abstract]; by Donald C. Barton.....	115
Some features of the Kansan drift in southern Iowa [abstract and discussion]; by George F. Kay.....	115
Triassic rocks of Alaska [abstract]; by George C. Martin.....	119
Lithogenesis and stratigraphy of the red beds of southeastern Wyoming [abstract]; by S. H. Knight.....	120
Experiment in the graphic presentation of the economic geology of bedded deposits [abstract]; by George H. Ashley.....	122
Brecciation effects in the Saint Louis limestone [abstract]; by Francis M. Van Tuyl.....	122
Vote of thanks.....	124
Register of the Washington meeting, 1915.....	125
Officers, Correspondents, and Fellows of the Geological Society of America.	127

SESSION OF TUESDAY, DECEMBER 28

The first general session of the Society was called to order at 9.15 o'clock a. m., Tuesday, December 28, at the George Washington University Medical School, Washington, District of Columbia, by President Coleman.

The report of the Council for the year ending November 30, 1915, was presented as follows:

REPORT OF THE COUNCIL

To the Geological Society of America, in twenty-eighth annual meeting assembled:

The regular annual meeting of the Council was held at Philadelphia, Pennsylvania, in connection with the meeting of the Society, December 29-31, 1914.

The details of administration for the twenty-seventh year of the existence of the Society are given in the following reports of the officers:

SECRETARY'S REPORT

To the Council of the Geological Society of America:

Meetings.—The proceedings of the annual general meeting of the Society held at Philadelphia, Pennsylvania, December 29-31, 1914, have been recorded in volume 26, pages 1-128; of the Cordilleran Section, pages 129-140, and of the Paleontological Society, pages 141-170, of the Bulletin.

Membership.—During the past year the Society has lost four Fellows¹ by death—Theodore B. Comstock, Orville A. Derby, Joseph A. Holmes, and William J. Sutton. One resignation has become effective. The names of the nineteen Fellows elected at the Philadelphia meeting have been added to the list, all of them having completed their membership according to the rule. The present enrollment of the Society is 376. Six candidates are before the Society for election and several applications are under consideration by the Council.

Distribution of Bulletin.—There have been received during the year 6 new subscriptions to the Bulletin, and 9 subscriptions have been discontinued, making the number of subscribers 115.

The irregular distribution of the Bulletin during the past year has been as follows: Complete volumes sold to the public, 15; sold to Fellows, 1; sent out to supply deficiencies, 1, and delinquents, 5; brochures

¹ Since the meeting the Secretary has received notice of the death of Frank A. Hill.

sent out to supply deficiencies, 8, and delinquents, 44; sold to Fellows, 3; sold to the public, 27.

Bulletin sales.—The receipts from subscriptions to and sales of the Bulletin during the past year are shown in the following table:

Bulletin Receipts, December 1, 1914–November 30, 1915

	Complete volumes.			Brochures.			Grand total.
	Fellows.	Public.	Total.	Fellows.	Public.	Total.	
Volume 1...							
Volume 2...							
Volume 3...							
Volume 4...		\$7.50	\$7.50				\$7.50
Volume 5...							
Volume 6...		7.50	7.50				7.50
Volume 7...		7.50	7.50				7.50
Volume 8...		7.50	7.50				7.50
Volume 9...		7.50	7.50				7.50
Volume 10...							
Volume 11...		7.50	7.50				7.50
Volume 12...		7.50	7.50				7.50
Volume 13...		7.50	7.50				8.70
Volume 14...					\$1.20	\$1.20	.90
Volume 15...		7.50	7.50		.90	.90	8.60
Volume 16...		7.50	7.50		1.10	1.10	7.50
Volume 17...							1.05
Volume 18...					1.05	1.05	1.80
Volume 19...					1.80	1.80	2.40
Volume 20...		7.50	7.50		2.40	2.40	8.90
Volume 21...					1.40	1.40	7.15
Volume 22...					7.15	7.15	5.95
Volume 23...				\$4.15	1.80	5.95	.80
Volume 24...		15.00	15.00		.80	.80	24.50
Volume 25...	\$7.50	22.50	30.00	1.95	7.55	9.50	38.25
Volume 26...		835.00	835.00		8.25	8.25	839.50
Volume 27...		60.00	60.00		4.50	4.50	63.75
Total....	\$7.50	\$1,015.00	\$1,022.50	\$6.10	\$43.65	\$49.75	\$1,072.25
Index 1-10...		2.25	2.25				2.25
Index 11-20...		3.50	3.50				3.50
Total....	\$7.50	\$1,020.75	\$1,028.25	\$6.10	\$43.65	\$49.75	\$1,078.00

Receipts for the fiscal year..... \$1,078.00

Previously reported..... 18,300.89

Total receipts to date..... \$19,378.89

Charged, but not yet received: On 1911 account..... 7.50

On 1915 account..... 4.90

Total sales to date..... \$19,391.29

Two subscriptions to volume 26 are still to be paid for.

Expenses.—The following table gives the cost of administration and of Bulletin distribution during the past year:

EXPENDITURES OF SECRETARY'S OFFICE DURING THE FISCAL YEAR ENDING NOVEMBER
30, 1915

Account of Administration

Note book.....	\$0.20
Printing (including annual meetings of 1914 and 1915).....	94.46
Messenger service.....	1.00
Telegrams	10.45
Telephone charges.....	1.03
Postage	37.36
Express charges.....	4.86
Letter-heads	2.75
Ribbon for typewriter.....	.75
Post-cards	5.50
Binding three copies of Bulletin.....	8.00
Fee to inscribe Society as member of International Engineer- ing Congress.....	5.00
Total.....	\$171.86

Account of Bulletin

Express and freight charges.....	\$44.78
Postage	6.70
Messenger service.....	1.55
Printing	2.00
Jurat	.25
Collection charges on checks.....	1.29
Wrapping paper.....	2.73
Total.....	\$59.30
Total expenditures for the year.....	\$230.66

Respectfully submitted,

CHARLES P. BERKEY,
Secretary pro tem.

TREASURER'S REPORT

To the Council of the Geological Society of America:

The Treasurer herewith submits his annual report for the year ending November 30, 1915.

The membership of the Society at the present time is 376, of whom 284 pay annual dues. One member died early in 1914, but the notice

was not received until 1915. Nineteen new members were elected at the last annual meeting, all of whom qualified. There have been 4 deaths during the year and 1 resignation. Twenty-four members are delinquent in the payment of dues—1 for six years, 2 for four years, 3 for two years, and are therefore liable to be dropped from the roll—and 18 for one year.

One Life Member died during the year, which, with the 15 previous deaths, leaves 92 living Life Members.

With the advice of the Investment Committee, the Treasurer bought during the year one New England Telephone and Telegraph Company five per cent bond, with interest, at a cost of \$1,015; and two American Agricultural Chemical Company five per cent bonds, and interest, at a cost of \$2,040.28. One bond of the St. Louis, Iron Mountain and Southern Railway Company was redeemed on June 1.

RECEIPTS

Balance in treasury December 1, 1914.....	\$1,055.28
Fellowship fees, 1912 (1).....	\$10.00
1913 (1).....	10.00
1914 (8).....	80.00
1915 (261).....	2,610.00
	<hr/>
	2,710.00
Initiation fees (19).....	190.00
Interest on investments:	
Iowa Apartment House stock.....	50.00
Ontario Apartment House stock.....	200.00
Texas and Pacific Railroad Company	
bonds	100.00
U. S. Steel Corporation bonds.....	150.00
St. Louis, Iron Mountain and Southern	
Railway Company bond.....	25.00
St. Louis and San Francisco Railroad	
Company equipment bond.....	50.00
Fairmont and Clarksburg Traction Com-	
pany bonds.....	100.00
Consolidation Coal Company bonds.....	100.00
Chicago Railways Company bonds.....	100.00
Southern Bell Telephone and Telegraph	
Company bonds.....	100.00
New England Telephone and Telegraph	
Company bond.....	50.00
American Agricultural Chemical Company	
bonds	50.00
Interest on deposits, Baltimore Trust	
Company	48.53
	<hr/>
	1,123.53

Case Library, accessions 1914.....	150.00	
Redemption of St. Louis, Iron Mountain and Southern Railroad Company bond.....	1,000.00	
Collection charge added to checks.....	.50	
Received from Secretary:		
Sales of publications.....	\$1,078.00	
Authors' separates.....	88.20	
Authors' corrections.....	17.45	
Collection charges added to checks.....	.87	
Binding Bulletin.....	1.60	
Postage on foreign subscriptions.....	6.80	
	<hr/>	1,192.92
		<hr/>
		<u><u>\$7,422.23</u></u>
EXPENDITURES		
Secretary's office:		
Administration	\$171.36	
Bulletin	59.30	
Allowance	1,000.00	
	<hr/>	\$1,230.66
Treasurer's office:		
Postage, bond, safe-deposit box.....	\$40.00	
Allowance for clerical hire.....	100.00	
	<hr/>	140.00
Publication of Bulletin:		
Printing	\$2,437.23	
Engraving	185.44	
Editor's allowance.....	250.00	
	<hr/>	2,872.67
Purchase of one New England Telephone and Telegraph Company five per cent bond, and interest.....	1,015.00	
Purchase of two American Agricultural Chemical Com- pany five per cent bonds, and interest.....	2,040.28	
	<hr/>	7,298.61
Balance in Baltimore Trust Company December 1, 1915.....		123.62
		<hr/>
		<u><u>\$7,422.23</u></u>

Respectfully submitted,

WM. BULLOCK CLARK,
Treasurer.

EDITOR'S REPORT

To the Council of the Geological Society of America:

The Editor submits herewith his annual report. The following tables cover statistical data for the twenty-six volumes thus far issued:

Cost.	Average— Vols. 1-20.	Vol. 21.	Vol. 22.	Vol. 23.	Vol. 24.	Vol. 25.	Vol. 26.
	pp. 610. pls. 55.	pp. 839. pls. 54.	pp. 759. pls. 31.	pp. 774. pls. 43.	pp. 755. pls. 36.	pp. 820. pls. 28.	pp. 525. pls. 27.
Letter press.	\$1,686.58	\$2,049.95	\$1,660.45	\$1,750.40	\$1,647.90	\$2,049.19	\$1,076.22
Illustrations.	390.99	404.27	260.81	274.70	288.80	342.67	171.79
Total	\$2,077.57	\$2,454.22	\$1,921.26	\$2,025.10	\$1,936.70	\$2,391.86	\$1,248.01
Average per page.....	\$3.41	\$2.93	\$2.53	\$2.62	\$2.56	\$2.91	\$2.37

Classification.

Volume.	Areal geology.	Physical geol- ogy.	Glacial geology.	Physiographic geology.	Petrographic geology.	Stratigraphic geology.	Paleontologic geology.	Economic geol- ogy.	Official matter.	Memorials.	Unclassified.	Total.
Number of pages.												
1.....	116	137	92	18	83	44	47		60	4	4	593+xii
2.....	56	110	60	111	52	168	47	9	55	1	7	662+xiv
3.....	56	41	44	41	32	158	104		61	15	1	541+xii
4.....	25	124	38	74	52	52	14		47	32	2	458+xii
5.....	138	135	70	54	28	51	107		71	14	9	665+xii
6.....	50	111	75	39	71	99	1		63	25	4	538+x
7.....	38	77	105	53	40	21	123	4	66	28	13	558+x
8.....	34	50	98	5	43	67	58	14	79	8		446+x
9.....	2	102	138		44	28	64	16	64	12		460+x
10.....	35	33	96	37	59	62	68	28	84	27	17	534+xiii
11.....	65	110	21	10	54	31	188	7	71	60	46	651+xii
12.....	199	39	55	53	24	98	5	5	70	2		538+xi
13.....	125	17	13	24	28	116	42	4	165	32	29	583+xii
14.....	48	47	48	59	183	118	22	1	80	14	1	609+xi
15.....	26	124	3	94	36	267			77	17	3	636+x
16.....	64	111	78	30	102	141	19		67	22	15	636+xiii
17.....	49	161	41	84	47	294	27		71	9	2	785+xiv
18.....	16	164	141	5	29	246	5		68	40	3	717+xii
19.....	106	108	29	66	30	155	32		56	15	20	617+x
20.....	43	54	35	29	37	45	303	8	60	3	132	749+xiv
21.....	72	234	75	48	85	70	106	1	111	11	10	823+xvi
22.....	23	54	28	28	23	403	74		63	49	1	747+xii
23.....	75	52	126	108	19	145	134		66	32	1	758+xvi
24.....	18	57	96	57	49	160	106	23	133	53	3	737+xviii
25.....	34	211	54	32	156	9	175		108	9	22	802+xviii
26.....		72	23	11	56	90	148		54	44	6	501+xxi

Respectfully submitted,

JOSEPH STANLEY-BROWN, *Editor.*

The foregoing report is respectfully submitted.

THE COUNCIL.

December 28, 1915.

On motion, the report was laid on the table as usual until the following day.

ELECTION OF AUDITING COMMITTEE

The Auditing Committee, consisting of John E. Wolff, George H. Perkins, and E. B. Mathews, was then elected, and the Treasurer's report was referred to it for examination.

ELECTION OF OFFICERS

The Secretary declared the vote for officers for 1916 as follows, the ballots having been canvassed and counted by the Council in accordance with the By-Laws:

President:

JOHN M. CLARKE, Albany, New York.

First Vice-President:

J. P. IDDINGS, Brinklow, Maryland.

Second Vice-President:

HARRY FIELDING REID, Baltimore, Maryland.

Third Vice-President:

RUDOLPH RUEDEMANN, Albany, New York.

Secretary:

EDMUND OTIS HOVEY, New York City.

Treasurer:

WILLIAM BULLOCK CLARK, Baltimore, Maryland.

Editor:

JOSEPH STANLEY-BROWN, New York City.

Librarian:

FRANK R. VAN HORN, Cleveland, Ohio.

Councilors:

FRANK B. TAYLOR, Fort Wayne, Indiana.

CHARLES P. BERKEY, New York City.

ELECTION OF FELLOWS

The Secretary announced the election in due form of the following Fellows, the ballots having been canvassed and counted by the Council:

THOMAS CLACHAR BROWN, A. B., A. M., Ph. D., Bryn Mawr College, Bryn Mawr, Pennsylvania.

CHARLES WILFORD COOK, A. B., M. S., Ph. D., University of Michigan, Ann Arbor, Michigan.

WILLIAM EBENEZER FORD, Ph. B., Ph. D., Sheffield Scientific School, New Haven, Connecticut.

CHARLES TOWNSEND KIRK, B. S., A. M., Ph. D., University of New Mexico, Albuquerque, New Mexico.

DONALD FRANCIS MACDONALD, B. S., M. S., LL. D., United States Geological Survey, Washington, D. C.

EDGAR THEODORE WHERRY, B. S., Ph. D., United States National Museum, Washington, D. C.

Announcement was then made by the Secretary that the Society had lost four Fellows by death during the year 1915: Theodore B. Comstock, Orville A. Derby, Joseph A. Holmes, and William J. Sutton. Since the 1914 meeting the Secretary had also received notice of the death of Arthur B. Willmott on May 8, 1914. Memorials of deceased Fellows were presented as follows:

MEMORIAL OF THEODORE BRYANT COMSTOCK

BY HEINRICH RIES

Theodore B. Comstock was born at Cuyahoga Falls, Ohio, on July 27, 1849. After graduating from school he attended the Pennsylvania State College, where he received the Bachelor of Agriculture degree in 1868. In 1870 he obtained the Bachelor of Science degree from Cornell University, and in 1886 the Doctor of Science degree from the same institution.

His first teaching position was that of Instructor in Botany in Cornell University, which he held from 1868-1870, and then left to accept a professorship of Natural Science at Pelham Priory from 1871-1872. Following this he held the following positions: Instructor in Natural Science in Cincinnati, 1873; Director of the Kirkland Summer School of Natural History in 1875, and then acting Professor of Geology in Cornell University until 1879.

Many of his former students at Cornell speak highly of the interest which he took in them and their work, and it may be of interest to note in this connection that he gave the first instruction in Economic Geology that was given at this institution. His lecture syllabus which he pub-

Theo. B. Comstock.

lished at that time contains many illustrations showing the application of geology to engineering problems.

After leaving Cornell he occupied several other teaching positions as follows: Instructor in Shaler's Harvard Summer School of Geology, 1876; Professor of Mining Engineering and Physics, University of Illinois, 1885-1889; Director Arizona School of Mines, 1891-1893; President University of Arizona, 1893-1895.

During the period that he was engaged in teaching he also was occupied with more or less field work, beginning as early as 1870, in which year he went as assistant on the Morgan Expedition to Brazil. In 1876 he served as assistant on the Kentucky Geological Survey, and in 1877 as member of an expedition to the Northwest Territories, in 1877 as assistant on the Arkansas Geological Survey, and following that in a similar capacity on the Texas Geological Survey. In the later years of his life he gave up most of his active geological work and settled in Los Angeles, California, where he died on July 26, 1915.

Doctor Comstock was a member of the following societies: Geological Society of America, of which he was an original Fellow; American Institute of Mining Engineers, Mining and Metallurgical Society, American Association for the Advancement of Science, National Geographic Society, National Educators' Association, Southern California Academy of Science, and New York Academy of Science. He was the author of many papers dealing chiefly with geological subjects, a list of which is appended.

BIBLIOGRAPHY

- 1873. On the geology of western Wyoming. *American Journal of Science*, third series, volume 6, pages 426-432; volume 7, page 151.
- 1875. Geological report. Abstract. *American Journal of Science*, third series, volume 10, pages 59-60.
- 1876. Remarks on the hot springs and geysers and other topics illustrating the scientific value of the Yellowstone Park. *Proceedings of the American Association for the Advancement of Science*, volume 24, part 2, pages 97-99.
- 1876. Formation of geyserite pebbles in pools adjacent to the geysers of the Yellowstone Park. Abstract. *Proceedings of the American Association for the Advancement of Science*, volume 24, part 2, page 97.
- 1877. On some unexplained phenomena in the geyser basins of the Yellowstone National Park. *Proceedings of the American Association for the Advancement of Science*, volume 25, pages 235-239.
- 1883. Notes on the geology and mineralogy of San Juan County, Colorado. *Transactions of the American Institute of Mining Engineers*, volume 11, pages 165-191, map.
- 1886. Supermetamorphism and volcanism. *American Naturalist*, volume 20, pages 1006-1008.

1886. Some peculiarities of the local drift of the Rocky Mountains. *American Naturalist*, volume 20, pages 925-927.
1886. Remarkable extinct geyser basin in southwest Colorado. *American Naturalist*, volume 20, pages 963-965.
1886. The veins of southwest Colorado. *American Naturalist*, volume 20, pages 1043-1044.
1886. Mining engineering at the University of Illinois. *Transactions of the American Institute of Mining Engineers*, volume 15, page 589.
1887. Supermetamorphism; its actuality, inducing causes, and general effects. Abstract. *Proceedings of the American Association for the Advancement of Science*, volume 35, pages 232-233.
1887. Hints toward a theory of volcanism. Abstract. *Proceedings of the American Association for the Advancement of Science*, volume 35, page 233.
1887. The geology and vein structure of southwestern Colorado. *Transactions of the American Institute of Mining Engineers*, volume 15, pages 218-265, plates 1-4, map.
1887. The fossil fuels of Illinois and their exploitation. *Engineering and Mining Journal*, volume 44, page 24, quarto.
1887. Notes on the region north of the Vermillion Lake district in British Columbia. *Transactions of the American Institute of Mining Engineers*, volume 16, pages 109-111.
1887. Engineering relations of the Yellowstone Park. *Transactions of the American Institute of Mining Engineers*, volume 16, page 46.
1888. A preliminary examination of the geology of western central Arkansas. Report of the Arkansas Geological Survey for 1888, volume 1, pages 1-320, 2 maps.
1889. Hot Springs formations in Red Mountain district, Colorado: A reply to the criticisms of Mr. Emmons. *Transactions of the American Institute of Mining Engineers*, volume 17, pages 261-264.
1890. A preliminary report on the geology of the central mineral region of Texas. First Annual Report of the Texas Geological Survey, pages 237-391, plate 2.
1891. Report on the geography and mineral resources of the central mineral region of Texas, chiefly south of the San Saba River, north of the Pedernales River, west of Burnet, and east of Menardsville and Junction City. Second Annual Report of the Texas Geological Survey, pages 553-664, 3 maps.
1891. Tin in central Texas. *Engineering and Mining Journal*, volume 51, pages 117-118, quarto.
1891. A preliminary report on parts of counties of Menard, Concho, Tom Green, Sutton, Schleicher, Crockett, Valverde, Kinney, Maverick, Uvalde, Edwards, Bandera, Kerr, and Gillespie, Texas. Second Report of Progress of the Texas Geological Survey, pages 43-54.
1892. Valuable experiments in vein formation. *Science*, volume 19, page 214.
1894. Notes on Arizona mines. I. Silver. *Engineering and Mining Journal*, volume 57, page 103.
1895. Notes on Arizona geology. *Engineering and Mining Journal*, volume 60, page 369.

Yours truly,
Orville A. Derby

- 1900. The chloride district, Arizona. *Engineering and Mining Journal*, volume 70, pages 97-98.
- 1901. The geology and vein phenomena of Arizona. *Transactions of the American Institute of Mining Engineers*, volume 30, pages 1038-1101, 1 figure.
- 1902. Edward Claypole, the scientist. *American Geologist*, volume 29, pages 1-23, 1 plate.
- 1903. Memoir of Edward Waller Claypole. *Bulletin of the Geological Society of America*, volume 13, pages 487-497.
- 1905. Superficial blackening and discoloration of rocks, especially in desert regions. *Transactions of the American Institute of Mining Engineers*, volume 35, pages 1014-1017.
- 1902. Geological notes. *Bulletin of the Southern California Academy of Sciences*, volume 1, page 74.
- 1907. The United States Geological Survey. *Science*, new series, volume 25, page 309.

MEMORIAL OF ORVILLE A. DERBY

BY JOHN C. BRANNER

Orville Adelbert Derby was born at Kelloggsville, New York, July 23, 1851; and died by his own hand at Rio de Janeiro, Brazil, November 27, 1915. He was the third son of John C. Derby and Malvina A. Lindsay Derby, and was reared on a farm near Kelloggsville, in Cayuga County, in the "Finger Lakes" region of New York State, about 16 miles south-east of Auburn.

Derby entered Cornell University in 1869, and while yet a freshman he became so interested in geology and was such a promising student that he was selected by Prof. Charles Fred Hartt, then professor of geology at Cornell, to accompany him on a trip to Brazil in the summer of 1870. An incident showing a characteristic trait of the man had something to do with his selection for assistant on that trip and for his subsequent promotions. Professor Hartt had to leave the university for an absence of two weeks. He was a bit uncertain as to what could be done during his absence with this very new student of his. At a venture he gave him Hall's volume on the fossil bryozoa of New York—a work that would certainly have cooled the unguided ardor of most beginners. When Hartt came back at the end of two weeks Derby was patiently pegging away on the bryozoa. Hartt's heart warmed to a student who had the grit to stick to his uninspiring work, and shortly thereafter the opportunity to visit Brazil was given him. Derby gladly accepted the invitation, and in doing so he determined both his career and the whole course of his life.

On his first voyage to South America he went to Pernambuco and made the first considerable collections of fossils ever made at Maria Farinha.

In the summer of 1871 he went to Brazil with Hartt again, this time visiting the Amazon Valley and making an important collection of Carboniferous fossils from the limestones at Itaituba, on the lower Tapajós River.

In the interval between 1871 and 1873 he was occupied with his studies, and in 1873 he graduated at Cornell University; the year following he continued his geological work for the master's degree, which he received in June, 1874. His thesis was "On the Carboniferous brachiopoda of Itaituba, Rio Tapajós," and was published as number 2 of Volume I of the Bulletin of Cornell University, Ithaca, 1874. That was Derby's first publication on the geology of Brazil, and it is not only a valuable paper in itself, but it is especially interesting in view of subsequent developments. The Itaituba fossils were in compact limestone, but as they were silicified, they could be obtained in satisfactory form only by dissolving away the surrounding rock—a long and tedious process which would have thoroughly discouraged most young men of Derby's age.

In 1873 Derby was appointed instructor in geology in Cornell, and in the summer of 1874 Professor Hartt made arrangements to go to Brazil again. Leave of absence was obtained, Derby was placed in charge of the work of instruction in the department, and in September, 1874, Hartt went to Brazil again, taking Branner with him as his only assistant.

Arriving in Rio de Janeiro, Hartt at once devoted all his energies to interesting the leading men in a geological survey of the empire, and by the end of a year the survey was provided for, and O. A. Derby, Richard Rathbun, and E. F. Pacheco Jordão were named as assistants of the new "Comissão Geologica do Imperio do Brasil." In December, 1875, Derby reached Rio de Janeiro and began his work under the government. He held this position less than two years, for, through a change of ministry, the survey was abolished in 1877, and Hartt died in Rio that same year. Shortly after the extinction of the survey, however, Derby was given a position in the National Museum at Rio as curator in charge of geology. He remained in the museum until 1886, when he was made State Geologist of the Brazilian State of São Paulo.

The establishment of the São Paulo Survey was a step of great importance to geological science in Brazil, for Derby's knowledge of and interest in the geology of the country as a whole enabled him to grasp more firmly the geological problems of that particular State, and at the same time he became and remained the leading authority on the geology of Brazil. He was State Geologist of São Paulo until 1904, when he resigned.

In January, 1907, a new federal geological survey was provided for under Dr. Miguel Calmon, who was then minister of public works, and

Derby was made its chief—a position he held during the rest of his life. The appropriations were necessarily small at the outset, but the work undertaken was of great importance to Brazil, for sooner or later it was to point the way to an intelligent and scientific development of the natural resources of the country.

The first edition of Branner's *Geologia Elementar* was thus dedicated:

"To Orville A. Derby, who has devoted his life to the study of the geology of Brazil, and has done more than any one else to solve its many problems, this work is affectionately dedicated."

This is a brief and mild statement of Derby's great service to Brazil and to the science of geology, without mentioning his many other services to science and to that country.

Visitors to Brazil who were interested in geology always found him helpful in connection with their work. Dr. J. B. Woodworth, of Harvard, who went to Brazil in 1908 to study the Permian glaciation of the Southern States of that country, says that "without his aid and personal attendance it would not have been possible for me to have carried on the Shaler Memorial Expedition in Brazil to a successful issue in anything like the time in which that work was accomplished. With true Latin-American courtesy he paved the way for me to find what he would have been proud himself to have discovered—the actual occurrence of glaciated pebbles in the tillite beds of Paraná."

First and last, Derby was a paleontologist. He had no fondness for administrative work; he was but little interested in structural geology or in its methods; he was forced by circumstances into some acquaintance with microscopic petrography; but his interest in paleontology was genuine, deep, and all-comprehensive. From all the cares of office and the worriments of life he found relief and happiness in boxes of fragments of fossils that most paleontologists would have put away as not worth while.

It was chiefly to this interest of his in paleontology that we owe Dr. C. A. White's "Contributions to the Paleontology of Brazil," published at Rio in 1887, and the following valuable works by Dr. John M. Clark: "Trilobites of the Ereré and Maecurú sandstones," Rio, 1896; "Upper Silurian fauna of Rio Trombetas," Rio, 1899; "Devonian mollusks of the State of Pará," Rio, 1899, and "Devonian fossils of Paraná," Rio, 1913. Besides these more important publications there are many smaller papers on paleontology that can not be mentioned here, and there still remains unpublished an important volume by D. S. Jordan on the Cretaceous fossil fishes of Ceará.

During the last eight years Derby has given much of his time to the study of *Psaronius* and its relationships. The last of his published papers was on the stem structure of *Tietea singularis*, which appeared in the *American Journal of Science* for March, 1915, pages 251-260.

Having to undertake work in regions but poorly supplied with maps, one of his first and most important duties, when he became State Geologist of São Paulo, was the inauguration of topographic work. This work was intrusted to Horace E. Williams, an able and energetic young American, to whom the State of São Paulo and the scientific world are indebted for an excellent series of topographic maps, on a scale of 1 to 100,000, to say nothing of his explorations of the western portions of that State, his work on the Serra da Canastra, etcetera.

Derby's own list of publications on the geology of Brazil numbers 125 papers. Naturally they embrace a wide range of subjects. Ten of his papers relate to the geology and genesis of the Brazilian diamonds. He became interested in the early cartography of Brazil and published a number of papers on the subject.

As an author and as a scientific reasoner, he was an extremely cautious man, so much so that the word "hedge" was constantly on his lips, both for his own guidance and as a warning to his assistants.

The last evening I spent in his rooms at Rio de Janeiro he referred to this personal trait and remarked that it had prevented his marrying; that he was too cautious to take the risk. This cautiousness of his was probably the real reason for some of the long delays in publishing his results, which led to the tying up of his own results and those of his assistants. Without doubt he hoped the delays would enable him to put everything beyond question and to make his reports final and complete instead of preliminary and tentative. But the delays were prolonged from year to year, until his assistants became discouraged and the government more or less exasperated at the lack of practical results for such great and long continued expenditures. It was largely this long delay that finally led to his resignation as State Geologist of São Paulo.

Derby never felt obliged to show results. After he had been State Geologist of São Paulo for ten or twelve years and had published next to nothing on the geology of that State, I asked him point blank and with some feeling where his results were. He replied: "They are in my head." We had to change the subject. But the important fact behind his delays is that the geology of São Paulo is difficult and involves problems that he had not been able to settle to his own satisfaction, and he was unwilling to commit himself to paper and thus lay himself open to adverse criticism.

It seemed unfortunate for Brazil, for himself, and for the cause of

science that he was unable to bring himself to take an active interest in the economic geology of the country; but his first and only interest in geology was in geology as a pure science. To him a fossil was a thing of beauty, of interest, and value, and a joy forever; but a mine or an industry was, after all, only an industry whose main object was money-getting.

It goes without saying that Derby and I did not always agree about geological questions, but our very disagreements tended to stimulate careful work and finally to disclose the truth. An interesting illustration was our disagreement in regard to certain beds in the black diamond regions of Bahia; he called them Paraguassú and I called them Caboclo. After a year of proving each other in the wrong, he was induced to send his assistant, Roderic Crandall, to the region in question to settle the dispute. Crandall went and reported that we were not speaking of the same things; that both series were legitimate, and that we were both right.

Derby was a man of unlimited gift. When once he decided on a course of action, nothing turned him to the right or to the left. His whole life is a demonstration of his power to make good in spite of obstacles that would have been insurmountable for most men—his determination to devote his life to the geology of Brazil, cost what it might.

How many of us would have lived for forty years in a foreign country, cut off, as he was, from all personal contact with the geologists of the world at large and from the people of his own race and from his own family? From the time he went to Rio, in 1875, to the day of his death—a space of forty years—he visited the United States only twice. The first of these visits was from January to June of 1883, when he went to Washington to arrange for the publication of Dr. C. A. White's Contributions to the Paleontology of Brazil, an epoch-making work on South American paleontology. He spent part of that time in Boston, New Haven, New York, and Philadelphia. His other visit was made in 1890, when he attended the Indianapolis meeting of the American Association, and returned to Rio by way of England.

When the Comissão Geologica was stopped, in 1877, the rest of us took to our heels. Not so Derby; he was not to be stampeded by a simple lack of funds or of employment; he meant to save the results of the work of Hartt and of his colleagues, and, so far as it could be done, he did it.

Personally Derby was one of the kindest hearted and most affectionate men I have ever known. His time, his sympathies, and his last dollar were at the service of his friends; and his right hand knew nothing of the kind deeds done by his left. The beggars in the streets found him their easiest victim.

He was held in the highest esteem in the community in which he lived. He stood for uprightness and honorable dealing, and he was never the willing tool of designing adventurers. For many years he has been justly regarded as the leading geologist in South America, and his standing is due not to the fact that there are but few geologists in South America, but to his ability and to his excellent work.

In 1892 he was awarded the Wollaston prize of the Geological Society of London, while his distinguished services led to his being made one of the associate editors of the *Journal of Geology* and to his election to membership in various learned societies in different parts of the world. He was a frequent contributor to the *American Journal of Science*. He was naturalized as a Brazilian citizen a few months before his death.*

The circumstances that led to Derby's suicide are not clearly known, or rather they are not clearly understood. Neither the published accounts of all the details available to the authorities nor the many private letters received from mutual friends throw much light on the case. The act was committed in his rooms in the Stranger's Hotel in Rio, where he had lived for eight years. The evening preceding his death was spent at the home of a Brazilian friend and he went to his own rooms about midnight. The next morning he was called as usual, took his bath, drank his coffee, and read the morning papers. About 10 o'clock a messenger going to his room found him lying across his bed with a bullet-hole through his head and the revolver still grasped in his dead hand. He left no word of explanation or complaint about anything or against any one.

The general impression seems to be that his suicide was due to disappointment on account of the reduction by the government of the appropriations for his work. A few feeble efforts have been made to find other explanations for his act, but this must be and is accepted as the only genuine one.

The history of his struggles to keep the scientific work intrusted to him out of politics and to make it efficient is nothing new. Scientific men the world over have often been in similar positions. In the present case there is no doubt but that the matter was complicated by the financial situation in Brazil brought about by the war in Europe. The government was hard pressed financially and it was absolutely necessary to reduce expenses to the lowest possible point. It was not unnatural under

* His successor as director of the Brazilian Survey is Dr. J. F. Gonzaga de Campos, who has been one of the geologists of the survey since it was begun. Doctor Campos is the author of a number of valuable papers on Brazilian Geology and a thoroughly conscientious man of wide personal acquaintance with the geology of Brazil.

the circumstances that the geological service should have been selected as the one that could be reduced without producing serious confusion in the administration of the government.

The Brazilian newspapers all speak of him in the highest possible terms, and his death is generally and properly looked on in Brazil as a great national loss.

His suicide itself was his last and his most emphatic protest against the extinction of geologic work—the final culminating expression of his profound interest in and devotion to the welfare of the country he had served faithfully for forty years.

A list of his papers on the geology of Brazil up to 1909 is given in this Bulletin, volume 20, papers 36 to 42. To that list should be added the following titles, that have appeared since the publication of that list:

BIBLIOGRAPHY

- Feições physicas e geograficas do Brasil. Boletino da Directoria da Agricultura da Bahia, volume X, pages 241-248. Bahia, 1907.
- Serviço Geologico e Mineralogico do Brasil. Boletino do Ministerio da Industria, Viação e Obras Publicas, volume I, pages 69-82. Rio, Abril de 1909.
- Os minerios de ferro do Brasil. Jornal do Commercio, Rio, August 25, 1909.
- Early iron-making in Brazil. Engineering and Mining Journal, New York, December 4, 1909.
- The iron ores of Brazil. The Times, London, December 28, 1909, page 56.
- The iron-ore resources of the world. Stockholm, 1910, pages 813-822.
- Physical and geological features of Brazil. The Brazilian Yearbook for 1909, pages 11-14. Rio de Janeiro n. d.
- Estudios geologicos en el Brasil. Santiago de Chile, 1911. (Publication of the fourth congresso científico Latino Americano en 1908.)
- On the mineralization of the gold-bearing lode of Passagem, Minas Geraes. Brazil. American Journal of Science, volume CLXXXII, September, 1911, pages 185-190.
- A notable Brazilian diamond. American Journal of Science, volume CLXXXII, September, 1911, pages 191-194.
- O aproveitamento do carvão brasileiro. Jornal do Commercio, 24 de Abril de 1912, page 5.
- Speculations regarding the genesis of the diamond. Journal of Geology, volume XX, July-August, 1912, pages 451-456.
- Observations on the stem structure *Psaronius brasiliensis*. American Journal of Science, November, 1913, pages 489-497.
- Observations on the crown structure of *Psaronius brasiliensis*. American Journal of Science, August, 1914, pages 149-156.
- Illustrations of the stem structure of *Tietea singularis*. American Journal of Science, volume XXXIX, March, 1915, pages 251-260.

MEMORIAL OF JOSEPH AUSTIN HOLMES¹

BY JOSEPH HYDE PRATT

The life of Dr. Joseph Austin Holmes was devoted to the development and welfare of his country, and in his death the people of the United States have lost one of their most efficient and valuable public servants. He was a man who put duty first, and in carrying out this ideal he gave his life in an endeavor to improve the condition and safety of the miners. He did not know the word "failure;" and, where other men would have failed, he has been able to accomplish the results desired. It is granted to but few men to be able in the few years of their life's activity to do that which will leave a permanent influence and impress on an industry; but to Doctor Holmes, whose life we are now commemorating, this distinction was allotted.

Due almost entirely to his energy and efforts, there has been created throughout this country an organized movement looking to the preservation of human life; and, although his first work was directed toward the prevention of mine accidents and safety and welfare of the hundreds of thousands of men who daily risk their lives in the production of fuel, so necessary to the nation's industry and commerce, it developed the "safety first" idea that has spread to nearly every industry and into all walks of life. These words are almost synonymous with the word "Holmes," and wherever we see "safety first" we are reminded of the wonderful achievements of this man. He has not only left his impress on an industry, but has also created an organization which will live as long as our government exists, and is a monument to the tireless energy, public-spiritedness, and unselfishness of the man who is responsible for its creation. I refer to the Bureau of Mines, whose foundation he laid by many feats of exacting labor and fruitful work, and who, by masterful generalship and arguments, as he only could use, carried the bill to establish the Bureau of Mines successfully through an unsympathetic Congress.

To Dr. Charles D. Walcott, former Director of the United States Geological Survey and now Secretary of the Smithsonian Institution, must be given the credit of recognizing those qualities of character and ability in Mr. Holmes which he realized were necessary in a man who could not only lay the foundation and build up an organization that would lead to a Bureau of Mines, but who would also be able to direct it after its creation. In a recent communication from Doctor Walcott, he wrote:

¹ Read at the annual meeting of the Geological Society of America, Washington, D. C., December 28, 1915.

J. Holmes

"About 1900 it became more and more evident that he (Doctor Holmes) was a man of broad conceptions and fitted to undertake work of national scope, and it was with great pleasure that I learned, in 1904, that he was willing to give all of his time and energy to the development of the Section of Mines and Mining in the Federal Survey. I told him that as soon as the work was sufficiently well organized it would be made a Division of the Survey and undoubtedly lead to the creation of a Bureau of Mines and Mining. He entered into the work with a zeal and intelligence that was not fully understood by his immediate associates; but the work steadily grew and, in 1910, he was appointed Director of the Bureau of Mines."

His appointment, however, was not attained without very severe opposition from a Secretary who was hostile to Doctor Holmes, and it is rumored that this important position was offered to several other men; but, to the credit of the men of science of this country, it can be said that they all refused to accept what all knew rightfully belonged to another. Those who knew Doctor Holmes, having confidence in his ability and believing that he was the logical head for the new Bureau, were persistent in their demand that he should receive the appointment. It is not generally known how near the Bureau came to losing Doctor Holmes as its Director and how near the University of West Virginia came to securing him as its president; and, as an incident bearing on this is illustrative of the loyalty of Doctor Holmes' friends, I wish to quote in part a few lines from a letter I recently received from Dr. I. C. White, State Geologist of West Virginia:

"It was during this discouraging period of his life, just before the appointment of a Director of the Bureau of Mines, when he had given up all hope of receiving the appointment, that he came up from Pittsburgh to spend the week end at the writer's home in Morgantown, West Virginia. He was weary and care-worn from the long and disappointing vigil, but gentle and loving as ever. No word of reproach or bitterness escaped his lips. If he could not serve his country in an edifice his own hands had so largely constructed, he was ready to give his services to a State that had stood by him in his long battle, and where he knew he would be among appreciative friends. The State University of West Virginia was seeking a president, and one of the purposes of Doctor Holmes' visit to my home was to acquaint the writer, who had ever been his trusted friend, with the fact that he had despaired of being appointed Director of the United States Bureau of Mines, and would accept the presidency of the University of West Virginia if the regents of the same would make the tender."

Fortunately for the industry, Doctor White and others, realizing that for the success of the Bureau of Mines it was necessary that Doctor Holmes should be its head, decided, out of genuine loyalty to him and appreciation of his work, that they would not place his name for action before the regents of the University until President Taft had actually

bestowed the directorship of the Bureau of Mines on some one else. His friends' belief in what President Taft would finally do was confirmed a few days' later, when the appointment of Doctor Holmes was announced from the White House.

That he was a wise selection is evidenced by the wonderful development of the Bureau under his administration. The work he had planned as Chief of the Technologic Branch of the United States Geological Survey developed rapidly, aided by Congress, which widened the scope and enlarged the purposes of the Bureau. The principal investigations taken up under Doctor Holmes' directorship and the results accomplished are as follows:

An investigation in regard to the improper use of explosives and the use of improper explosives.

Investigation in regard to better lights for mines. Result, the establishment of a permissible list of portable electric lamps for use in dangerous mines.

In developing rescue work Doctor Holmes introduced into this country the so-called "oxygen breathing apparatus." Result, such apparatus is now not only widely used in mine-rescue work, but is being adopted by manufacturing plants and by city fire departments. There are today six mine-rescue stations, eight mine-rescue cars, and one rescue motor truck operated by the Bureau of Mines. There are 76 mine-rescue stations that have been established by mining companies, at which there are 1,200 sets of artificial breathing apparatus in addition to the auxiliary equipment for first-aid and fire-fighting work. There are also twelve mine-rescue cars being operated by mining companies.

Investigations into the cause of disasters and the recommendations made by the Bureau have resulted in an ever decreasing death rate.

The investigation of coal dust and explosions therefrom was one of the most important lines of investigation that Doctor Holmes took up. The result today is that the entire mining industry, including operators and miners, is convinced that coal dust will explode, and recognize the danger from it; and mine operators and State officials are following the recommendations of the Bureau to prevent dust explosions.

Investigations have been conducted regarding smelter smoke wastes and wastes in the treatment of rare minerals and metals. Doctor Holmes emphasized the need of such investigations, indicating that there was at least \$1,000,000 a day being wasted or lost in the present methods of mining and utilization of our mineral resources.

Investigations regarding the extraction of radium from its ores have resulted in the development of a process through which it will be possible

to greatly reduce the cost of radium compounds to the consumer. "The process is to be patented and dedicated to the public."²

Investigations have been started to reduce the great loss of \$75,000,000 annually, due to coking coal in beehive ovens. As a result already some of this loss has been reduced through the use of by-product ovens and the utilization of the by-products obtained.

Doctor Holmes called attention to the annual waste of over \$4,500,000 in brass-furnace practice, and then had prepared a report showing how, by practical means, this waste can be largely prevented.

These are some of the investigations that Doctor Holmes has had taken up by the Bureau of Mines, and they illustrate the wide scope of the work he was planning for the Bureau to undertake. Its development into one of the most important of all the Federal Bureaus has been phenomenal and is due not only to the indefatigable work of the Director, but to the fact that he was a splendid judge of men and their capacity for work and was able to surround himself with the type of men who were able to carry out the plans his master mind had conceived, and these men were loyal and true to him.

He was thoughtful and considerate of his associates; and while he may have demanded much of them, he always gave them full credit for work done; and of the reports of the investigations carried out by the Bureau but very few bear his name as author. Credit is given to him who carried on the investigation. Doctor Holmes planned the character of the investigation, then put it up to one of his associates to do the detailed work. What he wanted was results. He had little time to write for publication or to think about personal advancement, and he left it to his associates to do the writing and give him the results—and results he surely obtained.

Although Doctor Holmes is the author of but comparatively few publications, yet he has been personally responsible for the publication of many important scientific and economic papers, because he has had the foresight to open up new fields of investigation and secure the properly trained men to carry on the work he outlined. I doubt if there has ever been a man who surpassed him in this respect.

This faculty of Doctor Holmes showed itself soon after he became State Geologist of North Carolina in 1891. In this position he had wide latitude for planning out a varied line of investigations relating to many subjects, inasmuch as the object of the State Survey was the investigation of all natural resources of the State. Almost as soon as he was appointed State Geologist, he began to plan new lines of work and to call in to assist

² Van H. Manning: Jour. Ind. and Eng. Chem., vol. 7, No. 8, p. 716, Aug., 1915.

him men who were fully qualified to carry on the investigation he desired. Thus you find associated with him during the first years of his directorship of the State Survey such men as Prof. George Williams, of Johns Hopkins; Prof. S. L. Penfield, of Yale; Dr. George F. Kunz, of New York; Prof. F. P. Venable, of the University of North Carolina; Dr. George P. Merrill, of the National Museum at Washington; Prof. George Swain, of the Massachusetts School of Technology; Prof. Thomas L. Watson, of the University of Virginia; Prof. H. V. Wilson, of the University of North Carolina; Prof. William Cain, of the University of North Carolina; Mr. H. B. C. Nitze, Mr. Gifford Pinchot, of Washington; Prof. Heinrich Ries, of Cornell, and others. The published reports of the State Survey, similarly as those of the Bureau of Mines, seldom bear the name of Holmes as one of the authors.

Doctor Holmes did a great deal to broaden the scope of the State Geological surveys and to demonstrate that there could be and should be a close cooperation between the State and Federal surveys. There was always most friendly cooperation between the North Carolina Survey and the Federal Survey; and, although the State received very largely from the Federal Survey, it gave very largely in return, for Doctor Holmes was always ready to give his time and energy to any work which promised to be of service to the Federal Survey; and he was often called in consultation regarding the work of that Survey. Doctor Walcott, who was then Director, states that he was early impressed with Mr. Holmes' thoroughness and the quality of his work as State Geologist. In the Geological Survey his most important work was probably the application of geology to the industrial development of the country. He started this in the State Survey, but later introduced it into the Federal Survey.

As State Geologist he became very much interested in the preservation of the forests of the southern Appalachian region, and it is due largely to his work as State Geologist that the Weeks bill was passed by Congress, which has resulted in the purchase of forest areas to be used for forest reservations in the southern Appalachian region and the White Mountain region. It was under the supervision of Doctor Holmes that the mass of evidence was collected, which proved to the congressional committees that it was absolutely necessary for Congress to take some action to prevent the destruction of the forests of these two areas in order to protect the flow of navigable streams.

In connection with an investigation relating to our turpentine industry, he had experimental work carried on in regard to the cup and gutter method, which is now in general use in this industry. He also had investigations made as to the practicability of the reproduction of the long-

leaf pine, and an actual demonstration in planting of seed proved the feasibility of such reproduction.

Doctor Holmes also started the "good roads" movement in North Carolina, and one of the first publications of the State Survey was a report on "Road materials and road construction in North Carolina." While his work in connection with the roads was almost entirely from the educational standpoint, yet it was this work that made it possible for his successor to obtain, through the North Carolina General Assembly, the creation of, first, a Highway Division of the Survey, and later of a State Highway Commission.

In the State work Doctor Holmes also began investigations in relation to the water powers, mineral waters, underground water supplies, timber resources, mineral resources, and fisheries of the State; but a limited treasury and lack of time prevented him from carrying these out as rapidly as he desired, and it was left to his successor to complete some of them.

During his term of office as State Geologist, 1891 to 1905, the Survey published twenty bulletins and economic papers, giving the results of investigations that he had started. In 1905 the act creating the Survey was repealed and a new act, which was prepared by Doctor Holmes, was passed by the General Assembly of 1905. This created the North Carolina Geological and Economic Survey.

Doctor Holmes brought geology to this people and made them realize its value and application in the arts.

In connection with the investigation of the fisheries of the State, Doctor Holmes was the leading spirit in the establishment of the Biological Laboratory at Beaufort. In June, 1897, after consultation with Doctor Holmes and Prof. H. V. Wilson, of the University of North Carolina, the United States Commissioner of Fisheries established at Beaufort, North Carolina, a temporary station for the investigation of the marine fauna and flora of the Southern coast. Professor Wilson was appointed director, and for the next three years he and Doctor Holmes devoted much time and thought to its development. Congress finally made an appropriation for the establishment of a permanent laboratory, but made no appropriation for the purchase of a site. Doctor Holmes recommended a site and arranged for its private purchase and its donation to the government. He, with Professor Wilson, drew up the outline plans for the laboratory buildings, and he remained in close touch with the work of the laboratory until his resignation as State Geologist. This work of Doctor Holmes had an important bearing on the fisheries of the State of North Carolina, as it started the interest of the people of the State in the value of the

fisheries, and finally resulted, some years after the resignation of Doctor Holmes as State Geologist, in the creation of the Fisheries Commission of the State of North Carolina.

Doctor Holmes' work as State Geologist had brought him prominently before the public, and in 1903 he was chosen Director of the Department of Mines and Metallurgy at the Saint Louis World's Fair. He accepted this appointment and had charge of and organized that department. He planned the exhibits and introduced new features for the exhibits, which have since been adopted by all succeeding expositions. These new features made the Mine Building of the Saint Louis Exposition the most successful and instructive mining exhibit that was ever made at any exposition. For special services rendered at this exposition he was decorated by several foreign governments. In connection with the mining exhibit, he suggested that an investigation be made of the fuels of the United States, and was successful in persuading Congress to authorize the investigation and make the necessary appropriation with which to carry on the work. Doctor Holmes and, at his suggestion, two representatives of the United States Geological Survey were created a committee to carry on the investigations which were made during the years 1904 and 1905. Although Director of the Department of Mines at Saint Louis, Doctor Holmes continued to have general supervision of the work of the North Carolina Geological Survey. Early in 1905 the Director of the United States Geological Survey appointed Doctor Holmes to take individual charge of the fuel investigations, and soon after he was appointed Chief of the Division of Technology of the Federal Survey, and then severed his connection with the State Geological Survey.

While connected with the Federal Survey, Doctor Holmes examined mine experiment stations and mine-rescue stations in Great Britain, Belgium, France, and Germany, and it was the result of these studies that led to the inauguration of the movement for mine-rescue work in this country.

In 1907 President Roosevelt, on Doctor Holmes' recommendation, secured the appointment by the governments of Great Britain, Germany, and Belgium of one expert engineer from each of these countries to visit the United States and then visit, with Doctor Holmes, the more important coal fields of this country. This was done in order to determine to what extent the safety practices used in other mining countries might be introduced into the United States. It was on the basis of the findings of these engineers that Doctor Holmes developed and organized his investigations relating to mine explosions, etcetera.

In 1908, when President Roosevelt took up the question of the conservation of our natural resources, Doctor Holmes was appointed a member of the National Conservation Commission, and he had charge of the inventory of the nation's mineral resources.

In all Doctor Holmes' work his central thought has always been the development of the mining industry and the improvement of conditions affecting the miner. In carrying out these great ideas, he thought only of the object to be attained and paid little or no heed to personal attacks or opposition, such as inevitably accompanies a forward movement or investigation that requires the cooperation of both the legislative and administrative departments of our government. When, however, an attack was made on him that appeared to endanger the work itself in which he was engaged, he was then ready to put forth all his efforts to meet and defeat the opposition.

Doctor Holmes was human as the rest of us and occasionally was forgetful in regard to certain things that were to be done. This characteristic of his sometimes led to severe criticism of his work by those who were not thoroughly acquainted with him. Whenever any apparent neglect on his part was called to his attention, the matter was instantly taken care of and ample apology made for the oversight. Doctor Holmes was excessively careful to observe all the little courtesies of life and was a splendid representative of the Southern Christian gentleman.

Doctor Holmes was born at Laurens, South Carolina, November 23, 1859, and died at Denver, Colorado, July, 1915, after nearly a year's illness and fight against tuberculosis. His ill health was undoubtedly brought on by severe exposure in connection with the examination of mines after explosions and of hardships endured in investigations regarding mining conditions in Alaska. His parents were Z. L. and Catherine (Nickles) Holmes.

His early education was in the schools of South Carolina, but his university work was at Cornell, where he graduated in 1881, taking the degree of B. S. Later he received the degree of D. Sc. from the University of Pittsburgh, and in 1909 the degree of LL. D. from the University of North Carolina. During his college course Doctor Holmes devoted especial attention to chemistry (including the chemistry of explosives), to metallurgy, geology, general physics, and mining. He visited mining regions and metallurgical plants in many parts of the United States, Germany, France, Great Britain, and Belgium.

In the fall of 1881 he became professor of Geology and Natural History in the University of North Carolina, and held this position until 1891, when he became State Geologist.

On October 20, 1887, Doctor Holmes married Miss Jeannie I. Sprunt, of Wilmington, North Carolina.

Doctor Holmes was a fellow and charter member of the Geological Society of America; fellow of the American Association for the Advancement of Science; member of the American Institute of Mining Engineers, American Society for Testing Materials, and American Society of Mechanical Engineers. He was appointed a member of the Mining Legislation Committee of Illinois; one of the founders of the Elisha Mitchell Scientific Society; member of the Sigma Xi Scientific Society; member of the Washington Academy of Science, Saint Louis Academy of Science, and the North Carolina Academy of Science; member of the Cosmos Club of Washington and the Engineers' Club of New York.

In closing this sketch let me further express my feelings and thought regarding Doctor Holmes in the words of several friends who were very close to him:

"Doctor Holmes stands as one of the finest examples of unselfish devotion to the cause which he championed, even to the extent of giving his life for it. Mining in America in its national aspect is more deeply indebted to him on its scientific, operating, and industrial sides than to any one other individual. It seems most unfortunate that Doctor Holmes did not live to aid the movement to improve the laws affecting mines and mining; but, with the Bureau of Mines firmly established, and cooperating with the thoughtful mining engineers and operators throughout the country, the results he hoped to see should be speedily obtained.

"CHARLES D. WALCOTT."

"Ever thoughtful, resourceful, a great organizer, a clear, logical, and eloquent speaker, a splendid judge of men and their capacity to do the work his master mind had planned, the United States Bureau of Mines, founded only in 1910, has under his leadership rapidly grown to be one of the most important of all government agencies. . . .

"His monument is the United States Bureau of Mines, and his memory will be cherished forever in the hearts of countless miners, whose lives he has rendered safer in the perilous occupation they follow, and without the product of whose busy hands our present civilization could not exist. Although cut down in but little beyond the prime of life, he has left us an example of what glorious achievements, indomitable will, and untiring work can accomplish. The great Bureau he so largely created and so successfully directed will continue its brilliant work along the path he so skilfully blazed, since, thanks to a very able and conscientious Secretary of the Interior, his successor is in thorough accord with the high ideals of the former chief, and was ever his efficient helper.

"I. C. WHITE."

"In the death of Doctor Holmes the people of the United States lose one of their most remarkable and efficient public servants. And the saddest part of it all is that Doctor Holmes is a victim of overwork—a too great devotion to

the duties which had been assigned to him in behalf of the safety of the million miners in the United States. He was one of the most enthusiastic, indefatigable workers I ever had the pleasure of associating with. His mind was continually on the yearly death toll of the miners, and, although taken away in the prime of his life, he has already accomplished much in reducing the terrible death rate. In the last five years of his life he saw a slowly but steadily decreasing death rate, and while it gave him much joy, it only added to his almost superhuman efforts in behalf of the men.

"VAN. H. MANNING."

A full list of Doctor Holmes' reports and more important scientific papers is given in his bibliography, which appears at the close of this sketch. The picture of Doctor Holmes that accompanies this sketch was selected by Mrs. Holmes.

BIBLIOGRAPHY

1. Agricultural education in North Carolina. Miscellaneous Special Report Number 2, United States Department of Agriculture, 1883, pages 84-87.
2. Notes on the tornado which occurred in Richmond County, North Carolina, February 19, 1884. Journal of the Elisha Mitchell Scientific Society, volume I, 1884, pages 28-34.
3. Notes on the Indian burial mounds of eastern North Carolina. Journal of the Elisha Mitchell Scientific Society, volume I, 1884, pages 73-79.
4. Occurrence of *Abies canadensis* and *Pinus strobus* in central North Carolina. Journal of the Elisha Mitchell Scientific Society, volume I, 1884, pages 86-87.
5. Notes on a petrified human body. Journal of the Elisha Mitchell Scientific Society, volume II, 1885, pages 59-60. (With Dr. T. W. Harris.)
6. Taxodium (cypress) in North Carolina. Journal of the Elisha Mitchell Scientific Society, volume II, 1885, pages 92-93.
7. Supplemental report on Sam Christian gold mine. Manuscript. North Carolina Geological Survey, 1886, 3 pages.
8. A sketch of Prof. Washington Caruthers Kerr, M. A., Ph. D. Journal of the Elisha Mitchell Scientific Society, volume IV, part 2, 1887, pages 1-24.
9. Temperature and rainfall at various stations in North Carolina. Journal of the Elisha Mitchell Scientific Society, volume V, 1888, pages 31-41.
10. Study of plants in the garden and field. The North Carolina Teacher, 1888, 6 pages.
11. Historical notes concerning the North Carolina Geological Surveys. Journal of the Elisha Mitchell Scientific Society, volume VI, 1889, pages 5-18.
12. The conglomerate and pebble beds of the Triassic and Potomac formations of North Carolina. Journal of the Elisha Mitchell Scientific Society, volume VI, 1889, page 148.
13. Mineralogical, Geological, and Agricultural Surveys of South Carolina. Journal of the Elisha Mitchell Scientific Society, volume VII, 1890, pages 89-117.

14. Hoover Hill gold mine in North Carolina. *Engineering and Mining Journal*, volume LIV, page 520.
15. Character and distribution of road materials. *Journal of the Elisha Mitchell Scientific Society*, volume IX, part 2, 1892, pages 66-81.
16. Road material and road construction in North Carolina. (With William C. Cain.) *Bulletin 4, North Carolina Geological Survey*, 1893, 88 pages.
17. Geology of the sand-hill country of the Carolinas. *Bulletin of the Geological Society of America*, volume 5, 1893, pages 33-34.
18. Economic geology of North Carolina. *Southern States*, volume I, 1893, pages 153-161.
19. Improvement of roads in North Carolina. *Yearbook*, 1894, United States Department of Agriculture, 1895, pages 513-520.
20. Notes on the kaolin and clay deposits of North Carolina. *Transactions of the American Institute of Mining Engineers*, volume XXV, 1895, pages 929-936, and *Journal of the Elisha Mitchell Scientific Society*, volume XII, part 2, 1895, pages 1-10.
21. Notes on the underground supplies of potable waters in the South Atlantic Piedmont Plateau. *Transactions of the American Institute of Mining Engineers*, volume XXV, pages 936-943, and *Journal of the Elisha Mitchell Scientific Society*, volume XII, part 1, 1895, pages 31-41.
22. Corundum deposits of the southern Appalachian region. *Seventeenth Annual Report of the United States Geological Survey*, part 3, 1896, pages 935-943.
23. Gold in the Carolinas. Gold fields along the Southern Railway. Published by the Southern Railway, 1897, pages 8-19.
24. Mica deposits of the United States. *Bulletin of the Geological Society of America*, volume 10, 1898, pages 501-503.
25. North Carolina mineral industry in 1898. *Engineering and Mining Journal*, volume LXVII, 1899, pages 50-51.
26. Mica deposits in the United States. *Twentieth Annual Report of the United States Geological Survey*, 1899, pages 691-707.
27. Water Power in North Carolina. (With Geo. F. Swain and E. W. Myers.) *Bulletin 8, North Carolina Geological Survey*, 1899, 362 pages.
28. Some recent road legislation in North Carolina. *Economic Paper Number 2, North Carolina Geological Survey*, 1899, 24 pages.
29. The deep well at Wilmington, North Carolina. *Journal of the Elisha Mitchell Scientific Society*, volume XVI, part 2, 1899, pages 67-70; *Science*, new series, volume XI, 1900, page 128.
30. Mica industry in North Carolina in 1900. *United States Geological Survey. Mineral Resources*, 1900, pages 853-954.
31. The Cretaceous and Tertiary section between Cape Fear and Fayetteville, North Carolina. *Science*, new series, volume XI, 1900, page 143.
32. Recent road legislation in North Carolina. *North Carolina Geological Survey. Economic Paper Number 5*, 1901, 47 pages.
33. Proceedings of the North Carolina Good Roads Convention. *United States Department of Agriculture, Office of Public Road Inquiries. Bulletin Number 24*, 1903, 72 pages.
34. Road building in North Carolina. *United States Department of Agriculture, Office of Public Road Inquiries. Bulletin Number 24*, 1903, pages 65-71.

- 35-41. Biennian reports of the North Carolina Geological Survey; 1891-1892; 1893-1894; 1895-1896; 1897-1898; 1899-1900; 1901-1902; 1903-1904.
42. The collection of mineral statistics in the United States of America. Cong. int. d'expansion econ. mondiale, Mons, 1905, sect. 2, Statist. int. Bruxelles, 2 pages.
43. Fuel investigations, Geological Survey; progress during year ending June 30, 1909. Proceedings of the American Society for Testing Materials, volume 9, 1909, pages 619-625.
44. Inspection of mines. Report of proceedings of the American Mining Congress, twelfth annual session, Goldfield, Nevada, September 27-October 2, 1909, pages 236-238.
45. Preliminary report of Committee on Standard Specifications for Coal. Proceedings of the American Society for Testing Materials, volume 9, 1909, pages 277-279.
46. A rational basis for the conservation of mineral resources. Bulletin 29, American Institute of Mining Engineers, May, 1909, pages 469-476.
47. Coal-mine accidents and their prevention. National Civic Federation circular, New York, November 23, 1909, 4 pages.
48. The Bureau of Mines and its work. Report of proceedings of the American Mining Congress, thirteenth annual meeting, Los Angeles, California, September 26-October 1, 1910, pages 219-227.
- 49-53. Annual Report of the United States Bureau of Mines, 1911-1915, 5 volumes.
54. The sampling of coal in the mine. Technical Paper 1, United States Bureau of Mines, 1911, 18 pages.
55. The mining industry. Report of proceedings of the American Mining Congress, fourteenth annual meeting, Chicago, Illinois, October 24-28, 1911, pages 69-71.
56. Diseases and accidents of miners and tunnel workers in the United States. Reprint from Transactions of the Fifteenth International Congress on Hygiene and Demography, September 23-28, 1912, 13 pages.
57. Saving miners' lives. Proceedings of the Fourth National Conservation Congress, Indianapolis, October 1-4, 1912, pages 200-205.
58. The national phases of the mining industry. Eighth International Congress of Applied Chemistry, volume 26, 1912, pages 733-750.
59. Speech concerning work of the Bureau of Mines. Report of proceedings of the American Mining Congress, seventeenth annual meeting, Phoenix, Arizona, December 7-11, 1914, pages 95-96.
60. Preliminary report on operations of coal-testing plant of United States Geological Survey at Louisiana Purchase Exposition, Saint Louis, Missouri, 1904. E. W. Parker, J. A. Holmes, and M. R. Campbell. Bulletin 263, United States Geological Survey, 1905, 172 pages.
61. Report on operations of coal-testing plant of United States Geological Survey at Louisiana Purchase Exposition, Saint Louis, Missouri, 1904. Professional Paper 48, United States Geological Survey, 1906, 3 volumes.
62. United States Geological Survey. Preliminary report on operations of fuel-testing plant of United States Geological Survey at Saint Louis, Missouri, 1905; J. A. Holmes in charge. Introduction and chapter on "briquetting tests," by J. A. Holmes. Bulletin 290, United States Geological Survey, 1906, 240 pages.

63. The San Francisco earthquake and fire of April 18, 1906, and their effects on structures and structural materials. Reports by G. K. Gilbert, R. L. Humphrey, J. S. Sewell, and Frank Soulé, with a preface by J. A. Holmes. Bulletin 324, United States Geological Survey, 1907, 170 pages.
64. Coal-mine accidents: their causes and prevention; a preliminary statistical report, by Clarence Hall and W. O. Snelling, with an introduction by J. A. Holmes. Bulletin 333, United States Geological Survey, 1907, 21 pages.
65. Washing and coking tests of coal and cupola tests of coke, conducted by United States fuel-testing plant at Saint Louis, Missouri, January 1, 1905, to June 30, 1907, by R. G. G. Moldenke, A. W. Belden, and G. R. Delamater, with introduction by J. A. Holmes. Bulletin 336, United States Geological Survey, 1908, 76 pages.
66. Organization, equipment, and operation of the structural materials testing laboratories at Saint Louis, Missouri, by R. L. Humphrey, with a preface by J. A. Holmes. Bulletin 329, United States Geological Survey, 1908, 84 pages.
67. United States Geological Survey. Report of United States fuel-testing plant at Saint Louis, Missouri, January 1, 1906, to June 30, 1907; J. A. Holmes in charge. Introduction by J. A. Holmes. Bulletin 332, United States Geological Survey, 1908, 299 pages.
68. Mining conditions under the city of Scranton, Pennsylvania. Report and maps, by William Griffith and E. T. Conner, with a preface by J. A. Holmes and a chapter by N. H. Darton. Bulletin 25, United States Bureau of Mines, 1912, 89 pages.
69. United States Congress, House of Representatives; Committee on Mines and Mining. Hearing before committee, January 11, 1912; contains statement of J. A. Holmes, Director of Bureau of Mines, on existing law and new bill proposed to meet claims of Western mining men. Washington, D. C., Government Printing Office, 1912, 48 pages.
70. Hearing before committee, Sixty-second Congress, second session, on H. R. 17260, an act to amend an act entitled "An act to establish in Department of Interior a Bureau of Mines," approved May 16, 1910, June 12, 1912; contains statement of J. A. Holmes, Director of Bureau of Mines. Washington, D. C., Government Printing Office, 1912, pages 4-16.
71. Analyses of coals in United States, with descriptions of mine and field samples collected between July 1, 1904, and June 30, 1910, by N. W. Lord, with chapters by J. A. Holmes, F. M. Stanton, A. C. Fieldner, and Samuel Sanford. Bulletin 22, United States Bureau of Mines, 1913, 2 volumes, text, and plates.
72. The use and misuse of explosives in coal mining, by J. J. Rutledge, with a preface by J. A. Holmes. Miners' Circular Number 7, United States Bureau of Mines, 1913, 53 pages.
73. United States Congress, House of Representatives; Committee on Mines and Mining. Hearing (on H. R. 6063, appropriation for mining schools), Sixty-third Congress, second session, December 4, 1913; contains statement of J. A. Holmes, Director of Bureau of Mines. Washington, D. C., Government Printing Office, 1913, 19 pages.
74. Committee on Public Lands. Hearing on bill (H. R. 13137) to provide for leasing of coal lands in Territory of Alaska, and for other purposes, Feb-

W. J. Sutton

ruary 23-26, 1914; contains statements of J. A. Holmes, Director of Bureau of Mines, with an abstract of all bills on opening of coal lands in Alaska. Washington, D. C., Government Printing Office, 1914, part 2, 267 pages.

75. United States Navy Department. Report on coal in Alaska for use in United States Navy. Report of survey and investigation by experimental tests of coal in Alaska, etcetera; contains general statement by J. A. Holmes, Director of Bureau of Mines. House Document 876, Sixty-third Congress, second session, 1914, 123 pages.

MEMORIAL OF WILLIAM JOHN SUTTON

BY WILLIAM FLEET ROBERTSON ¹

William John Sutton was born in Kincardine, Ontario, on January 19, 1859.

His earlier education was acquired in the public schools of Walkerton, Ontario, and later at Trinity College School, Port Hope, Ontario. This was followed by special courses in geology and assaying at Cornell and the Columbia School of Mines.

In 1887 Mr. Sutton moved to British Columbia on the first flood of the "to the West" movement created by the completion of the Canadian Pacific Railway. Settling in Victoria, he was very shortly after appointed the official government assayer, which position he held for some two years.

The report of the Minister of Mines of British Columbia for the year 1888 contains Mr. Sutton's first published report, "A description of the mineral resources of the province" and "List of economic minerals found in the province."

In those earlier days in British Columbia, although placer gold mining had been extensively developed in the interior and coal mining was in operation on the seaboard of Vancouver Island, lode mining had hardly gained even a foothold and the work was in the hands of "practical" men, who, it is more than suspected, classed geologists, chemists, and poets together as purely "ornamental frills," to be respected individually, but of little economic value.

Finding scant financial encouragement in his chosen work, Mr. Sutton, like most of the earlier pioneers, not having "come West for his health," felt obliged to resign from his position as government assayer and entered the lumber business, subsequently securing large tracts of timber and land.

Having thus acquired some little competency, his love for his earlier studies in geology and mineralogy again asserted itself, and feeling the

¹ Written by Mr. Robertson at the request of Dr. R. W. Brock.

need for further instruction in these pursuits, which at that time could not be had here in the West, he went, in 1894, to the Michigan School of Mines at Houghton, Michigan, where he remained four years, taking a technical course and continuing as instructor in mineralogy and crystallography.

Returning to British Columbia, Mr. Sutton entered the service of the Wellington Colliery Company as geologist, and on the company's interests being acquired by the Canadian Collieries (Dunsmuir), Limited, he continued with the new company in the same capacity until the time of his death, on May 19, 1914.

During his connection with the Wellington Colliery Company, Mr. Sutton also acted as consulting geologist for Mr. James Dunsmuir, who had numerous interests in the various metalliferous mining camps of the province.

The Wellington Colliery Company owned the coal rights under a large portion of Vancouver Island, and these were unsurveyed and undefined; it therefore fell to Mr. Sutton to survey and geologically map all these areas and to prospect portions of these for workable coal. In this way he mapped geologically the greater part of the island, an extensive and valuable piece of work, but of such a confidential nature that its publication would have been contrary to the interests of the company.

In connection with this work and his personal timber interests, he acquired a knowledge of the geology of Vancouver Island probably more extensive than that possessed by any other person. He was regarded as *the* authority on the Cretaceous coal-bearing strata of Vancouver Island, and his opinions were often sought by the government departments.

It was always a matter of sincere regret on Mr. Sutton's part and that of his friends that his official position prevented the publication of the mass of valuable geological information he had acquired. It seemed as though he had been obliged by his commercial duties to "keep his light under a bushel," and few but his personal friends and professional *confrères* realized the extent of his geological work.

So irksome to Mr. Sutton had become this feeling of restraint that he had planned to retire from commercial life and to devote his time to the preparation of his geological and mineralogical data for publication and had practically arranged with the writer that such should be done, under the auspices of the British Columbia Bureau of Mines. This intention was, however, frustrated by his sudden death, at Ucluelet, on the west coast of Vancouver Island, while in the active pursuit of his work.

Mr. Sutton was a man of unusually strong physique and his death was probably due to heart failure, as he was never sparing of himself in his

A. B. Willmott

work, and there were few professional woodsmen who could keep pace with him through the dense forests.

While geology was his work, mineralogy was his hobby, and he had personally collected one of the best collections extant of British Columbia rocks and minerals, which he had amplified with rarer and more unique specimens acquired during a trip to Europe.

His interests in science were varied. He was an energetic member of the Natural History Society of British Columbia, being president of the Society in 1912 and 1913, before which he read a number of papers, chiefly on forestry and its conservation.

He was also a member of the Royal Astronomical Society of Canada, the Canadian Mining Institute, and the American Institute of Mining Engineers.

Mr. Sutton's death is a great loss to scientific investigation in British Columbia, where devotees to science are few, and it seems a great pity that he was unable to leave more published records of his store of information.

MEMORIAL OF A. B. WILLMOTT

BY A. P. COLEMAN

Arthur Brown Willmott, son of Rev. J. C. Willmott, of Milton, Ontario, passed away, after a long illness, on May 8, 1914, in his forty-eighth year. He graduated from Victoria University, Cobourg, Ontario, in 1887, taking his arts and science degrees at the same time. He attended Harvard University in 1891, and became professor of chemistry and mineralogy in McMaster University, Toronto, in 1892, continuing in that position until 1900, when he turned his attention to economic geology, becoming field geologist and later manager of mines for the Lake Superior corporation at Sault Ste. Marie, Ontario.

In 1893 he married Mina B. Sanders, daughter of W. B. Sanders, of Stouffville, Ontario.

Mr. Willmott was for some years in the employ of the Bureau of Mines of Ontario, working in Precambrian areas. His greatest interest was in the iron deposits of the Keewatin of Ontario, especially in the Michipicoten region, and he became a recognized authority on the difficult geological and economic problems of the region.

In 1910 Mr. Willmott returned to Toronto to take up work as a consulting mining engineer and to assume the *active* management of several mining companies. In 1913 he was again connected with the Lake Superior corporation in a consulting capacity, which position he held till his death.

The late Mr. Willmott was a most genial and lovable man, who made friends among all classes, and his premature death has been a sorrow to his many friends, as well as a serious loss to science. He leaves a widow and a son and daughter.

BIBLIOGRAPHY

- Michipicoten Mining Division. Bureau of Mines, Ontario, volume 7, 1897, pages 184-206.
- Michipicoten iron range. A. P. Coleman and A. B. Willmott. Bureau of Mines, Ontario, volume 8, part 2, 1899.
- Mineral industries of Sault Ste. Marie. Bureau of Mines, Ontario, volume 11, pages 91-100.
- The Michipicoten iron region. (A. P. C. and A. B. W.) Bureau of Mines, Ontario, volume 11, 1901, pages 152-185.
- The nomenclature of the Lake Superior formations. Journal of Geology, volume X, No. 1, 1902.
- The contact of the Archean and post-Archean in the region of the Great Lakes. Journal of Geology, volume XII, No. 1, 1904.
- The exploration of the Ontario iron ranges. Journal of the Canadian Mining Institute, volume VII, 1904.
- The iron ores of Ontario. Journal of the Canadian Mining Institute, volume XI, 1908.

The Society then proceeded to the consideration of scientific papers.

TITLES AND ABSTRACTS OF PAPERS PRESENTED BEFORE THE MORNING
SESSION AND DISCUSSIONS THEREON

*GEOGRAPHIC HISTORY OF THE SAN JUAN MOUNTAINS SINCE THE CLOSE OF
THE MESOZOIC ERA*¹

BY WALLACE W. ATWOOD AND KIRTLEY F. MATHER

(Abstract)

This paper gave a preliminary summary of physiographic studies which have been in progress during the last six field seasons.

At the close of the Mesozoic era and the opening of the Cenozoic era there were mountain-making movements which affected the entire Rocky Mountain province of North America, and the great dome which was then formed in the San Juan area was at once subjected to vigorous erosion. As the mountain mass rose erosion began, and as the great dome was more and more deeply dissected a mountain topography must have been produced, and those mountains may be thought of as the first generation of the San Juan Range. Certain deposits of Eocene till in this region indicate that during the dissection of these early San Juan Mountains ice formed in the range and descended to the bordering lowlands.

¹ Presented with permission of the Director of the U. S. Geological Survey.

After the retreat and disappearance of the early Tertiary ice, stream erosion continued, and the western portion of the San Juan Mountain area was reduced to a surface of slight relief which may be thought of as a peneplain. After the deposition of the Telluride conglomerate on this peneplain there was further erosion in the range, and then came the three great epochs of volcanism—the San Juan, the Silverton, and the Potosi. During these epochs of volcanism a great volcanic plateau was developed. By this time the Miocene epoch had been reached and possibly passed, and with the quieting down of volcanic activity began the erosion and dissection of the volcanic plateau. During this period of dissection another generation of San Juan Mountains was carved, this time out of volcanic debris and great lava flows.

The San Juan Mountains that were first carved out of this great volcanic plateau should then be thought of as surmounting those of today. Perhaps, if replaced, they would rise 3,000 or 4,000 feet above the present summits. They rose above the present summit peneplain.

With the redoming of the area, which involved the warping or doming of the summit peneplain, another cycle of erosion was begun. Valleys were again formed, and in these valleys snows collected which in time formed glaciers that advanced to the lowlands bordering the range. These earliest Pleistocene glaciers retreated and disappeared. The range continued to be uplifted, and the streams were so rejuvenated that they cut great canyons below the broad troughs occupied by the Cerro glaciers. Again, climatic changes favored the formation of ice among the summits, and that ice (the Durango glaciers) descended through the main canyons to the foothills and later retreated and disappeared. The canyons were still more deeply cut into the mountain mass, and then climatic conditions favorable for glaciation once more returned and the Wisconsin or third series of Pleistocene glaciers formed and descended through the great canyons, nearly as far as those of the Durango stage. These glaciers have now disappeared, and there is no true glacier ice remaining in the region today; but the streams are vigorously dissecting the mountain mass to still greater depths. The Black Canyon of the Gunnison has been largely cut during and since Pleistocene time. The vigor of the stream work is illustrated in many a sharp V-shaped notch cut below the depth of ice-action. High among the mountains are the remarkable landslides and great accumulations of talus.

The studies suggest somewhat continuous mountain growth in this region during late geologic time.

Presented in abstract extemporaneously by the senior author.

*DOMINANTLY FLUVIATILE ORIGIN, UNDER SEASONAL RAINFALL, OF THE
OLD RED SANDSTONE*

BY JOSEPH BARRELL

(Abstract)

The old red sandstones of the British Isles have been commonly interpreted as deposited in great lakes. Walther has, however, urged that these formations are desert deposits; Goodchild regards them as laid down partly in large inland lakes, partly as torrential deposits, partly as old desert sands.

Extensive volcanic material is intermixed. The present writer interprets them as dominantly fluviatile deposits, formed under semi-arid climatic conditions. This stands between the extreme interpretations, but is essentially different from either.

The subject is important from the standpoint of general and historical geology, but perhaps especially from that of organic evolution, since these deposits contain the oldest well known fish faunas in contrast to the rare and fragmentary fossils of the previous periods. If the stratigraphic record is misinterpreted, there is given, in consequence, an erroneous conception of the habitat and life adaptations of the advancing evolutionary wave of the vertebrate phylum when it first clearly appears in the geologic record.

A reinterpretation must rest on the criteria as to the mode of origin of sediments as much as on the actual stratigraphic features. It is the matching of more recent criteria to the older known facts of the stratigraphy of the old red sandstone which makes the basis of this paper.

Presented in abstract extemporaneously.

Remarks were made by Messrs. A. W. Grabau and J. M. Clarke, with reply by the author.

INFLUENCE OF SILURIAN-DEVONIAN CLIMATES ON THE RISE OF AIR-BREATHING VERTEBRATES

BY JOSEPH BARRELL

(Abstract)

The relationships of ancient faunas to their environments is a field wherein paleontology and physical geology meet. It is a field which has been commonly cultivated by the former, but it is one in which the latter may as logically enter. It was as a physical geologist, with ideas sharpened and made definite by previous study of the nature of Devonian sediments, that the present writer took up this subject.

The Devonian formations indicate the general presence of warmth and seasonal rainfall. In the Upper Devonian the general climatic conditions became more markedly semi-arid. There is found to be a concurrent elimination of sharks from the fresh waters. As a result, dipnoans and crossopterygians come to dominate the fauna. It is apparently in the Upper Devonian, furthermore, that amphibians began to expand. An examination is made of the various possible causes for this advance in evolution. The only one which is found adequate is the compulsion of seasonal dryness.

The actual line along which air-breathing developed was only one of several possible lines. The directions and limitations of later evolution were, however, more or less determined by this choice. Slightly different conditions of environment and organic response, both readily possible, might apparently have been more favorable for the future of air-breathing vertebrates. By appreciating these lost opportunities of the remote past a better perspective is obtained, on the one hand, of the devious and groping and not always most successful nature of organic progress; a better appreciation, on the other hand, of the dangers of stagnation or extinction which have been happily passed by.

This and the previous paper were presented by the writer in abstract to the

American Society of Vertebrate Paleontology on December 26, 1907, but have been withheld from publication. The present presentation will place the emphasis on the more general aspects and consequences of the problems of the Devonian climates in relation to the rise of air-breathing vertebrates. It is intended to publish the papers in the near future.

Presented in abstract extemporaneously.

*SOME LITTORAL AND SUBLITTORAL PHYSIOGRAPHIC FEATURES OF THE
VIRGIN AND NORTHERN LEEWARD ISLANDS AND THEIR BEARING ON
THE CORAL-REEF PROBLEM*

BY THOMAS WAYLAND VAUGHAN¹

(Abstract)

The ocean bottom off the shores of the Antilles shows three distinct types of profiles, and a fourth type is furnished by Saba and other banks. The first is that found off the volcanic islands, such as Saba and the members of the Saint Christopher Chain, into the sides of which the sea has cut relatively narrow platforms; but there are suggestions of submerged flats off the northwest end of Saint Eustatius and southeast of Nevis.

The second type of submarine profile is well represented off the north shore of Saint Croix and the south shore of Cuba. The precipitous character of these profiles indicates faulting, and the geologic structure supports this interpretation. There is a down-thrown block between the Virgins and Saint Croix and another between Cuba and Jamaica.

The third type of profile, represented by shores off which are extensive shallow flats, occurs where planation agencies have long been active. Here the rocks often, if not usually, dip under the sea at relatively gentle angles.

The fourth type of profile is represented by the extensive submerged banks or platforms which have no bordering lands and whose upper surfaces range in depth from 9 to 30 fathoms. Good examples are Saba Bank, southwest of Saba Island; Pedro Bank, southwest of Jamaica, and Rosalind Bank, off Mosquito Bank, which is the continental shelf northeast of Nicaragua and Honduras. That the depth of water on these banks is essentially the same as in many atolls of the Pacific, especially the Paumotu, has been repeatedly pointed out, but apparently the fact has not yet been sufficiently emphasized.

The third type of profile (that showing submarine terraces around islands) will now be discussed in some detail. From shoreline characters and other evidence the conclusion was reached that the Virgin Islands, the members of the Saint Martin group, and Antigua and Barbuda have recently undergone submergence to an amount of about 20 fathoms.² Assuming this conclusion to be correct, should the sealevel have remained stationary for a period of appreciable length antecedent to this submergence, there should be a submerged scarp or facet indicating its former stand; should there have been a succession of temporary stands, there should be a series of submarine terrace flats separated by scarps. The available sources of information were the charts of

¹ This article, illustrated by more than fifty profiles and a map, is published in the Jour. Wash. Acad. Sci., vol. 6, pp. 53-56, Feb. 4, 1916.

² Bull. Am. Geog. Soc., vol. 46, 1914, pp. 426-429.

the United States Hydrographic Office and of the British Admiralty. The Virgin Bank and the Saint Martin Plateau were selected for special study. The charts of the former, on a scale of slightly more than 1 mile to an inch, and that of the latter, on a scale of about $2\frac{1}{2}$ miles to an inch, were contoured on a 2-fathom interval from the shore to a depth of 40 fathoms, and on an interval of 10 fathoms in depths between 40 and 100 fathoms.

The shoreline of the Virgin group shows indentations indicative of submergence, and that the sea has stood at its present level long enough for alluvial filling of the heads of harbor digitations, while sea-cliffs occur at the ends of promontories. The chart of the near-by sea-bottom shows that south of Saint John, Tortola, and Virgin Gorda there are two distinct submerged terraces and a less definite third terrace. The outer terrace flat lies at depths between 26 and 28 fathoms on its landward and between 28 and 30 fathoms on its seaward margin, and it ranges in width from a half mile to 3 miles. On its sea front is a ridge which is inferred to be a submerged barrier coral reef. On its landward side a scarp rises from a depth of 26 or 28 fathoms to about 17 fathoms. Above this scarp is a second terrace flat, which has a depth of 14 to 15 fathoms on its landward and a depth of 14 to 20 fathoms on its seaward face, and ranges in width from one-third of a mile to 2 miles. Apparently the outer margin of this flat also bears a coral reef. These are the two principal terrace flats. The scarp separating them is indicated by crowded contours, and chart number 1832, United States Hydrographic Office, shows its continuity for 36 nautical miles, or about $1\frac{1}{2}$ land miles farther than from Washington to Baltimore. A third still higher terrace flat is suggested between depths of 6 and 10 fathoms, above which a fourth terrace may now be in process of formation, but the information regarding these is at present not definite enough to warrant a positive statement. The *continuity* of the upper one of the two well marked flats needs to be emphasized. It should be noted that east of Virgin Gorda there has been an uptilt.

On the windward side of Saint Thomas there is an extensive outer flat, bounded on its landward side by a steep escarpment which in places is nearly 160 feet high. The landward margin of the plain is between 26 and 28 fathoms in depth; the seaward margin has a depth between 30 and 34 fathoms; the width is as great as 10 miles and for distances as great as $8\frac{1}{2}$ miles, in depths between 29 and 31 fathoms, the range in relief of the surface may be as small as 2 fathoms. Its outer margin is cut by reentrants which have bottoms about 40 fathoms deep and simulate hanging valleys. There are also near the outer margin of this flat banks or ridges, the upper surfaces of which are relatively flat, between 17 and 20 fathoms in depth. One of these banks has a total basal width of about 4 miles and a length of more than 5 miles. As its form is not that of a coral reef, it can only be the base of what was an island, which had been reduced almost to a smooth surface by marine planation and then, as indicated by other evidence, submerged. As all the other shoals, with one exception, are truncated at nearly the same level, it seems that most of them should be ascribed to a similar origin. These shoals usually show escarpments between 20 and 30 fathoms on their windward sides and more gradual slopes on the leeward sides. The outer flat on the north side of Saint Thomas corresponds to the lower flat on the south side of Saint John, Tortola, and Virgin Gorda. Both are submarine plains, which several lines of evidence

show were developed when sealevel was about 20 fathoms, or slightly more, lower than now. The escarpment extending from the islands north of Culebra Island, east of Porto Rico, across the Virgin Passage, and along the north side of Saint Thomas, and the escarpment on the seaward face of the outlying shoals apparently can be explained in no other way.

The indentations on the outer margin of the outer flat may have been caused by emergence and stream cutting after its formation, or they may be due to initial marginal irregularities which have not been obliterated.

The approximate accordance in level of the tops of the outlying shoals at depths between 17 and 20 fathoms has been mentioned. These summits accord in height with a flat or gently sloping zone, which lies above and nearer shore than the deeper flat and represents the 14 to 20 fathom flat south of Saint John and Tortola. It is scarcely represented on the seaward side of the promontories, namely, Cockroach and Cricket rocks and Outer Brass and Little Hans Lollick Islands. However, it spreads out on the flanks of the promontories and ranges from half a mile to nearly $1\frac{1}{2}$ miles in width; it is separated on its seaward side by a steep slope or escarpment from the deeper flat and on its landward side by a less distinct escarpment, in places about 26 feet in height, from a less developed flat, which has a depth of 7 to 10 fathoms. The descent is sudden from the shore to about 6 fathoms, which is near the landward margin of the highest submarine flat. This flat also is narrow on the tips of the promontories mentioned, but widens on their flanks and along the shores of the main island. The submerged valley in Charlotte Amalia Harbor has a depth of 10 fathoms.

The narrowness or absence of the 14 to 20 fathom flat on the promontory tips, while it is so well preserved in protected places, especially off the south sides of Saint John and Tortola, shows that it is older than the deeper flat, and in exposed places was cut away during the formation of the latter, subsequent to the formation of which, after perhaps a brief interval of still lower stand of sealevel, the entire area has been resubmerged to an amount about the same as that of the initial submergence.

There is doubt as to the interpretation of the 7 to 10 or 12 fathom flat. In places it seems to be distinct and older than the one next lower, but it may represent the submarine terrace being formed at present sealevel.

According to the physiography of the sea-bottom, the Virgin Islands were joined to Porto Rico during the cutting of the scarp separating the deepest from the next higher flat. The biogeographic evidence shows conclusively that the two were united and have been severed in Recent time by submergence. Stejneger says in his *Herpetology of Porto Rico*: "It is then plain that the 16 species of reptiles and batrachians found in Saint Thomas and Saint John form only a herpetological appendix to Porto Rico." Doctor Bartsch informs me that the testimony of the land Mollusca is the same as that of the reptiles and batrachians. The biogeographic evidence substantiates the deductions based on the purely physiographic study.

There are three tiers of coral reefs in the Virgin Islands. They rise above (a) basements 10 fathoms or less in depth; (b) above the outer edge of the 14 to 20 fathom flat; (c) above the outer edge of the 28 to 34 fathom flat. As the escarpment within the outermost reef could not have been cut during the presence of such a reef, the flat must be older than the reef and the reef

must have developed during subsequent submergence. The flat, therefore, can not be due to the growth of the reef.

The members of the Saint Martin group have indented shorelines, sea-cliffs, and an unusually fine development of bay-bars. The relations on the windward side of the Saint Martin Plateau are similar to those north of Saint Thomas. The outer, deeper flat, from 26 to 36 fathoms in depth, has a maximum length, east and west, of over 30 miles. It seems composed of two terraces. The scarp on its landward side is distinct and in places is about 50 feet high, between 20 and 28 fathoms, as off the east end of Scrub Island, east of Anguilla Island.

As some of the submerged valleys on the east side of the Saint Martin Plateau resemble valleys in the Upper Cretaceous Anacacho limestone Texas, it appears that not only must the scarp line which has been pointed out be interpreted as a former shoreline, but that these channels with steep heads must be interpreted as former drainage lines which were subaerially cut and afterward submerged. The Anacacho limestone in the Brackett quadrangle is similar in general character to the limestone which composes Anguilla and Tintamarre.

While the shoreline stood some 20 fathoms lower than now, the Saint Martin Plateau must have been entirely above sealevel. The biologic evidence is in accord with this interpretation, but at present it alone is not sufficient to be decisive.

Antigua is another island with an indented shoreline. It shows typical instances of submerged valleys and fairly good examples of pouch-shaped harbors. Profiles off the southeast shore exhibit essentially the same features as the profiles on the Virgin Bank and the Saint Martin Plateau. If sealevel stood 20 fathoms below its present stand, Antigua and Barbuda would be united. Doctor Bartsch has especially studied the land mollusca and says: "The land shells show that these islands must have been connected in very recent time."

The deduction that there has been in Recent geologic time submergence to an amount of about 20 fathoms in the Virgin Islands, on the Saint Martin Plateau, and on the Antigua-Barbuda Bank, it seems to me, may be accounted demonstrated.

A set of profiles on the same vertical scale—(a) across Havana Harbor, showing depth of filled channel; (b) off the north side of Saint Thomas; (c) off the west side of Anguilla; (d) off the southeast coast of Antigua; (e) Mosquito Bank, off Nicaragua—all indicate a rise of sealevel by an amount of about 20 fathoms. There is in the Virgin Islands and in Cuba clear evidence of a lowering of sealevel by about 20 fathoms, perhaps more, previous to resubmergence. Although the evidence for the other areas is not definite as to the return of sealevel to a former stand, the similarity of the profiles suggests that it also occurred in them. As this lowering and subsequent rise of sealevel affects a large area, it appears too wide-spread to be explained by local crustal movement. The changes in position of strand-line here noted are more reasonably explained by the lowering of sealevel, due to the withdrawal of water in the Pleistocene ice epochs to form to great continental glaciers, and the raising of sealevel after each epoch through the melting of the glaciers; but the volume of evidence supplied by this area is perhaps not large enough

to justify a general conclusion as to relations of Recent coral-reef development to glaciation and deglaciation.

A brief comparison will now be made with the Great Barrier Reef of Australia. Fifteen profiles, on the same horizontal and vertical scales, the latter about 70 times the former, were drawn on the British Admiralty charts across the continental shelf south of the Great Barrier Reef and across the reef. These profiles show the continuity of the platform from the area south of the Great Barrier, that there is an outer, deeper flat about 200 feet deep, and that, except near its north end, the reef stands back from the seaward edge of the continental shelf. Therefore, apparently the idea that the platform was formed by infilling behind the reef may be permanently set aside. There is striking general similarity of the conditions presented to those off Nicaragua and in the West Indies. The evidence in favor of a shoreline between about 25 and 30 fathoms below present sealevel antecedent to Recent submergence is strong, if not conclusive, and supports the deduction that the living barrier reef is growing on what was a land surface in Pleistocene time—an interpretation essentially that proposed by E. C. Andrews in 1902.

The relations around the Pacific islands off which barrier reefs occur are those of continuous platforms surmounted or margined by discontinuous reefs. These relations indicate the superposition of reefs on antecedent platforms which have undergone geologically Recent submergence. E. C. Andrews so interprets the conditions of formation of the barrier reefs off the Fiji Islands.² It appears to me that the conditions governing the development of the living reefs in the West Indies, Central America, Brazil, Florida, and Australia are clear. The reefs have grown on antecedent basements during Recent submergence. The history of these basements is complex, but during Pleistocene time they stood higher with reference to sealevel than now; their outer margins were remodeled by marine cutting and marine planation, and they were then resubmerged. These changes in height of sealevel accord with the demand of the glacial control theory. It would be remarkable if the conditions in the tropical western Pacific Ocean were exceptional, and the present available facts indicate that they conform to the principles governing reef development in the other areas. Here it should be said, regarding the charts for the Pacific, that as they have been made primarily for navigation purposes, the depths of lagoons and lagoon channels are often given in a way fairly satisfactory, but on only a few charts can the submarine profiles outside the reefs be determined. The coral-reef problem can not be regarded as satisfactorily solved until the relations in the Pacific islands have been ascertained. In my opinion, but little further advance in understanding the problem can be expected from purely biologic studies or from physiographic investigations of the dry land surface alone. As apparently the greatest present need is for more accurate information on the detailed submarine relief in depths between 15 and 50 fathoms, especially on the seaward margins of the platforms, both outside the reefs and off the breaks in the reef lines, the efforts of those interested in such investigations should be concentrated on getting additional hydrographic surveys in coral-reef areas.

Presented by title in the absence of the author.

² *Am. Jour. Sci.*, vol. 41, Jan., 1916, pp. 135-141.

CORAL-REEF PROBLEM

BY W. M. DAVIS

(Abstract)

The author, having reviewed various coral-reef theories preparatory to his Shaler Memorial voyage of 1914, and having visited on that voyage thirty-five reef-encircled islands in the Pacific, is now preparing a report on his observations and inferences. He finds in this connection only three theories that demand serious consideration—Darwin's theory of subsidence, Daly's theory of glacial control, and Vaughan's theory of submerged platforms. The present paper sets forth the difficulties that stand in the way of accepting two of these theories. The theory of glacial control seems inadequate because it excludes subsidence on insufficient grounds, because it assumes unwarranted conditions for preglacial islands, and because certain essential consequences of its main process—the truncation of low preglacial islands by the lowered and chilled ocean of the Glacial period—are not found on Pacific islands. The theory of submerged platforms is unsatisfactory because it assumes without sufficient warrant the origin of the platforms independent of coral agencies, because it assumes the absence of reef-building corals during the production of the platforms, because it provides no adequate cause for the production of flat platforms in close association with steep volcanic islands, and because it excludes all but small changes of level in reef-encircled islands during recent geological periods.

Presented in abstract extemporaneously.

Remarks were made by Professors R. A. Daly and J. P. Iddings, with reply by the author.

TERTIARY-QUATERNARY OROGENIC HISTORY OF THE SIERRA NEVADA IN THE LIGHT OF RECENT STUDIES IN THE YOSEMITE REGION¹

BY F. E. MATTHES

(Abstract)

It is a noteworthy fact that the hanging tributaries of the Merced are not all graded with respect to the same former profile of the master stream. Those flowing over the massive and resistant granites of the Yosemite region indicate an old profile situated about 2,800 feet above the present canyon floor. Those flowing over normally jointed and relatively easily eroded rocks indicate a more strongly concave profile of a later date, situated 600 to 1,200 feet lower. The older profile of the Merced appears to be coordinate with the profiles determined by Lindgren for the Tertiary rivers of the northern Sierra Nevada (by means of the auriferous gravels). The younger profile of the Merced is believed to correspond to that stage of erosion which Lawson has recognized in the Chagoopa Plateau of the Kern River country. It clearly antedates the cutting of the inner gorge of the Merced, just as the mature valleys of the Chagoopa Plateau antedate the cutting of the Kern's narrow canyon. Inner

¹ Published with permission of the Director of the U. S. Geological Survey.

trenches of the same order occur in most of the great canyons of the western Sierra slope. Their youthful aspect, coupled with the fact that they are still actively being deepened by the rivers, points strongly to their initiation in consequence of rapid tilting at a time which can scarcely have been more remote than the beginning of the Quaternary period. The amplitude of this uplift at the eastern edge of the Sierra block may be roughly estimated from the inclination of the younger profile of the Merced. That profile appears at least four times as steep as a graded profile of the river might be expected to have been. It is to be inferred, therefore, that the range crest at the head of the Merced stood, prior to the last uplift, only one-fourth as high as it now stands. Mount Lyell, instead of 13,090 feet, had an altitude of only some 4,400 feet. This, it will be observed, is 2,000 feet less than the present altitude of Mono Lake (6,417 feet). It would follow, then, that the regions to the east of the Sierra Nevada have also been uplifted broadly by several thousand feet since the end of Tertiary time.

The Sierra uplift indicated by the older profile of the Merced is considerably less than that indicated by the younger profile and probably did not exceed 2,500 feet.

Presented in abstract extemporaneously.

The Society adjourned about 12.30 o'clock and reconvened at 2 o'clock.

TITLES AND ABSTRACTS OF PAPERS PRESENTED BEFORE THE AFTERNOON
SESSION AND DISCUSSIONS THEREON

The Society reconvened at 2 o'clock, with President Coleman presiding and Charles P. Berkey acting as Secretary, and took up the consideration of scientific papers.

GEOLOGICAL TRANSFORMATIONS OF PHOSPHORUS

BY ELIOT BLACKWELDER

(Abstract)

Phosphorus migrates widely in and upon the earth, passing meanwhile through a varied and interesting series of metamorphoses. An attempt will be made to outline these changes, beginning with the crystallization of apatite and other phosphates in igneous rocks and primary veins, following them through the circulation of ground-water to the ocean, there to undergo an all but endless series of reincarnations in the bodies of organisms, and ending temporarily with the fixing of the element in the form of phosphatic sediments. Subsequent changes take place in the latter, both near the land surface and in the deep interior of the earth. The various known types of phosphatic deposits may be placed rather definitely in this metamorphic cycle.

Presented in abstract extemporaneously.

*DIFFUSION IN SILICATE MELTS*BY N. L. BOWEN ¹*(Abstract)*

This paper gave a brief description of some determinations of the rates of diffusion of molten rock-forming silicates and a discussion of the significance of the results in petrologic problems.

Presented in abstract extemporaneously.

DISCUSSION

Prof. J. P. Iddings remarked that Mr. Bowen's investigations were made on stationary liquids, but it is not to be supposed that large bodies of magma having different temperatures in different parts will exist without convection currents for any appreciable length of time, and his remark that convection currents may play a prominent rôle in differentiation processes is very much to the point. The effect of gaseous constituents in molten magmas, or of other components which reduce the viscosity of such liquids, must also be taken into account.

PETROGRAPHY OF THE PACIFIC ISLANDS

BY E. A. DALY

(Abstract)

The total number of named islands in the open Pacific is about 3,000. Of these, only 22 are reported to show outcrops of quartzose rocks. Five other islands are reported to have outcrops of crystalline schists, serpentine, or deformed limestone.

Excluding New Guinea and New Zealand with their immediate satellitic islands, Oceania has 345 islands which have been definitely described as wholly or largely volcanic in origin. Probably the whole number showing volcanic rocks above sealevel is at least twice as great. Among these islands only 152 have yet afforded any petrographic data, and not one has been examined with desirable thoroughness. To illustrate the scrappiness of our information as well as certain problems regarding the origin of Pacific lavas, the igneous-rock types of each island, so far as recorded, have been tabulated.

The compilation illustrates the advisability of a systematic exploration of all the smaller islands of the Pacific. The task is quite feasible, since the total land area involved is less than 75,000 square miles. If Hawaii, Viti Levu, Vanua Levu, New Pomerania, New Mecklenburg, and New Caledonia be excluded, the total land area to be covered would be only 50,000 square miles. In fact, with relatively small effort and cost, the natural history (including the petrography and geology) of all the land areas within one-eighth of the earth's surface could be investigated by a single organization. The Geological Society may well use its influence in developing a plan for such comprehensive exploration in the Pacific, under private American auspices.

Presented in abstract extemporaneously.

¹ Introduced by C. N. Fenner.

DISCUSSION

Prof. J. P. Iddings heartily approved of the proposition of Professor Daly regarding the systematic survey of the islands of the Pacific and suggested the desirability of having a properly equipped steamer for carrying out the survey.

Further remarks were made by Messrs. A. C. Lane and N. L. Bowen, with reply by the author.

SOME FACTORS WHICH AFFECT THE DEPOSITION OF CALCIUM CARBONATE

BY JOHN JOHNSON ¹

(Abstract)

This paper gave a brief discussion of the operation of the inorganic factors which affect the solubility, hence the precipitation of calcium carbonate, and therefore are important in connection with the origin and mode of formation of limestones. Of these factors there are two in particular, namely, the concentration of free carbon dioxide in the water and the temperature, which exert an influence more far-reaching than has hitherto been generally recognized; indeed, it appears that by considering their effects alone we are enabled to coordinate a large number of the phenomena of the deposition of limestones.

Presented in abstract extemporaneously.

Remarks were made by Dr. A. C. Lane, with reply by the author.

SPECIFIC WEIGHT OF DRILL CORES

BY ALFRED C. LANE

(Abstract)

It has been found that the specific gravity of drill cores, or volume per cubic foot, can be readily obtained by measuring their dimensions. This can be done to an accuracy of within 1 per cent, in pieces of core over 100 mm. long, without grinding off the ends, by using the micrometer gauge for the diameter and taking the average of six measurements of the diameter and the average of about four measurements of the length made with a finely graduated ruler. Thus, after weighing them, the weight in grams per cubic centimeter or (the same thing) ounces per cubic foot is obtained. Systematic tests seem to show that this may be of considerable practical value, as the variation in one lava flow between the denser, more crystalline, less glassy, and the more glassy, less crystalline, and perhaps more altered upper and lower parts, can be distinctly and continuously followed, even when the amygdaloidal or vesicular structure is not conspicuous.

Presented in abstract extemporaneously.

¹ Introduced by H. S. Washington.

CHEMICAL AND MINERALOGICAL COMPOSITION OF METEORITES

BY GEORGE P. MERRILL

(Abstract)

The paper gives a brief résumé of researches on the subject indicated by the title, which were made with especial reference to the reported occurrences of minor constituents. No traces were found in the stones and irons examined of antimony, arsenic, barium, gold, lead, strontium, tin, tungsten, uranium, zinc, or zirconium. On the other hand, the presence was shown, beyond an apparent reasonable doubt, of the rarer elements iridium, platinum, palladium, ruthenium, and vanadium. Comparisons are made between the meteorites and terrestrial rocks, consideration being given to the efficacy of the former as world-forming materials. A continuation of work, preliminary reports and abstracts of which have been published in the *American Journal of Science* and the *Proceedings of the National Academy of Sciences*, and final results of which are to appear in one of the memoirs of the Academy.

Read in abstract from notes.

DISCUSSION

Prof. O. C. FARRINGTON remarked that the long series of investigations on meteorites which Doctor Merrill had made had yielded many valuable results and were likely to yield more. To the list of elements found in meteorites given by Doctor Merrill it seems certain that radium can be added, since it has been found in one stone meteorite by an English analyst and is indicated by some unpublished results obtained in this country of which I had been notified. It has, so far, not been found in any iron meteorites. In comparing stony meteorites with the crust of the earth, I urged that the general average of stony meteorites could not properly be used, since most stony meteorites were of higher specific gravity than the rocks of the earth's crust. For the purpose of comparison, the class of stony meteorites known as eukrites should be used, since meteorites of this class most nearly resemble the rocks of the earth's crust in specific gravity and are composed of feldspars and pyroxenes, which are the dominant minerals in the earth's crustal rocks.

Doctor Merrill replied briefly to Professor Farrington's remarks.

IMPORTANCE OF WATER AS A MAGMATIC CONSTITUENT

BY GEORGE W. MOREY¹*(Abstract)*

That water is an original constituent of the magma is now generally admitted, but the importance of its effects has not received adequate recognition. The erroneous assumption is often met with that, since the critical temperature of pure water is about 370°, it can exist, at temperatures higher than this, only in the gaseous state, no matter how great the pressure to which it

¹ Introduced by Henry S. Washington.

EXAMPLES OF AUSTRALITES

Figure 1 is a dumb-bell type of australite from Victoria, Australia. After Dunn. Figure 2, Plate V, Bulletin Geological Society of Victoria. 4/5 natural size.

Figure 2, A-G, are different forms of "Pele's Tears" from Kilauea, Hawaii. Enlarged three times. Drawn by Edw. C. Selbert.

is subjected. On the contrary, water can exist in solution in liquid magmas, and as such unquestionably plays a very important rôle in geologic processes which go on at temperatures up to 1,000° or higher. The presence of water in such solutions has three main effects, namely, it increases the fluidity of the melt, lowers the temperature at which crystallization begins, and facilitates the process of crystallization; these effects are marked even when the quantity of water is relatively small. These facts, which have an important bearing on certain phenomena of volcanism and igneous intrusion, were discussed and illustrated.

Presented in abstract extemporaneously.

DISCUSSION

Prof. J. VOLNEY LEWIS: It is a well known fact that fragmental volcanics of acid character are far more abundant than basic ones. Some of us are accustomed to attribute this to the greater abundance of water in siliceous magmas, as a rule, or to their greater viscosity and hence greater resistance to the escape of this water during eruption. May it not be possible that the difference is in reality due, in part at least, to the greater capacity of basic magmas for retaining water in solution until it is gradually released by crystallization or imprisoned permanently in glass? I should be glad to know whether or not the investigation of which Doctor Morey has spoken might throw light on this question.

Doctor Morey replied briefly to Professor Lewis's remarks.

"PELE'S TEARS" AND THEIR BEARING ON THE ORIGIN OF AUSTRALITES

BY E. S. MOORE

CONTENTS

	Page
Introduction.....	51
Characteristics and origin.....	52
Comparison with similar bodies.....	53
Chemical constituents.....	54
Conclusion.....	55

INTRODUCTION

Among the problems confronting the Australian geologists few offer more interesting material for speculation than the question of the origin of the australites. These peculiar bodies have been known by various names, such as obsidian buttons, obsidian bombs, and obsidianites. They somewhat resemble in composition and form the moldavites and billitonites, so Suess¹ has applied the name australites to such bodies from Australia, while he uses the term tektites to cover all bodies of this type.

It is not the object of this paper to go into a detailed discussion of the australites, as they have already been described very fully by several of the

¹ Franz Suess: Die Herkunft der Moldavite und verwandter. Gläser Jahrb. d. k. k. Geol. Reichsanstalt. Heft 2, vol. 50, 1900.

Australian and other geologists. Numerous analyses and a valuable discussion of the origin of these bodies will be found in a paper by H. S. Summers,² and an excellent set of figures accompanying a discussion of the genesis has been published by E. J. Dunn,³ former Director of the Geological Survey of Victoria, Australia. In these publications various references to the bibliography on this subject will also be found.

CHARACTERISTICS AND ORIGIN

The australites are small bodies of glass, often of remarkably perfect form, so perfect in some cases as to suggest an artificial origin. They are sometimes button-shaped, sometimes dumb-bell in shape, sometimes spherical, and at other times quite irregular in form. Many of them show flow structures and often pitted surfaces are developed. In chemical composition they are much like obsidian, although they are peculiar, in having in most cases a higher lime and magnesia content in proportion to potash and soda than is generally found in rocks so high in silica and in which the potash is usually higher than the soda. Summers, therefore, endeavors to show that they do not correspond to any terrestrial rock type. They are undoubtedly related to one another in chemical composition, as he points out, and most of them can doubtless be regarded as having a common source, yet comparison of the analyses with other special types show a close resemblance to some types of obsidian.

There have been advanced several theories to account for the origin of these peculiar bodies, the two main theories being the volcanic and the meteoric. The greatest difficulty so far confronting the volcanic theory is the distribution of the australites, which occur over a band stretching across Tasmania and over the whole of the southern part of Australia from the east to the west, covering the greater part of the island south of the tropic of Capricorn. Various suggestions have been made to account for their transportation, among them Mr. Dunn's ingenious bubble hypothesis, transportation by aborigines, emus, etcetera. These means have never appeared entirely satisfactory because, as pointed out by Summers, the australites are believed to be of Cenozoic age—although I believe they have never been found embedded in any solid formations—and that while there may be known volcanoes of this age in Victoria, some of the specimens are still found at least 2,000 miles from these volcanoes, and so far no deposits of obsidian have been found connected with them.

To account for the distribution, therefore, the meteoric hypothesis has been advanced, but this is open to as great objections as the other. In the first place, the composition, form, and internal structure of the bodies are quite different from those of known meteorites, and in the second place it seems hard to realize how such perfectly formed bodies, which give every indication of having been in a liquid condition, could strike the earth from some extra-terrestrial source and have their outline so well preserved. If the heat which caused them to become liquid were due to atmospheric friction, it would be

² H. S. Summers: Obsidianites—their origin from a chemical standpoint. Royal Soc. of Victoria, vol. xxi (new series), 1909.

H. S. Summers: On the composition and origin of australites. Report of the Australian Association for the Adv. of Sci., vol. xiv, 1913, pp. 189-199.

³ E. J. Dunn: Australites. Bull. No. 27, Geol. Survey of Victoria, 1912.

greatest at the moment of impact and the body must certainly suffer deformation. On the other hand, if they had been heated, so that they could assume their present form before entering the atmosphere, it would be expected that the heat of atmospheric friction would cause the exterior to scale off and the perfect form to be destroyed.

Other suggestions concerning the genesis have been made by Gregory and G. P. Merrill. Gregory has stated that these bodies may have been due to lightning discharges in the great dust storms of Australia; but this theory has not received serious consideration, because it has neither the merit of satisfying the demand for distribution nor explaining the form and composition of the bodies.

In his article entitled "On the supposed origin of the moldavites and like sporadic glasses from various sources," after examining various obsidian pebbles, moldavites, and billitonites, Merrill concludes with the statement that "whatever may have been their original source, the Bohemian and Moravian specimens are now simply water-worn pebbles of weathered glass, originally etched by corroding vapors or solutions, the results being indistinguishable from those produced by artificial etchings on obsidian with fluorhydric acid. The Australian forms are likewise, to me, simply pebbles of glass which have been water-worn or abraded by wind-blown sands. In their contours there is nothing even suggestive of meteoric markings, nor do I find any semblance of such an origin, so far as the surface markings alone are concerned, in the examples from Billiton."⁴ He explains, however, that this statement is not to be construed as meaning that he is opposed to the cosmic origin of these bodies.

COMPARISON WITH SIMILAR BODIES

Having thus briefly considered the australites, let us turn to some bodies of similar form whose origin is established beyond a doubt. There have been found two types of such bodies. One of these types, as mentioned by Mr. Summers,⁵ has been taken from the smoke-boxes of locomotives, and the other is found near Kilauea, Hawaii, in the form of "Pele's tears." The latter are little masses of lava of various shapes, to which I believe Mr. Perret first applied the name "Pele's tears" because of their relation to Pele's hair and the tear-drop form of many of them. My attention was first attracted to these through the kindness of Dr. T. A. Jaggar, and they seemed to show such a marked similarity in form to certain of the australites that they may have an important bearing on their origin. So far as observed, they are confined to a small spatter cone close to the main crater of Kilauea. They vary from one-sixteenth to about one inch in length and in diameter according to the shape. They exhibit various forms. Such shapes as rings, half rings, pears, tear-drops, hearts, spheres, spindles, and dumb-bells are common. In one case an oval central body surrounded by a ring, which seems, however, to be more or less distinct from it, was found. They consist of a pumiceous dark brown glass, with a smooth glazed coating covering the surface of the body where the original surface is unbroken. A little evidence of flow structure may be seen on the surface in some cases, but no such perfect button forms as exhibited by some of the australites have been found (see figures).

⁴ G. P. Merrill: Proc. U. S. National Museum, vol. 40, 1911, p. 486.

⁵ Loc. cit.

Several determinations for specific gravity were made and the results were found to vary considerably. Examples are 1.00, 1.25, and 1.36. In making these determinations it was desired to obtain the density of the body—that is, glass sponge with included air, rather than that of the glass composing the body.

CHEMICAL CONSTITUENTS

In order to determine whether these bodies represent a special differentiated phase of the magma, with greater silica content and higher viscosity or other particular properties, a partial analysis was made. The writer is indebted to L. J. Youngs, of State College, Pennsylvania, for the following figures:

	Per cent
SiO ₂	49.94
Al ₂ O ₃	13.52
FeO (total iron calculated as ferrous iron)	10.79
TiO ₂	1.26
CaO	9.74
MgO	11.80
Total	97.05

There was a very slight gain on ignition. The alkalies were not determined, owing to scarcity of material on hand, but by difference they would correspond closely with amounts found by other analysts in rocks from this region. The rock is a basalt in composition, and under the microscope the crushed material is found to consist of 70 to 90 per cent of brownish glass, the proportions varying in different bodies, with small phenocrysts of feldspar and pyroxene and possibly some olivine.

A comparison of the analysis with various analyses of Kilauean rocks found in Washington's tables⁶ and made by Silvestri, Phillips, and Lyons shows that the magnesia is very much higher and, as a rule, the lime is considerably higher than for most of these rocks. It compares better with an analysis of Pele's hair made by A. H. Phillips and is similar in most respects to an analysis by Merwin of the lava dipped from the crater Halemaumau.⁷

Although the figures obtained by Mr. Youngs were carefully checked and there seems to be an established high percentage of magnesia, it is not safe to draw conclusions, without further analyses, as to whether a high magnesia content is always characteristic of "Pele's tears." If it be a special feature, then may not their composition have some bearing on the question of the temperature at which this lava becomes liquid and possibly on the origin of these particular bodies? This point is more interesting because of the proportionally high percentage of lime and magnesia in the australites and the high temperature at which they dissolve and pass into the liquid condition. The calcium and magnesium bearing minerals, such as the olivines, pyroxenes, and more calcic feldspars, are always considered among the high-temperature minerals.

⁶ Chemical analyses of igneous rocks. Professional Paper 14, U. S. Geol. Survey.

⁷ Day and Shepard: Water and volcanic activity. Bull. Geol. Soc. Am., vol. 24, 1913, p. 586.

CONCLUSION

The forms of some of "Pele's tears" are so similar to the dumb-bell types of australites as to demonstrate the probability of the latter being of volcanic origin. These bodies of distinctly volcanic origin demonstrate the possibility of bodies with such shapes as spheres, dumb-bells, etcetera, being formed in the atmosphere from a rotating liquid body. There is no evidence in the composition of the lava that a specially viscous type of liquid is necessary to develop these forms; but the magnesia and lime contents are high, and it appears that we must look for some specially favorable condition of temperature, pressure, or other physical circumstance to account for their origin, since they are not, so far as known to the writer, common products of volcanoes. Thus may some special condition also have given rise to the australites.

No attempt is made here to throw fresh light on the distribution of the australites in Australia, although it is the opinion of the writer that they are of volcanic origin and the method of transportation will be discovered as the geology of Australia becomes better known.

TRIASSIC IGNEOUS ROCKS IN THE VICINITY OF GETTYSBURG, PENNSYLVANIA

BY GEORGE W. STOSE AND J. VOLNEY LEWIS

(Abstract)

CONTENTS

	Page
Distribution and mode of occurrence.....	55
Petrography.....	56
General character.....	56
Texture.....	56
Order of crystallization.....	56
Varieties of diabase.....	56

DISTRIBUTION AND MODE OF OCCURRENCE (BY G. W. STOSE)

The igneous rocks are all intrusive. No basalt flows. One main sill crosses the area from southwest to northeast just east of Gettysburg. Average thickness, 2,500 feet. Dips under shales on west at 20°. Irregular outline in places may be due in part to cross-faulting. Average width of outcrop about 1 mile. Local widening to over 2 miles, probably due to local thickening and corresponding displacement of overlying shale.

Two main cross-cutting bodies nearly at right angles to bedding. Larger one, 1 mile thick. Offshoots from these form minor sills and cross-cutting bodies.

A sill along the plane of the flat, overlapping western contact of Triassic sediments on Paleozoic limestone exposed at many places.

Thin dikes and sills are in part later than the large sill and cross-cutting bodies.

The igneous magma seems to have entered the Triassic rocks near their western border; to have spread along the flat, western contact as a sill; to have extended laterally in a thick sheet between the layers as the Gettysburg sill; to have broken across the bedding in several places between these two sills as cross-cutting bodies. Judging from the coarseness and thickness of the

intrusive bodies, the molten rock probably reached the surface, but all traces of the lava have been removed by erosion.

PETROGRAPHY (BY J. V. LEWIS)

GENERAL CHARACTER

Dominantly diabasic. Remarkable diversity of differentiation facies. Range from coarse-grained granitic texture and pink to light or dark gray colors in the larger bodies to dense black rock in thin sheets and dikes and in contact facies.

Chief constituents greenish black pyroxene and whitish to gray plagioclase, the former generally preponderating. Approximately equal at many places and locally the feldspar is in excess. The microscope shows plentiful magnetite and minute apatite crystals and generally some quartz and orthoclase. Locally in darker varieties much hypersthene or olivine or both. In lighter facies quartz and orthoclase abound, chiefly in micrographic intergrowth. Here and there biotite and, far less commonly, titanite.

Pyroxene not uncommonly altered in part to uraltic amphibole or serpentine and chlorite with granular magnetite. Corresponding alteration of feldspars yields fine scaly (apparently sericitic) aggregates and, less commonly, kaolin. Epidote abundant in places.

TEXTURE

Typically diabasic—pyroxene filling angular interstices in a felted ground-mass of slender plagioclase crystals. By coalescence of pyroxene into larger areas, in which feldspars are imbedded, the texture becomes ophitic. Dense varieties grade into typical basalt with glassy ground-mass; some with scattered phenocrysts of pyroxene and, less commonly, feldspar and olivine. In acid facies much quartz and orthoclase, in separate grains or micrographic intergrowth, occupy angular spaces among plagioclase crystals and there is much less pyroxene.

ORDER OF CRYSTALLIZATION

Prevailing diabasic—plagioclase completed before pyroxene. Two marked exceptions: (1) Prismatic pyroxene crystals in some coarser quartz-orthoclase facies with subordinate plagioclase. (2) Pyroxene phenocrysts in dense black dikes, thin sheets, and contact facies, with few large feldspars. This earlier crystallization, probably before intrusion, followed the usual order in plutonic rocks and would have produced a gabbro. In the normal diabase the order has been: (1) apatite, (2) magnetite, (3) olivine, (4) plagioclase, (5) pyroxene, (6) micrographic quartz-orthoclase, (7) orthoclase, (8) quartz.

VARIETIES OF DIABASE

(1) *Normal diabase*, the most common pyroxene-plagioclase rock; (2) *feldspathic diabase*, or *anorthosite*, chiefly plagioclase feldspar; (3) *quartz diabase*, with abundant quartz, chiefly in micrographic intergrowth with orthoclase; (4) *micropgmatite*, mainly micrographic quartz-orthoclase; (5) *aplite*, essentially dense granular quartz-orthoclase rock; (6) *hypersthene diabase*, much hypersthene, replacing in part monoclinic pyroxene; (7) *olivine diabase*, with abundant olivine; (8) *basaltic diabase*, or *basalt*, dense black facies, in

places vesicular and having glassy ground-mass; (9) *olivine basalt*, dense black variety, with abundant olivine.

Presented in abstract extemporaneously by both authors.

DESERT REGOLITH AND ITS GENETIC RELATIONS TO MAXIMUM EPIROTIC DEPOSITION

BY CHARLES KEYES

(Abstract)

It was remarked by many who listened to the admirable illustrated lecture on the "Characteristics of Egyptian deserts," given by Dr. W. F. Hume before the meeting of the Twelfth International Geological Congress in 1913, that it was a very great surprise to learn that arid lands were so dominantly bare-rock plains rather than rolling sand wastes, or tracts deeply mantled by rock debris, as so commonly regarded. That the mantle of decayed rock materials which so widely distinguished most pluvial lands of the globe and which is so aptly denominated the regolith should appear to be so largely wanting in this most famous and typical of deserts was a fact that was directly ascribed to the peculiarities of arid land depletion.

The recognition of the prevalence of bedrock surfaces over broad tracts does not preclude the existence of frequent and often extensive accumulations of rock-waste in the deserts. Enormous amounts of these soil materials are manifestly not only constantly moved about over wide areas of the arid country, but they are directly exported far beyond desert confines. The areas of greatest accumulation of continental or epirotic deposits appear to have a close genetic relation to the areas of greatest arid deflation. Concrete illustrations are drawn from four continents.

Presented by title in the absence of the author.

ORIGIN OF FOLIATION IN THE PRECAMBRIAN ROCKS OF NORTHERN NEW YORK

BY WILLIAM J. MILLER

(Abstract)

It has been generally assumed that the Adirondack Precambrian rocks, including the Grenville strata and the syenite-granite intrusive series, have been severely compressed and folded as well as thoroughly metamorphosed and foliated by the compression.

Evidence will be presented to show that the Grenville strata have never been highly folded or severely compressed. Various broad Grenville belts are known to be only very slightly folded, while many masses, large and small, are merely tilted or domed at various angles, though some local contortions do occur. These structural relations are best explained as having been the result of slow irregular upwelling of the more or less plastic magmas (probably under very moderate compression), whereby the Grenville strata, previously deformed little or none at all, were broken up, tilted, and lifted or domed. Some bodies of strata, caught between batholithic magmas, were locally squeezed or contorted.

The Grenville sediments, which are thoroughly crystallized, were certainly reorganized into new minerals under deep-seated conditions; but, since the strata were never highly compressed, it is evident that they were subjected to essentially static rather than dynamic metamorphism. This explains not only the retention of stratification surfaces to the present time, but also the invariable parallelism of stratification and foliation.

Regarding the foliation of the syenite-granite series, it is believed that during the process of intrusion the magmas were under only moderate lateral pressure, if any; that the process of intrusion was long continued; that the foliation was developed essentially as a flow-structure, under moderate pressure, during the intrusion, and that the almost universal but varied granulation of these rocks was produced mostly by movements in the partially solidified magma, and possibly in part by moderate pressure applied after complete consolidation.

The usual parallelism of both Grenville masses and foliation with adjacent masses and foliation of the syenite-granite intrusives are readily accounted for because the Grenville masses were swung into general parallelism with the slow-moving magmatic currents.

Presented in abstract extemporaneously.

DISCUSSION

Prof. J. E. WOLFF: In the Archean highlands of New Jersey, with a nearly constant northeast striking and easterly dipping foliation and frequent northeast pitching linear structure, it seems necessary to suppose strong lateral compression.

Further remarks were made by Professors R. A. Daly, M. B. Baker, and George H. Chadwick, with reply by the author.

Professor COLEMAN remarked: It seems to me that both sides in this interesting discussion are entirely right, but in different areas. From my own field experience at one point or another I can corroborate the statements made by all who have joined in the discussion. There is no real contradiction between them, since what took place in one region differed from what took place in another.

LANDSLIDES IN UNCONSOLIDATED SEDIMENTS

BY DAVID H. NEWLAND

(Abstract)

The paper discusses landslides as an agency of degradation in regions of low relief and loose sediments like the larger stream valleys in the glaciated district. Occurrences in the terraced Pleistocene clay and sand beds of the Hudson-Champlain Valley are referred to in particular, on account of the number of available observations which extend over a considerable period of time.

Gravity disturbances in bedded clays and sands often occur on small gradients. The materials as a whole possess less stability under varying conditions of moisture content and climate than the unsorted heterogeneous ac-

cumulations of rock weathering that are commonly involved in slides in mountain regions. Their forms are correspondingly varied and complex, in some instances embodying very puzzling mechanical features. The gravity stress which is the fundamental cause of dislocation may be transmitted long distances through the medium of a practically fluid stratum below the zone of rupture, as has not infrequently happened in the Hudson Valley. Unlike the usual condition in mountain forms, there need be no essential variations in the character of the material displaced and the undisturbed beds. Any structural change that could be of significance in the formation of such slides, in the very nature of the case, is scarcely to be looked for, and the same is true also with respect to a slipping surface.

The conditions attendant on the disturbances can generally be determined by observation or by testing the ground in the vicinity, from which some conclusion may be drawn as to the causes leading up to the slides. The exact impetus or proximate causes, however, is seldom to be ascertained. Usually several factors have to be taken into consideration in determining the origin of individual slides, and their relative importance is difficult to estimate. The problem may be further complicated by the entrance of some external influence into the situation, either of natural development or arising from the agency of man.

Of the conditions which govern the form taken by the movement, those of more immediate concern are the nature of the beds—that is, whether clay, sand, or mixture of the two; the moisture content, and the surface contour. The forms that have come under observation in the Hudson Valley are as follows:

1. Surface creep, involving soil, sand, and gravel; little active in plastic clays.
2. Slumping and flows; peculiar to clays and silts.
3. Earth slides; materials of any sort, but not fluent; the movement takes place on the face of slopes that are oversteepened.
4. Subsidence of surface through squeezing out of a wet clay substratum on the plane of its bed.
5. Subsidence of surface from unbalanced pressure on confined liquid substratum, leading to an upward movement at a distance.

The influence of the various kinds of movement on the process of degradation is too important to be left out of account in a region like the Hudson Valley. Their importance, of course, can not be estimated quantitatively, although there is reason to believe that locally they have a predominant part in the work of surface leveling. On the nearly flat tops of the terraces erosion ordinarily is unable to make much headway, especially when the surface is heavily sodded, whereas a very light slope suffices to cause the precipitation of masses of earth in slides, some of which may attain large proportions. There is record of 10 or 12 catastrophic landslides in the Hudson Valley in a period of 75 years; the larger ones involved upward of 100,000 cubic yards of earth. The inconspicuous forms, no doubt, accomplish the largest share of leveling, since they are widely active with cumulative effects.

Presented in abstract extemporaneously.

The Society adjourned at 5.45 o'clock p. m.

ANNUAL DINNER

The annual dinner of the Society was held at Rauscher's, about 227 persons participating. Dr. John M. Clarke acted as toastmaster and the speakers of the evening were Messrs. William N. Rice, H. P. Cushing, Frank D. Adams, Charles D. Walcott, George O. Smith, W. G. Miller, and Joseph Barrell.

SESSION OF WEDNESDAY, DECEMBER 29

The Society convened at 9.15 o'clock a. m., with President Coleman in the chair.

REPORT OF AUDITING COMMITTEE ¹

The Auditing Committee begs to report that they have examined the papers and vouchers of the Treasurer and find them to be correct and in good order.

The investment securities will be examined at a later date.

JOHN E. WOLFF,
GEORGE H. PERKINS,
For the Committee.

The report was accepted.

The printed report of the Council was then taken from the table and, on motion, accepted.

The Society then took up the consideration of scientific papers.

TITLES AND ABSTRACTS OF PAPERS PRESENTED BEFORE THE MORNING
SESSION AND DISCUSSIONS THEREON*FERROUS IRON CONTENT AND MAGNETIC PROPERTIES OF THE NATURAL
OXIDES OF IRON AS AN INDEX TO THEIR ORIGIN AND HISTORY*

BY R. B. SOSMAN AND J. C. HOSTETTER ²

(Abstract)

Practically all natural iron oxide contains more or less ferrous iron, the percentage varying from a few hundredths of 1 per cent up to the percentage found in magnetite. The magnetic susceptibility likewise varies over a very wide range and depends in part on the ferrous iron content. By measuring

¹ Under date of February 19, 1916, Edward K. Mathews reports that, acting as a member of the Auditing Committee of the Society, he examined the Society's securities in the hands of the Treasurer and found them to be as listed in the Treasurer's report under date of December 1, 1915.

² Introduced by Arthur L. Day.

these quantities, both for natural oxides and for oxides made under known conditions in the laboratory, it is possible to draw some conclusions concerning the conditions under which the natural oxides were formed or to which they have been subjected since their formation.

Presented in abstract extemporaneously by the senior author.

VARIABLE COMPOSITION OF MELANOCHALCITE

BY W. F. HUNT AND E. H. KRAUS

(*Abstract*)

Our previous knowledge of this rare mineral has been restricted to that of a single paper by the late Prof. G. A. Koenig. An investigation of recently acquired material shows, however, considerable variation in the chemical composition from that previously reported. Koenig's interpretation of the composition as a basic salt of an ortho-silico-carbonic acid is questioned, and evidence is given for considering melanochalcite as a mechanical mixture of tenorite, malachite, and chrysocolla.

Presented by title in the absence of the authors.

DEFINITION AND DETERMINATION OF THE MINERAL HYDROXIDES OF IRON

BY H. E. MERWIN AND EUGEN POSNJAK

(*Abstract*)

Specimens from a large number of localities were grouped according to optical characteristics. Those found to be sufficiently homogeneous were studied chemically and thermally. Definitions are based on a correlation of characters. Those characteristics most readily determined are emphasized.

Presented in abstract extemporaneously by the senior author.

Brief remarks were made by Prof. J. E. Wolff.

SALINE FUMAROLE DEPOSITS OF THE SOUTH ITALIAN VOLCANOES

BY HENRY S. WASHINGTON

(*Abstract*)

After a brief account of previous work on volcanic salts, the occurrences observed in the summer of 1914 at Vesuvius, Etna, and Vulcano were described. Analyses were given showing the characteristics of the salts found at each volcano. They present certain rather remarkable features—among them the practical absence of chlorides at Vulcano; the occurrences of thiosulphates (no sulphites) within the crater, but not on the outer slope at the same volcano; the prevalence of chlorides, with sulphates, at Etna; the relatively large amount of iron at Vulcano and its paucity at Etna, though the lavas show the converse relations. An interpretation of the results and their application to our views on the constitution of the magma was attempted. The purely

mineralogic description, in collaboration with Dr. H. E. Merwin, will be published later.

Presented in abstract extemporaneously.

CRYSTALS AND CRYSTAL FORCES

BY F. E. WRIGHT

(Abstract)

A general-discussion of crystals as systems of vectorial forces, with special reference to their individuality and their behavior with respect to other systems of forces, special emphasis being placed on the individuality of crystals as it finds expression in the so-called false equilibria of thermodynamics. Methods for the measurement of crystal forces were considered briefly.

Presented in abstract extemporaneously.

EXTENSION OF THE MONTANA PHOSPHATE DEPOSITS NORTHWARD INTO CANADA

BY FRANK D. ADAMS AND WM. J. DICK

(Abstract)

The paper described an investigation undertaken for the Commission of Conservation of Canada for the purpose of ascertaining whether the great phosphate deposits recently discovered in Montana, Idaho, Wyoming, and Utah could be traced northward into Canada. Three lines of section across the Rocky Mountains in Canada were examined and were described. These are those on the North Kootenay Pass, the Crows Nest Pass, and on the main line of the Canadian Pacific Railway. In the most northerly of these sections, crossing the Rocky Mountain National Park at Banff, in the province of Alberta, the phosphate was discovered.

Presented in abstract extemporaneously by the senior author.

DISCUSSION

Mr. L. D. BURLING: The contact between the Devonian and the Cambrian in the North Kootenay Pass section was quite naturally ascribed by Dawson to faulting, but the direct superposition of the two systems described by Professor Adams is strikingly corroborated in three other widely scattered localities: (1) at Elko, British Columbia; (2) in the mountains near Upper Columbia Lake, and (3) along the main line of the Canadian Pacific Railway, just west of Banff, Alberta, where the Sawback formation has been shown to directly underlie the intermediate limestone of the Devonian and to be of Cambrian age; 25 miles to the west, however, the Cambrian is overlain by 10,000 or more feet of Ordovician and Silurian strata.

Of interest also in this connection will be the statement that the small "downfaulted block" which has been described as representing the "Jurassic" in the Canadian Pacific Railway section west of Banff is now believed to

represent an outcrop of the Upper Banff shale in its normal position with respect to the underlying Rocky Mountain quartzite.

The finding of the Albertella fauna in place is not the least interesting of Professor Adams' results. The reference of this elusive drift-block fauna to the Middle Cambrian instead of the Lower Cambrian has been confirmed during the past summer by its discovery in place in the type section on Mount Bosworth, British Columbia, some 700 feet above the base of the Cathedral limestone. The definite placement of this fauna is of value in connection with the discussion of the Precambrian age of the rocks of the Galton and Purcell series, since the latter are now known to be older than the Albertella fauna. That they are still older is evidenced by Dawson's early and hitherto unrecorded discovery of Lower Cambrian fossils in the Kootenay Valley south of Upper Columbia Lake.

Brief remarks were made by Prof. Alfred C. Lane, with reply by the author.

EMERALD DEPOSITS OF MUZO, COLOMBIA

BY JOSEPH E. POGUE

(Abstract)

The paper is a result of a field study of the deposits made in July, 1915. The geological and mineralogical relations of the emerald are discussed and evidence presented to show that the emerald originated as a result of gas-aqueous emanations from an intrusion that is not exposed, but is indicated by the presence of contact rocks, pegmatites, and a significant mineral association.

Presented in abstract extemporaneously.

CRYSTALLINE MARBLES OF ALABAMA

BY WM. F. PROUTY

(Abstract)

The crystalline marbles of Alabama are largely confined to an area about 35 miles long and $1\frac{1}{4}$ miles in maximum width. This area is for the most part a fault block, with the strike of its rocks in some places similar to and in other places differing from both that of the Ocoee phyllite, which bounds it on the southeast side throughout the length of the field, and the Knox dolomite formation, which bounds the field for the larger part of the distance on the northwestern side. Topographically the area is a well defined valley, except locally, where it is crossed by ridges of dolomite or more resistant rock.

Although there is no direct fossil evidence to indicate the age of the marble, the general characteristics lead to the conclusion that it varies in age from Cambrian to Ordovician in different parts of the field.

The thickest deposits of marble are in the central and southwestern parts of the field, and it is here that the chief developments of the area are being made at the present time.

The quarry methods employed in some of the openings are considerably

different from the usual methods in such work, the object being to take advantage of the unusual structural conditions.

The Alabama marble is medium to fine grained, with distinctly interlocking crystals. It is unusually translucent, sonorous, and resistant to abrasion and weathering. These qualities, together with its warm coloring, explain its growing popularity in the market.

A careful study of the structural conditions in different parts of the field show numerous examples of drag folding and, locally, schistosity in the marble.

The sharp line of demarcation between calcite and dolomite beds is suggestive of a distinct differentiation in the original sediments.

Presented by title in the absence of the author.

ORISKANY IRON ORE

BY R. J. HOLDEN

(Abstract)

This ore occurs in quantity only in Virginia. Its chief development has a definite stratigraphic position at the top of the Lewistown limestone. The ore bodies have thicknesses up to 40 feet and extend on the strike for distances of a few hundred feet to half a mile and down the dip for several hundred feet. Various theories of its origin have been given, but mining has shown that the ore is secondary, and it now seems certain that the most productive phase of the ore is a limestone replacement, the iron being derived from the overlying shale.

Presented in abstract extemporaneously.

GEOLOGIC MAP OF THE FORT HALL INDIAN RESERVATION

BY GEORGE R. MANSFIELD

(Abstract)

The sedimentary rocks of the Fort Hall Indian Reservation include representatives of all the great Paleozoic and later systems except the Cretaceous. Igneous rocks, mainly extrusives, are present in great abundance and considerable variety, including a single occurrence of nepheline basalt. Some of the Triassic and Jurassic rocks are so well developed as to require new subdivision. The structure is quite complex, with both folding and faulting, especially the latter, and there is an important thrust-fault, the Putnam overthrust. Three epochs of deformation and four of igneous activity have been recognized. There may be others. Important deposits of phosphate occur in the eastern part of the reservation.

Presented by title.

*PRELIMINARY GEOLOGIC MAP OF THE WAYAN QUADRANGLE, IDAHO-
WYOMING*

BY GEORGE R. MANSFIELD

(Abstract)

The rocks of the Wayan quadrangle include some basalt and a long sequence of sedimentary formations representing all the great Paleozoic and later systems except the Cambrian and Silurian. The strata have been complexly folded and faulted. The most noteworthy structural features are the great bifurcated syncline that occupies the central portion of the quadrangle, bending off toward the northwest, and the Bannock overthrust that traverses the quadrangle in an irregular course. There are many minor folds and faults. Besides structural problems, there are many of stratigraphic and physiographic interest. The quadrangle is of great economic importance because of the large body of high-grade phosphate rock that it contains.

Presented by title.

*GLACIAL LAKES AND OTHER GLACIAL FEATURES OF THE CENTRAL
ADIRONDACKS*

BY HAROLD L. ALLING¹

(Abstract)

Although the glacial geology of the foothills of the Adirondacks has been investigated, the central area, specially the Mount Marcy, Lake Placid, Ausable, and Elizabethtown quadrangles, has not yet received the attention it deserves. Here Pleistocene phenomena are beautifully shown in great abundance. The most interesting features are two series of local glacial lakes, or water levels, ranging from 2,000 down to 500 feet in altitude through a dozen successive stages. They are indicated by deltas, terraces, strong beaches, and channels. In several of the outlets beautiful cataract plunge basins are exhibited. The post-lacustrine deformation of a number of the levels has been determined and indicates that the amount of uplift decreases in passing to the lakes of lower altitude.

Presented by title.

*PLEISTOCENE FEATURES IN THE SCHENECTADY-SARATOGA-GLENS FALLS
SECTION OF THE HUDSON VALLEY*

BY HERMAN L. FAIRCHILD

(Abstract)

A map shows (1) the deep flooding by the sealevel waters of the Hudson Valley; (2) the vast areas of detrital plains; (3) the glacial indrainage, especially the Iromohawk; (4) the ice-block kettles of Saratoga and Round lakes. The relation of the Iromohawk distributary channels to the Round Lake

¹ Introduced by H. L. Fairchild.

kettle shows that a large portion of the drift-buried ice block, which produced the basin of Round Lake, persisted until the locality had been raised 175 feet, the total Pleistocene uplift being 375 feet.

Differential uplift diverted the Iromohawk flow from its northward course, as mapped by J. H. Stoller, into the present Mohawk channel.

Presented in abstract extemporaneously.

PLEISTOCENE UPLIFT OF NEW YORK AND ADJACENT TERRITORY

BY HERMAN L. FAIRCHILD

(Abstract)

The paper published in the Bulletin, volume 25, pages 219-242, was intended to emphasize the fact of deep submergence in sealevel waters of the Connecticut and Hudson valleys. With further study of the marine plane and with precise levels on the international boundary, it is now possible to locate with approach to accuracy the isobases of land uplift across New York and the adjoining areas, east and west.

A fixed vertical relation in the Ontario basin between the Iroquois and the marine planes give us a key, for the Iroquois area, to (1) the amount of post-Iroquois uplift; (2) of Iroquois or glacial uplift; (3) of initial altitude; (4) of local warping; (5) of flooding, due to differential uplift of the Rome outlet.

The uplifting of the area, in time and amount, shows very close relation to the latest ice-body, and the uplifting appears to have been a progressive wave, subsequent to the far removal of the ice.

Presented in abstract extemporaneously.

Published in full in this volume.

DISCUSSION ON THE TWO PRECEDING PAPERS

Prof. J. W. SPENCER: Some years ago I devoted much time in the study of the terraces of New England, but only published a note on them. From the highlands of northern New England extensive terraces, sometimes 20 miles long, were found to occur in all the great valleys, with slopes much less than the gradients of the rivers. These terraces descended not in warpings, but by steps (overlapping one another); not merely to the south in the Connecticut Valley, but westward down the Lamoille, eastward down the Saco, northward in the valleys leading to the Saint Lawrence. Consequently, if they were due to the rise of the mountain mass, there should be some agreement between the terraces of the different valleys. If such had been the case, the movements were per saltum in steps and not by warping curves. This does not support the hypothesis of Professor Fairchild, that the Connecticut terraces record the warping. Also long ago I found that the Iroquois beach extended east of Watertown, and correctly traced it to East Pitcairn; but in 1902 Professor Fairchild terminated late Iroquois some 20 miles within the explored zone, as shown on his map.

Prof. R. D. SALISBURY: We can congratulate ourselves on the fact that Professor Fairchild has been able so long to follow up his studies on glacial

drainage and associated problems, and that he can now present to us the ripe conclusion of his long and careful study. The generalizations based on such detailed and long-continued work are the generalizations which we have come to trust. The fact that his conclusions tie up harmoniously the conclusions of many individual workers who have studied local areas intensively is gratifying and seems to be good evidence of the soundness of the results at which Professor Fairchild has arrived. I have but one comment to add. I think the east end of isobase O will have to be shifted somewhat farther south.

Professor Fairchild replied briefly to Professor Spencer's and Salisbury's remarks.

STUDIES OF GLACIATION IN THE WHITE MOUNTAINS OF NEW HAMPSHIRE

BY JAMES WALTER GOLDTHWAIT

(Abstract)

The field studied in 1915 is the northwestern part of the White Mountains, more particularly the Ammonoosuc Valley between Littleton and Bretton Woods. It is the purpose of the paper to show that glacial phenomena, which were regarded by Louis Agassiz and by Charles H. Hitchcock as records of local mountain glaciers, and by Warren Upham as records of a local White Mountain ice-cap, remaining at the close of the last Glacial epoch, are in reality records of the North American ice-sheet, which retired into Canada without leaving either mountain glaciers or a local ice-cap in its wake. Striae, dispersion of boulders, terminal moraines, outwash plains and kame terraces, mapped in detail, are offered as evidence.

Observations on the Mount Washington Range, the Franconia Mountains, and Mount Mooslaue support the view advanced in 1912, that there were cirque cutting glaciers in the White Mountains at a time prior to the last regional glaciation, but that these glaciers were small and not very numerous.

Presented by title in the absence of the author.

GLACIATION AND STORMY PERIOD OF THE FOURTEENTH CENTURY

BY ELLSWORTH HUNTINGTON

(Abstract)

A recent study of the salt lakes at the eastern base of the Sierra Nevadas shows that their later strands can be dated in terms of the growth of the big trees on the other side of the mountains. Gale has shown that chemical evidence indicates that Owens Lake must have overflowed not more than "4,000 years ago or considerably less." Jones has shown the same to be true of Pyramid Lake. Since they did not use the entire drainage area and made no allowance for increased solution of mineral matter from the rocks with increased precipitation, it seems necessary to reduce the time to approximately 2,000 years. In that period Owens Lake has decreased to 40 per cent of its former size, and Pyramid has similarly shrunk, although not so much. Below the old outlet strands there is in each case a series of younger strands whose

height and relative age agree with the fluctuations in the growth of the sequoia trees 50 miles west of Owens Lake. The most notable strand appears to date from the fourteenth century—a time at which the big trees made a peculiarly rapid growth. It is remarkable for its size and strength. It could have been formed only under the influence of unusually high winds. What seems to be the same strand can also be recognized at Mono and Pyramid lakes, both by its relative location and its evidences of peculiarly strong wave action.

The fourteenth century period was also marked by climatic stress in other regions. In central Asia the Caspian Sea and Lop-Nor expanded with great rapidity, as is known from well authenticated historic records. In north-western Europe storms of unusual severity afflicted the countries around the North Sea. Floods were of frequent occurrence, and extraordinarily cold winters caused the Baltic Sea to be frozen over completely. Norway suffered great economic distress because of persistent failure of the crops. In England the rains were so abundant that the average production of wheat per acre fell off one-third and the agricultural population was in great distress. In Iceland and Greenland similar occurrences took place. Petterson holds that increased severity of climate and the crowding down of the ice were the cause of the final abandonment of Greenland by the Norsemen.

It is noteworthy that in both the eastern and western hemispheres the chief evidences of climatic stress come from the semi-arid and desert regions, where a subtropical climate prevails, or else from the northern border of the belt of cyclonic storms. These are the regions where the Glacial period also produced the most noteworthy results, either by the expansion of salt lakes or by the production of ice-sheets. The conditions in the fourteenth century seem to have been of essentially the same nature as those of the Glacial period, the only apparent difference being in degree. It is possible that a study of such climatic fluctuations during historic times may lead to a final solution of the problem of the nature and cause of Glacial periods.

Presented in abstract extemporaneously.

PLEISTOCENE DEPOSITS OF MINNESOTA AND ADJACENT DISTRICTS

BY FRANK LEVERETT

(Abstract)

The extension of glacial investigations into Minnesota from districts farther east has shown that in the last, or Wisconsin, stage of glaciation the Labrador ice-sheet reached its culmination and began to wane before the ice which spread over northern Wisconsin and neighboring parts of Minnesota reached its culmination, and that this ice in turn began to wane before the Keewatin ice-sheet, which lay still farther west, reached its culmination. It is suggested in explanation of this westward wave of ice culmination that the highland of Labrador was the natural starting point of glaciation, and that because of storms coming to it from the southwest the ice-sheet grew westward to such an extent as to eventually receive more snowfall in the region south of Hudson Bay than in the Labrador district; so that there resulted a waning of ice movement from the latter district. Still later the ice-sheet grew westward into central Canada and caused the culmination of the Keewatin ice movement which spread into Minnesota and Iowa.

The same sort of westward growth of the ice-sheet may have taken place in the Illinoian stage of glaciation, for it now appears probable that the Illinoian drift from the Labrador center is somewhat older than drift which was carried southward from the region south of Hudson Bay through the Lake Michigan basin into Illinois, and also older than the Iowan drift brought in from districts still farther west.

This interpretation, if correct, may do away with some of the difficulty hitherto found in explaining the development of an ice-sheet in the low area west of Hudson Bay.

Presented in abstract extemporaneously.

Society adjourned at 12.45 o'clock p. m.

AFTERNOON SESSION

The Society met at 2.10 o'clock p. m., with Mr. F. B. Taylor in the chair.

RESOLUTION REGARDING THE TAKING OF EXPERT TESTIMONY

The Secretary *pro tem.* read a resolution adopted by the Council at its noon meeting as follows:

"In view of the fact that there is almost a universal desire among scientific men for some sort of reform in the manner of presenting expert opinion in legal procedure, and in view of the fact that the geologists of the country are certainly as deeply interested in this matter as any other scientific body, the following resolutions are presented for consideration:

"*Resolved*, That the Geological Society of America recognizes the urgent need of reform in the methods of securing evidence or expert opinion in judicial procedure;

"That the Geological Society of America approve the efforts of the American Association for the Advancement of Science in this behalf, and

"That the Council of the Society is hereby authorized and directed to cooperate with the Committee of the American Association for the Advancement of Science in an endeavor to bring about such a reform."

On motion, the resolution was adopted by the Society.

From 2.10 to 2.30 o'clock p. m. the Society met in joint session with the Paleontological Society and listened to the address of Dr. E. O. Ulrich, President of that Society, on "The use of fossils in correlation."

The Society then proceeded to the consideration of scientific papers.

Many of the attending geologists took advantage of the invitation of the Geophysical Laboratory to visit that institution during the afternoon. Arrangements were made by the staff of the laboratory to exhibit the work of the different departments, and a most instructive afternoon was spent by those interested in the geophysical and petrographic lines.

TITLES AND ABSTRACTS OF PAPERS PRESENTED BEFORE THE AFTERNOON
SESSION AND DISCUSSIONS THEREON*PENNSYLVANIAN OF TENNESSEE*

BY L. C. GLENN

(Abstract)

The paper described briefly the formations into which the Pennsylvanian is divided in the State, gave their areal distribution, discussed the principal coals in each, and attempted to correlate these coals. Brief consideration was given to the age of the deposits found in the State.

Presented in abstract extemporaneously.

*SUBDIVISIONS OF THE THAYNES LIMESTONE AND NUGGET SANDSTONE,
MESOZOIC, IN THE FORT HALL INDIAN RESERVATION, IDAHO*

BY GEORGE R. MANSFIELD

(Abstract)

A mineral examination of the Fort Hall Indian Reservation by a Geological Survey party in 1913 involved a detailed mapping of some of the Mesozoic formations. It proved desirable to subdivide the Thaynes limestone, Lower Triassic, and the Nugget sandstone, Jurassic or Triassic. The thickness of these strata, including the intervening Ankareh sandstone, is about 6,800 feet.

The Thaynes limestone was raised to the rank of a group consisting of three formations—the Ross limestone at the base, the Fort Hall formation, and the Portneuf limestone. The Ankareh, which at the type locality is a shale, was here found to be a sandstone. The Nugget sandstone was subdivided into four members—the Higham grit at the base, the Deadman limestone, the Wood shale, and the main sandstone member.

The paper described briefly the formations subdivided, explained the use of the names, and included a discussion by G. H. Girty of the interesting faunas of the three formations of the Thaynes group.

Presented in abstract extemporaneously.

*STRATIGRAPHY OF SOME FORMATIONS HITHERTO CALLED BECKWITH AND
BEAR RIVER, IN SOUTHEASTERN IDAHO*

BY GEORGE R. MANSFIELD AND P. V. BOUNDY

(Abstract)

In the Montpelier and Wayan 30-minute quadrangles of southeastern Idaho, parties of the Geological Survey have found great thicknesses of strata, aggregating 17,000 feet or more, that have hitherto been assigned to the Beckwith and Bear River formations. On the maps of the Hayden Surveys both forma-

tions are included in the Laramie. The Beckwith has been assigned to the Cretaceous or Jurassic and the Bear River to the Upper Cretaceous.

There is a considerable lack of agreement, both lithologically and faunally, between the formations in the quadrangles named and the Beckwith and Bear River formations in their type localities. The discrepancy is so great that it now seems inadvisable to continue the use of the names Beckwith and Bear River in the district discussed. Three groups of strata are recognized, the lowest of which is marine Jurassic, and rests unconformably on the Twin Creek limestone, the main Jurassic formation of the region. The two higher groups are non-marine and probably Lower Cretaceous. They are separated from each other by an unconformity, but the lower group appears to be conformable on the Jurassic beds below. The two higher groups have some resemblances to the Kootenai of Montana and Canada, but the data are at present insufficient for their correlation with that formation. No characteristic Bear River fossils have been found in the district, though such have been found farther north, and there is a possibility that the doubtful beds may grade upward into the true Bear River in that direction.

The beds formerly called Beckwith are divided into seven formations and a new name is given to the strata hitherto called Bear River. The paper gave a statement of the stratigraphic problems involved, together with a description of the formations.

Presented in abstract extemporaneously by the senior author.

SEDIMENTATION ALONG THE GULF COAST OF THE UNITED STATES

BY E. W. SHAW

(Abstract)

Sedimentation along the Gulf Coast of the United States proceeds in three general and markedly different ways, each of which prevails over a large area. On the west coast the sediment delivered to the sea by streams is being re-worked, some of it many times over, but is not carried far away. At the mouths of the Mississippi silt, clay, and fine sand are accumulating, layer on layer, almost precisely where dropped by the river. Along the Florida coast comparatively little sediment is carried into the sea by streams, and the littoral deposits consist largely of very clean sand and calcium carbonate extracted from sea-water by invertebrates, algæ, bacteria, etcetera.

The lagoon and barrier beach conditions which prevail along the west coast and a part of the north coast are perhaps the most common, and those at the mouths of the Mississippi most unique, though the processes in operation at the southern end of Florida differ in some respects from any in operation elsewhere in the world.

The object of the paper was to compare and contrast individual processes and results affecting each region. On shore and off shore samples of sediment from each of the three general regions were exhibited.

Presented in abstract extemporaneously.

RELATIVE AGE OF THE DETROIT RIVER SERIES

BY CLINTON R. STAUFFER

The Detroit River series is that part of the so-called Monroe formation which lies above the Sylvania sandstone. In an article on the "Nomenclature and subdivisions of the Upper Siluric strata of Michigan, Ohio, and Western New York," by Lane, Prosser, Sherzer, and Grabau,¹ this series was subdivided as follows:

Upper Monroe or Detroit River Series	{ Lucas dolomite Amherstburg dolomite Anderdon limestone Flat Rock dolomite
---	--

The series is overlain by the Onondaga (Columbus or Dundee) limestone or the lowest generally recognized Devonian of the region. Doctor Prosser originally defined the Lucas limestone as the upper portion of the Monroe and states that "it includes all the rocks between the top of the Sylvania sandstone and the base of the formation which Doctor Lane in Michigan has named the Dundee limestone."² It is thus evident that the Lucas has been much restricted in the later paper. However, it seems probable that the Upper Monroe, or Detroit River series, includes beds not very well known at the time the definition of the Lucas limestone was written. There is not only some doubt as to the stratigraphic order of these subdivisions, but very conflicting opinions regarding the real age of the whole series. It may be that part of the confusion lies in the fact that the faunas of these subdivisions are not entirely distinct, and that they have been mistaken for each other in the various outcrops. Only a small portion of any of the faunas of the series is as yet known.

The Detroit River series is rather widely distributed over Michigan, Ontario, Ohio, and doubtless Indiana as well. In Ohio the Anderdon outcrops in the quarries at Castalia; the Amherstburg probably occurs at Fremont, certainly at West Liberty, and the Lucas outcrops at numerous places along the margin of the Onondaga (Columbus), in northwestern Ohio. It is especially well shown and quite fossiliferous at Silica and Centennial, in Lucas County. In Ontario the most extensively distributed division of this series is probably the Amherstburg, at least so far as at present known. In addition to those along the Detroit River, outcrops of this dolomite may be found near Woodstock, Saint Marys, Wingham, Formosa, and McRae Point. The Lucas dolomite is less perfectly known, but appears to be represented in the outcrops at Kincardine and perhaps the upper part of the outcrop above Beachville. The Anderdon is not certainly known in Ontario, outside of Essex County, although it is quite well developed in Monroe County, Michigan. Probably the best development of the series is along the Detroit River, but it is by no means confined to that region. The salt shaft at Oakwood,³ near Detroit, cut one of the best and most complete sections through it that is thus far known, excepting, of course, the numerous well records. But the most accessible sections are those that have been exposed in the vicinity of Amherstburg, Ontario.

¹ Bull. Geol. Soc. Am., vol. 19, 1907, p. 556.

² Jour. Geol., vol. xi, 1903, pp. 540-541.

³ Mich. Geol. and Biol. Survey, Pub. 12, Geol. ser. 9, 1911, fig. 21, opp. p. 278.

FIGURE 1.—THE UNEVEN UPPER SURFACE OF THE ANDERDON LIMESTONE IN THE AMHERSTBURG STONE COMPANY'S QUARRY

FIGURE 2.—ROUGH UPPER SURFACE OF THE ANDERDON LIMESTONE, SHOWING JOINTS WIDENED BY SOLUTION AND LATER FILLED WITH ONONDAGA MUD

THE ANDERDON SURFACE AT AMHERSTBURG, ONTARIO

The Flat Rock dolomite is not exposed at Amherstburg, unless the lowest beds of the Amherstburg Stone Company's quarry belong to it; but the other divisions are well shown. These are all more or less fossiliferous and at certain places have yielded an abundant fauna. All divisions of the Detroit River series contain Devonian faunal elements. This is most marked in the Amherstburg dolomite, and probably least in the Lucas dolomite. The Amherstburg dolomite is typically developed in the bottom of Detroit River at Amherstburg, Ontario, and within the last seven or eight years this has been excellently exposed in the dry cut of the Livingston Channel. Near the north end of this cut the rocks are very fossiliferous and nearly all of the abundant forms are not only of marked Devonian aspect, but resemble decidedly the Onondaga fauna. Some of the most striking resemblances are to be found among the species of *Cladopora*, *Cystiphyllum*, *Favosites*, *Hederella*, *Romingeria*, *Synaptophyllum*, *Syringopora*, *Crania*, *Meristella*, *Pentamerella*, *Productella*, *Reticularia*, *Rhipidomella*, *Schizophoria*, *Spirifer*, *Stropheodonta*, *Conocardium*, *Modiomorpha*, *Paracyclas*, *Schizodus*, *Bellerophon*, *Callonema*, *Eotomaria*, *Loxonema*, *Platyceras*, *Tentaculites*, *Ryticeras*, *Dawsonoceras*, *Proetus*, etcetera. Such an aggregation of genera would alone be sufficient to demand comparison with the Onondaga fauna, but in many cases even the similarity of the species is so close as to have caused their identification with the Onondaga forms. Certainly one can have no quarrel with the man who insists that this fauna must be placed within the Devonian system. It is, in fact, a Devonian fauna. But the massive layers of brown dolomite containing this fauna are overlain, toward the south end of the cut, by similar beds carrying the Lucas fauna. A close study of this latter fauna, which is the one that has usually been considered to have especially strong Silurian affinities, shows that it is different from those to which it has been compared, and that the majority of its species, that have been considered Silurian, were really described from the Lucas dolomite of Michigan or from the Ohio outcrops of this same formation; 34 per cent of the species listed by Grabau and Sherzer⁴ are also found in undisputed Silurian deposits; 9 per cent occur in typical lower Devonian, and 57 per cent are apparently not known outside of the Detroit River series. While the Lucas fauna at first seems impossible as a Devonian aggregation, this latter consideration decidedly minimizes the relationship to other known Silurian faunas.

In northwestern Ohio the Onondaga (Columbus) limestone rests unconformably on the Lucas dolomite, but at Amherstburg and at the Sibley quarry, nearly straight across the river in Michigan, it rests on the Anderdon limestone. The dip of the rocks at the Amherstburg Stone Company's quarry is to the west-southwest, although well records show that it evidently is reversed at no great distance. A similar dip occurs at the Sibley quarry across the river. In the dry cut of the Livingston Channel the dip is south at the rate of about 100 feet per mile, thus seeming to indicate the rising axis of an anticline of which the limbs slope away to east and west. This has led some men to the belief that the Anderdon belongs immediately under the Onondaga, as it has usually been found in surface outcrop. Also very serious doubt exists in the minds of some as to whether there is a stratigraphic break or an erosion surface at the top of the Anderdon limestone or between the Onondaga and

⁴ Mich. Geol. and Biol. Survey, Pub. 2, Geol. ser. 1, 1910, pp. 211-213.

the Detroit River series. The question involved is not confined to the Detroit River region alone, but is contingent on observed conditions over the whole southwestern part of Ontario, as well as much of Ohio, and which conditions are undoubtedly similar to those in other adjoining States to the west. Near the eastern end of Lake Erie the relation of the Onondaga to the Silurian is one of unconformity (disconformity). This is well illustrated in the outcrops at Buffalo and to the westward in Ontario. There are, however, two distinct erosion periods represented at or near the base of the Onondaga. One of these preceded and the other followed the Oriskany sandstone.⁵ It is probable that it was the latter of these that scattered the sands of Oriskany origin over a much wider region than that originally covered by the formation itself. This is indicated by the fact that none of the sandy remnants at the base of the Onondaga, west of North Cayuga township, in Ontario, carry even the slightest trace of Oriskany fossils. At most places a well developed basal conglomerate may be found in the lowest Onondaga beds, but occasionally it is rather imperfectly formed or sometimes entirely wanting. Even in such cases, however, not one of the Ontario contacts, and few, if any, of those in Ohio thus far examined, is without abundant evidence of the erosion period that intervened between the formations in contact. Near Springvale this is abundantly shown by the reworked beds of Oriskany, that now contain an Onondaga fauna, and the widened cracks in the pre-Oriskany dolomites, which are now filled with sand to a depth of several feet below the contact. At Goderich, Ontario, the basal conglomerate of the Onondaga limestone is especially well developed. Pebbles of the Upper Monroe as large as a man's fist are mingled with Onondaga corals, brachiopods, trilobites, etcetera, in an arenaceous limestone which rests on an uneven surface of Monroe. In the early days of the Goderich salt industry Mr. Attrill drilled a test well near the Lake Huron shore, across the Maitland River from Goderich. In reporting on the results of this experiment, Dr. T. Sterry Hunt says: "We now come to the consideration of an unexpected result of the examination of the cores from the Goderich boring, namely, the occurrence beneath 278 feet of beds, chiefly dolomite, which, according to the Geological Survey, underlie the Corniferous (Onondaga) limestone of the region, of not less than 276 feet, chiefly of gray, non-magnesian, coralline limestone, abounding in chert and seeming like a repetition of the Corniferous (Onondaga). Beneath this lower fossiliferous limestone, it will be noted, are dolomites with gypsum, succeeded by variegated marls, with an aggregate thickness of not less than 364 feet before reaching the saliferous strata, which latter have been penetrated 520 feet without reaching the underlying Guelph formation. Prof. James Hall, who has kindly examined such specimens of the corals as I have obtained from the limestone, recognizes in them two species of *Favosites*, *Favosites winchelli* and *Favosites commonsi*, together with a section of *Acervularia* or *Diphyphyllum*."⁶ This fossiliferous horizon is undoubtedly a part of the Detroit River series and very probably includes the Amherstburg dolomite and the Anderdon, as these beds outcrop about 35 miles farther north. The interesting point here is that it lies 278 feet below the base of the Onondaga, and that this latter has a well

⁵ Bull. Geol. Soc. Am., vol. 23, 1912, p. 373; also Geol. Survey Canada, Memoir 34, 1915, p. 60 and pl. iii.

⁶ Geol. Survey of Canada, Rept. Prog. for 1876-1877 (1878), p. 242.

FIGURE 1.—JOINTS IN THE ANDERDON LIMESTONE WIDENED BY SOLUTION AND FILLED WITH ONONDAGA MUD

FIGURE 2.—INTERSECTING JOINTS IN THE ANDERDON LIMESTONE, SHOWING LATER FILLING WITH ONONDAGA MUD

PRE-ONONDAGA JOINTING AT AMHERSTBURG, ONTARIO

developed basal conglomerate in which pebbles of the Detroit River series are mingled with sand and Onondaga fossils and which rests on an uneven surface of Detroit River series. Certainly there can be no doubt but that at this place the time interval between the fossiliferous Detroit River series and the overlying Onondaga was very long.

The most important consideration, however, is the condition at Amherstburg, since it is there that doubt has been expressed regarding the occurrence of an extensive unconformity (disconformity) at the base of the Onondaga. At Amherstburg the development of the basal conglomerate is weak; in fact, various geologists to whom specimens from this horizon were submitted expressed grave doubt as to the real conglomeratic nature of the deposit, although admitting the intermingling of the Anderdon and Onondaga limestones at the contact. This basal layer of the Onondaga is composed of more or less angular fragments of the compact drab Anderdon intermingled with the rather porous brown magnesian Onondaga, and in which there is sometimes so much sand that a thin layer might be called a real sandstone. In addition to the unevenness (see plate 7, figure 1), the upper surface of the Anderdon is rough and uneven. This roughness is attributable to differential weathering, since recently weathered surfaces of the Anderdon show the same pitted surface, although the rock is fresh at a small fraction of an inch below the surface. In some of these irregularities considerable accumulations of sand occur and fragments of the Onondaga often cling to these irregularities after the overlying layer has been removed. The Anderdon is affected by a system of joints which do not pass upward into the Onondaga. Some of these joints have been widened by solution to three or more inches before the Onondaga was deposited, and then into these cracks the Onondaga muds, intermingled with sand, filtered as the sea advanced over the region (see plates 7, 8, 9). In some cases this sand has penetrated several feet below the contact and may now be found in cavities among the Anderdon fossils. The upper surface of the Anderdon is often pretty well covered with rather large, low-spined gastropods and somewhat similar cephalopods (see plate 9). These are almost always of the brownish gray Onondaga limestone, which has led some one to suggest that the underlying mud must have been soft when the Onondaga sea advanced over it; and at which time these gastropods and cephalopods were pressed into this soft mud. Unfortunately these fossils are poorly preserved and have not been satisfactorily determined. Grabau and Sherzer say that the gastropod is probably *Trochonema ovoides*,¹ but they do not even mention the cephalopod. Possibly it is the *Trochoceras anderdonense* which they have described. The indications are that these gastropods and cephalopods were already fossil when the Onondaga sea advanced over the region; in fact, they were molds of the exterior and interior of the shells into which the Onondaga mud was pressed. Frequently it is possible to find casts of the shelly portion made up of the Onondaga material, which contrasts strongly with the other, while the internal mold of the Anderdon material is still retained and forms a perfect core for the fossil. Many of the infillings of these gastropod and cephalopod molds are in large part sand. One or two somewhat similar fossils were found in the basal Onondaga above the contact, but they were so poorly preserved that it was impossible to make out whether

¹ Mich. Geol. and Biol. Survey, Pub. 2, Geol. ser. 1, 1909, p. 44.

they were in reality the same forms. These may have been internal molds loosened from the Anderdon and incorporated into the Onondaga at the time the latter was being deposited.

There is thus an accumulation of positive evidence demonstrating the existence of a period of erosion of long duration at the basal contact of the Onondaga in the Amherstburg quarry, and very similar evidences may be found on the Michigan side of the river. It is, of course, impossible to say positively that the sand at the contact represents the Oriskany in this region, remote from any known deposits of that formation, or that it is even material derived from the erosion of an Oriskany deposit, as is certainly the case farther northeast in Ontario. It is equally possible that the sand at Amherstburg was derived from outcrops of the Sylvania, subjected to erosion in pre-Onondaga time, since the Onondaga (Dundee) limestone is in contact with the Sylvania sandstone at the National Silica Company's quarry, 7 miles northwest of Monroe, Michigan,⁸ and hence may have a similar relation at other near-by localities. It should be pointed out also that the Onondaga of extreme southwestern Ontario, like the Columbus limestone of northern Ohio, probably does not represent the whole of the formation as developed in western New York. In Ontario, immediately across the Niagara River from Buffalo, where the Onondaga limestone is probably the exact equivalent of the same deposit in New York, the lower layers are characterized by the relative absence of corals and the abundance of brachiopods. Some of the most characteristic fossils of this horizon are *Amphigenia elongata*, *Anophia nucleata*, *Anoplothea camilla*, *Centronella glansfagca*, *Chonetes hemisphericus*, *Cypriocardinia indenta*, *Platyceras dentalium*, and numerous others that are abundant at higher horizons. The beds carrying these species have been traced across the province and are last found near the shore of Lake Huron to the south of Port Elgin. These species are not found in the Onondaga at Goderich or at Amherstburg; neither are they found in the outcrops on the islands of Lake Erie and at Marblehead. The probability is that the beds which should contain them were never deposited in those regions, and that the lowest Onondaga is wanting. Some of these species reappear in the Columbus limestone of central Ohio, where the Onondaga fauna is again more like that of western New York.

It is therefore evident that the time interval represented by the unconformity (disconformity) between the Anderdon limestone and the Onondaga was a long one. If we may trust the record of the Goderich well, as described by Hunt, and the excellent section of the Oakwood salt shaft, the stratigraphic order as given by Sherzer and Grabau cannot be disputed. The period of erosion at Amherstburg therefore removed all the Lucas and Amherstburg dolomites and lasted through as much of the Onondaga as is represented by the lower zone of that formation at the eastern end of Lake Erie. The particular division of the Detroit River series which shows most marked Middle Devonian faunal characters therefore preceded its derivative fauna, the Onondaga, by the Lucas dolomite interval and this long erosion period. If the arenaceous material at the base of the Onondaga at Amherstburg and near-by localities in Michigan represents the Oriskany horizon, as believed by Sherzer

⁸ W. H. Sherzer and A. W. Grabau: Geol. and Biol. Survey Michigan, Pub. 2, Geol. ser. 1, 1909 (1910), p. 40.

FIGURE 1.—THE SURFACE OF THE ANDERDON LIMESTONE, SHOWING MOLDS AND JOINTS FILLED WITH ONONDAGA MUD

FIGURE 2.—UPPER SURFACE OF THE ANDERDON LIMESTONE, SHOWING ONONDAGA CASTS IN THE ANDERDON MOLDS

FOSSILIFEROUS SURFACES OF ANDERDON AT AMHERSTBURG, ONTARIO

and Grabau,⁹ the Amherstburg dolomite can not very well be considered younger than Helderbergian. But since the basal Onondaga at Amherstburg does not carry the oldest fauna belonging to that formation, there is still room for doubt that the arenaceous contact is that representing the Oriskany farther to the east in Ontario and New York.

As is well known and as already pointed out above, the Oriskany sandstone itself rests unconformably on the Silurian (Cobbleskill and Rondout), at the eastern end of Lake Erie. It may be that the unconformity (disconformity) at the base of the Oriskany in western New York represents the one occurring at the base of the Detroit River series, and that the unconformity (disconformity) at the top of the Oriskany is the one so prominent at the base of the Onondaga in Ohio, Michigan, and extreme southwestern Ontario. If this be the case, these two unconformities run together at various places in Ontario near the eastern end of Lake Erie, but diverge rapidly to the westward from Springvale. There is thus represented an unknown interval during which it is probable the Detroit River series was deposited; for to maintain that such a fauna as that found in the Amherstburg dolomite is Silurian is more impossible than to find a place for it among the recognized Devonian formations. While the fauna of the Detroit River series is not altogether unlike that of the Helderbergian, it differs so markedly from it that it can hardly have lived in that sea which spread westward over New York and even into Illinois during Lower Devonian. The occurrence of some of these same fossils in the dolomites of Alberta, as shown by a small collection from the headwaters of the north fork of the Saskatchewan River,¹⁰ suggests that there may have been another embayment from the north or northwest during early Devonian with a very different fauna from that found in the Helderbergian of the Atlantic embayment, although in part contemporaneous with it. The fact that a fauna similar to that of the Detroit River series, especially in its Mollusca, and to a less extent in the Anthozoa and Brachiopoda, occurs in the lowest Devonian of the Ural Mountain region of Russia¹¹ lends some support to this suggestion. Even in the *Idiostroma* limestone of the Cedar Valley of Iowa there is a suggestion of the recurrence, at a much later date, of the coral and hydrozoan reefs of the Anderdon.

Presented in abstract extemporaneously.

DISCUSSION

Dr. A. C. LANE: Over a large area there is no trouble in drawing the line between Devonian and Silurian, as there is such an unconformity that the Onondaga (Dundee) comes almost directly, separated only by the Oriskany sandstone from the Salina or Lower Monroe. But near Detroit River we find that, while there remains an unconformity, or rather disconformity, under the Dundee, the Oriskany seems also to be almost or quite continuous as a transgression sandstone with the Sylvania sandstone. If we turn to Schuchert's

⁹ Mich. Geol. and Biol. Survey, Pub. 2, Geol. ser. 1, 1909 (1910), p. 46.

¹⁰ A. W. Grabau and W. H. Sherzer: Mich. Geol. and Biol. Survey, Pub. 2, Geol. ser. 1, 1909 (1910), pp. 102 and 116.

¹¹ Th. Tschernyschew: Mémoires du Comité Géologique, vols. iii, 1885, and iv, 1893.

Von N. Lebedew: Idem., vol. xvii, 1902, tables opp. p. 132.

curves we see that between the marked depression of the middle of the (Upper) Silurian and the Devonian there was a long period of relatively low sealevel or high stand of the continent, not due to any disturbance in Michigan, but perhaps due to disturbances in the other hemisphere. The question is whether the base of the Devonian shall be drawn at the beginning or the end of this time of uplift. If we assume, as seems likely, that the beginning is the best line, the most sharply marked, and the one most nearly coeval throughout the world, then the Detroit River series would be Devonian.

We have, however, an alternation of a fauna so Devonian that it was at first taken for Hamilton (Traverse) and then a recurrence of Silurian fauna. But this Silurian fauna is of a local peculiar type—that associated with brines—that may easily be assumed to have lingered on in some Dead Sea or Aral Sea. The real difficulty is that in New York we have a Lower Devonian fauna of a quite different type. We must either assume these to have formed during the unconformity, so well described by Stauffer, or assume that New York and Michigan were in different provinces for a length of time that seems unlikely. But the division line between Devonian and Silurian must depend on diastrophic studies outside of Michigan.

RECESSION OF NIAGARA FALLS REMEASURED IN 1914

BY J. W. SPENCER

(Abstract)

In October, 1914, just ten years after my previous survey, which had been the fifth (the fourth was that of Kibbe, in 1890), I remeasured the crest-line of the main cataract of Niagara. The principal changes were not at, but adjacent to, the apex. Here had been a narrow V-shaped incision in the upper strata only, which has widened to one of U-shape 50 feet broad. To the west blocks of rock 400 feet long and from zero to 55 feet wide had fallen, thus straightening the line. The fallen area here and elsewhere aggregated about two-thirds of an acre, corresponding to a mean recession for the full width of the falls of scarcely 2.5 feet a year. The diversion from the river had been more or less compensated by the late prevailing high water. This reduces the mean annual rate of recession during 72 years to 4 feet a year, compared with 4.2 feet during the previous 62 years, but adds 5 per cent to the computed age of the falls.

The eastern end of the great cataract at the time of Hennepin (1678) has been investigated, the location of the western end having been previously found. Thus it appears that the mean rate of recession during 236 years was approximately 3.75 feet a year. This factor would increase the calculated age of the fall by 10 per cent, which figure still comes within the limit originally provided for ($39,000 \pm 4,000$ years), but on the side of increased age.

Additional soundings in the upper part of the gorge were also made. In no case did I find depth of 100 feet, in contrast with those of 186 to 192 feet a short distance below. These confirm the evidence of a late reduction in the height of the falls, as previously described, and also mentioned by Kalm in 1750. The reduction in height was due to the falling of the gorge walls at the Whirlpool Rapids, thereby damming and raising the river level above them.

I made another sounding (of 210 feet) near that of 183 feet, previously de-

scribed, in the channel just beyond the end of the gorge. This shows that the level in Lake Ontario was, after the birth of the falls, even lower than I had previously announced.

These are new facts leading to further precision, besides confirming earlier results.

Presented in abstract extemporaneously.

TERRESTRIAL STABILITY OF THE GREAT LAKE REGION

BY J. W. SPENCER

(Abstract)

Among the results obtained by studying the lower lake terraces is one relating to the cessation of earth-movements. The terraces in question are those most strongly developed at or near the mouths of tributary streams, recurring all the way from the head of the lake to below the outlet of Lake Ontario, at 12 to 20 feet and 2 to 5 feet above its level. In them no deformation is determinable, in contrast with the differential movement of 540 feet between the head of the lake and Parishville, New York, as seen in the Iroquois beach, with tilting varying from 2 to 6 feet per mile. Having previously proved from the daily fluctuations of lake levels that there has been absolutely no earth-movement since 1854, these terrace features show that the cessation dates back a long time. The movements were in operation after the diversion of the Huron drainage to that of Lake Erie, which I have placed at 3,500 to 4,000 years ago. At present this would give near the date for their cessation.

Presented in abstract extemporaneously.

SCOUR OF THE SAINT LAWRENCE RIVER AND LOWERING OF LAKE ONTARIO

BY J. W. SPENCER

(Abstract)

I am not aware of previous investigations on this problem. When the waters of the Ontario basin were subsiding from the earlier glacial lake, the Saint Lawrence River channel did not come into existence until they fell to a level of not more than 20 feet above the present stage. This figure represents the total lowering of the lake due to the river scour acting on its general bed of drift. At the upper rapids the river has exposed very little rock. Thus at the first, or Gallops Rapids, the river flows over beds of boulders in two of its branches, while in the third the rock has been channeled for only a quarter of a mile (in Ordovician strata). These features show the relative youthfulness of the modern Saint Lawrence River. The present shoreline of the lake was examined for evidence of deformation, but neither in it nor the lower terraces could such be found.

To one who had primarily observed the coast of the lake at Hamilton and Toronto, where great beaches occur, he would be surprised to find how widespread has been the cutting away of the shores or the small development of the beach. It is much less developed than the Iroquois. Indeed, the interven-

ing terraces do not suggest such long pauses as were required for the building of the Iroquois beach. So far no time measurement is of any value. The most important problem is the date when the waters fell below the Iroquois beach, as this would most nearly coincide with the last days of the Glacial period. But these new data show the newness of both the Saint Lawrence and the modern shores of Lake Ontario.

Presented in abstract extemporaneously.

PLEISTOCENE DRAINAGE CHANGES IN WESTERN NORTH DAKOTA

BY ARTHUR G. LEONARD

(Abstract)

The continental ice-sheet produced important drainage changes in western North Dakota. Its effects are particularly well shown in the case of the Missouri, Yellowstone, and Little Missouri rivers, since all these streams were forced to seek new channels.

The southerly course of the Missouri River below old Fort Stevenson has been attributed to the later or Wisconsin ice-sheet, but evidence is presented that the valley is preglacial. This evidence is based on the presence of glacial boulders and drift on the valley bottom and at many points on a terrace representing a former floodplain of the Missouri. The river must have flowed in its broad, terraced valley at the time of the earlier ice-invasion, and on the floor of this valley the glacier deposited boulders and till. When the ice-sheet invaded the region, it blocked the valleys of both the Missouri and Yellowstone rivers, and also the preglacial valley of the Little Missouri, forcing these streams to seek new channels. Lakes were formed in the valleys of the Little Missouri and Yellowstone rivers, the waters rising until they overflowed the divide to the Knife River. The combined waters of the three rivers flowed east across Dunn County and southeast across Morton County to the mouth of the Cannon Ball River. The length of this Pleistocene valley of the Yellowstone and Missouri rivers, which extends from the head of the Knife to the mouth of the Cannon Ball and crosses the divides between the Knife and Heart rivers and that between the Heart and Cannon Ball, is 155 miles.

The lower 50 miles of the Yellowstone Valley was blocked with ice during the Glacial period, and the waters flowed east to the valley of the Little Missouri, forming at least two broad, flat-bottomed valleys connecting these streams. One of these valleys, 28 miles long, is now occupied in part by Bennie Pierre Creek, a tributary of the Yellowstone.

The Little Missouri was also forced out of its preglacial valley, which is now occupied by Cherry and Tobacco Garden creeks. It probably flowed for a time through the Pleistocene valley of the Missouri and Yellowstone rivers, but later took an easterly course and formed its present postglacial valley, which extends from the mouth of Bowling Creek to the Missouri River—a distance of 100 miles. Evidence is given that this lower valley of the Little Missouri is much younger than the portion above the mouth of Bowling Creek.

Read by title in the absence of the author.

LANDSLIPS AND LAMINATED LAKE CLAYS IN THE BASIN OF LAKE BASCOM

BY FRANK B. TAYLOR

(Abstract)

Lake Bascom was a sprawling glacial lake which filled the valley of Hoosick River and its branches in northwestern Massachusetts, southwestern Vermont, and adjacent parts of New York. The lake was 500 feet deep near Williamstown and slightly deeper toward Pownal. Many landslips have occurred around the sides of the main deep part of the lake and a few in ravines which held narrow bays. Laminated pebbleless lake clay and silt covers a considerable part of the lake floor around Williamstown and eastward toward North Adams, and it also occurs between North Adams and Briggsville and northward and at Petersburg and near North Pownal. During some of the later, lower stages of the lake, pebbleless clay and silt were deposited near Hoosick, around Hoosick Falls and North Hoosick and near North Bennington. Other laminated clays on the Owl Kill and on the Hoosick below Eagle Bridge are not related to landslips.

At its highest level Lake Bascom stood at an altitude of 1,125 or 1,130 feet above sealevel. With three or four exceptions, all the landslips observed in this region occur in the basin of Lake Bascom at some distance below its highest level. Some of the largest slips are very old, surely Pleistocene in age. In all probability they occurred soon after the fall of the lake waters. They are now represented by a series of steplike benches at certain places on the hillsides. Other slips are more recent and some are still in active movement. In one place they are just beginning, the bluff back of the edge of a previous slip being riven by deep cracks.

Nearly all of these landslips are related to underlying beds of laminated pebbleless lake clay and silt. The slips generally occurred on steep slopes where a stream was cutting at the bottom and where the underlying beds at or above the stream level were lake clay and silt. Some, however, are not related to any stream. The weakness of these beds, increased by softening in seasons of prolonged rain, caused a yielding and a slipping down of the heavy overlying masses. In some slips the laminated beds are partly exposed in vertical sections in the bluff at the back of the slipped mass; in others the clay beds are exposed in the banks or bed of the stream, where they are generally distorted.

A few landslips of a different character, perhaps more appropriately called landslides, occurred recently (August 20, 1901) on the east face of Mount Greylock. These were much smaller masses than some of the slips in Lake Bascom, but the amount of their descent was much greater, being in one case 1,500 feet. These slides occurred after a prolonged wet period and have no relation to lake clays.

Presented in abstract extemporaneously.

DISCUSSION

Mr. GEORGE C. MARTIN stated that Lake Bascom was evidently very similar to some of the marginal lakes on the borders of the existing Alaskan piedmont

glaciers. These lakes are not only filled with icebergs, but in one case¹ a lake is known to contain a large arm of the glacier, which floats on the surface of the lake and forms the larger part of its area. The occurrence of till overlying the clays in the former bed of Lake Bascom may possibly be explained as the residue from the melting of a large amount of heavily debris-laden glacial ice that floated on the surface of the lake and that was left stranded on the lacustrine deposits when the lake was drained. The presence of many bergs and possibly of a floating glacial arm in Lake Bascom may also explain in part the poor development of beaches.

Further remarks were made by Messrs. M. B. Baker and H. F. Cleland, with reply by the author.

TYPES OF LOESS IN THE MISSISSIPPI VALLEY

BY B. SHIMEK

(Abstract)

The gray loess and two types of yellow loess of the Upper Mississippi Valley and the red and yellow loesses of the South are discussed with reference to geographical distribution, differences in structure, color, and fossils, and the significance of these differences. The conclusion is drawn that these types indicate in part differences in time of deposition and in part differences in source of materials. Incidental references are made to deposits which resemble loess and have been so classed.

Presented by title in the absence of the author.

Society adjourned at 5.20 o'clock p. m.

PRESIDENTIAL ADDRESS

At 8 o'clock p. m., at the Cosmos Club, Prof. A. P. Coleman delivered his address as retiring President, his topic being

DRY LAND IN GEOLOGY

Published as pages 175-192 of this volume.

The address was followed by the complimentary smoker given in honor of the Geological Society of America and the Paleontological Society by the local members of the former organization. The address of the President was the most largely attended session of the meeting, about 300 persons being present.

¹ G. C. Martin: Geology and mineral resources of the Controller Bay region, Alaska. U. S. Geol. Survey, Bull. 335, 1908, pp. 47-48, pl. vii.

SESSION OF THURSDAY, DECEMBER 30

The Society convened at 9.15 o'clock a. m., with President Coleman in the chair.

The Society proceeded immediately to the consideration of scientific papers.

TITLES AND ABSTRACTS OF PAPERS PRESENTED BEFORE THE MORNING
SESSION AND DISCUSSIONS THEREON*STAGES IN THE GEOLOGIC HISTORY OF PORTO RICO*

BY CHESTER A. REEDS

(Abstract)

In March, 1915, Prof. C. P. Berkey in his report, "Geological Reconnaissance of Porto Rico," Annals New York Academy of Sciences, stated that an early Tertiary peneplain separated an "older" from a "younger" series of formations in Porto Rico. Observations made in the field during the following June and July suggest that more recent erosion cycles and periods of uplift and subsidence are also in evidence, and that they are responsible, for the most part, for the present configuration of the island. The positive and negative oscillations of the strand-line were induced in all probability by diastrophic movements within the island mass and adjacent sea-floor.

Previous to the early Tertiary peneplanation beds of shale, limestone, conglomerate, and volcanic rocks were intruded by igneous masses, producing a basal complex of folded and faulted strata and igneous intrusions. Following this peneplanation no igneous intrusion or ejectments have occurred, for none were observed in the basal lignitic shales and overlying chalky and cross-bedded limestones of the younger series. Folding and faulting are, however, in evidence in certain places in the beds of the "younger" series.

Evidences of marine transgression are to be seen in the marginal coastal plains and the wave-cut escarpments. Suggestions of an earlier transgression are to be seen in the even-crested cliffs and headlands on certain coasts and the isolated hills, ridges, and plateaus of near-by tracts. This older plain is typically developed in the district to the south and west of Yauco.

While successive transgressions and recessions of the sea were affecting the coastal and marginal tracts, subaerial denudation was busily sculpturing the present mature topographic features of the interior out of the exposed portion of the older warped peneplain and accompanying monadnocks. Features which indicate that stream erosion has been varied are to be found in the abandoned erosion levels in valley walls and at water gaps on rivers like the Descalabrado and the Jacaguas. At one stage the lower courses of these rivers were filled with sand and gravel. At the present time these sediments are being eroded and at some points the stream bed was observed to be 50 to 60 feet below the base of the gravel deposits.

The stages in the geologic history of Porto Rico were varied in length and kind of work performed.

Approximately 10,000 fossil specimens were obtained from the "older" and "younger" series of formations. In certain localities algæ, foraminifera, and corals were collected from the limestone members in the "older" series. They suggest Cretaceous age. The larger portion of the collection, however, was

STAGE	ELEVATION	EROSION	SUBSIDENCE	DEPOSITION	COMMENTS
XIV		Delta and Coastal Terraces		Coastal Barriers, Dune Sand, San Juan Formation River Flood Plains	Present
XIII	Slight Uplift	Rejuvenescence of Rivers			Partial Withdrawal of Sea from Platform
XII				Alluvial and Coastal Conglomerates, Sands and Shales	
XI			Marginal Transgression of the Sea		3 ^d Cycle of Erosion incomplete
X		Inauguration of present Upland Dissection & Inception of present River-Valleys			
IX	Uplift				Tilting Sedimentary Series, Folding & Faulting
VIII		Development of Peneplain			2 ^d Cycle of Erosion incomplete
VII	Uplift				Tilting Sedimentary Series, Folding & Faulting
VI				Local Deposition of green Shales and Limestones	Long Duration
V			Subsidence affecting greater Portion of Island		Unconformity between older & younger Series
IV		Development of Peneplain		Deposition beyond Edge of present Platform	1 st Cycle of Erosion
III	Uplift				Injected Porphyries, etc. affecting both Igneous & Sedimentary Series
II				Local Deposition of Conglomerates, Black Shales, interbedded Turfs and Breccias, and Limestones on submerged Platform	Long Duration
I	Initial Volcanic Cones				

Chester A. Reed, December 1915

FIGURE 1.—Stages in the geologic History of Porto Rico

gathered from the lignitic shales and white limestone of the "younger" series. The sea-urchins collected from these beds indicate late Eocene or early Oligocene age. It would be premature to assign ages to the various formations and stages in the geologic history of Porto Rico before this large collection of fossils has been worked up.

Presented in abstract extemporaneously.

DISCUSSION

MR. EDWIN T. HODGE: The "older series" in Porto Rico has long been looked on as non-fossiliferous, and therefore of undeterminate age. Last summer, during my investigation in the south central portion of the island, I was fortunate enough to find some fragmentary fossils, which, it is true, do not constitute a whole flora or fauna, yet are sufficient to establish with certainty the following statements:

At the crest of the Sierra Cayey, the rocks of which lie near the base of the series, I found the calises of two *Cladophyllia furcifera* characteristic of the

Edwards of the Gulf. An adjacent bed of hematite contained some fossil leaves. These I have submitted to Doctor Berry and Doctor Knowlton, who tell me that while they are not critical, yet they tend to substantiate the evidence of the corals.

A few miles farther south and higher in the stratigraphic series, as worked out by purely structural methods, I found the lower portion of a *Venaricardia alticosta* index for the Chickasawan and Claibornian of the Gulf.

The Cretaceous and Eocene strata in which these fossils were found have been folded to high angles—as much as 90° in places—and peneplained. The evidence of this peneplanation is decisive. It is shown by the even heights and flat crests of the mountains, the discordant relation existing between the drainage and the structure, which shows superimposition, and by the presence of deep residual soils.

On this peneplain was laid down the “later limestones.” From fossils collected in Porto Rico by Doctor Berkey and studied by Miss O’Connell, we know these to be of Oligocene age.

Thus in Porto Rico we have positive proof that the Eocene is separated from the Oligocene by a period of time during which there was uplift and intense folding, followed by long erosion. This is entirely a new fact to science and throws light on the small thickness and extent of the Oligocene, both in the eastern United States and Porto Rico. It follows that the Oligocene present is probably very late Oligocene.

Doctor REEDS replied that as the collection was very large nothing definite could be said at present regarding the age of the beds.

CRETACEOUS OF ALBERTA, CANADA

BY JOSEPH H. SINCLAIR¹

(Abstract)

Recent explorations by the author in the foothills of southern Alberta, Canada, have resulted in the identification of a very complete Benton fauna and accurate measurements of 4,500 feet of Cretaceous sediments.

The application of the Missouri River succession of Cretaceous rocks to the foothill district is questioned in several horizons, and Stebinger is borne out in his prophecy that the Claggett does not exist there as a lithologic unit.

The presence of the Bearpaw is also uncertain, although it may exist as a thin, brackish-water formation containing coal seams.

The adoption of the name “Dakota” for a thick formation lying between the Kootanie and the Benton is questioned on the ground that a not sufficiently great variation exists between the few specimens found in it by Cairnes and the flora of the Kootanie formation.

While the adoption of the section of the Missouri by Dawson may appear correct farther east in the Great Plains, it appears that continental conditions have pinched out certain marine phases of the eastern section, in the foothills of Alberta. The unsatisfactory line of demarcation between the Cretaceous and the Tertiary is also shown. Certainly no lithologic break is seen and the

¹ Introduced by Charles P. Berkey and A. W. Grabau.

lack of fossils makes a paleontological differentiation difficult. The total thickness of the Cretaceous appears to be 7,000 feet.

Presented in abstract extemporaneously.

SEDIMENTARY SUCCESSION IN SOUTHERN NEW MEXICO

BY N. H. DARTON

(Abstract)

In connection with the study of the Red Beds and associated strata in New Mexico many stratigraphic data have been obtained of the entire Paleozoic and Mesozoic succession. The distribution and local variations of the formations present some novel features which throw light on the geologic history of the region. Especially notable are the overlap relations of the formations of Carboniferous age, particularly the Pennsylvanian division, which overlaps Cambrian, Ordovician, Silurian, Devonian, and Mississippian rocks. The relations of the Red Beds of Permo-Pennsylvanian and Triassic age were studied in detail. Considerable new evidence was also obtained on the distribution of the Comanche group and overlying formations of Cretaceous age.

Read by title in the absence of the author.

*DIVISIONS AND CORRELATIONS OF THE DUNKARD SERIES OF OHIO **

BY CLINTON R. STAUFFER

The youngest Paleozoic deposits in Ohio have long been known as the Upper Barren Coal Measures and more recently as the Dunkard series. These rocks are quite extensively developed in adjacent States to the east and have doubtless been removed by erosion over wide areas in which no trace of them now remains. It is only the comparatively thin western edge of the Dunkard that extends into Ohio and covers a rather narrow strip along the Ohio River from Jefferson to Meigs county. At the northern end of the Ohio portion of the Dunkard basin there is no appreciable break between the Monongahela and the overlying Dunkard series. From the stratigraphic relations the basal plant beds (Cassville) ought therefore to continue the same flora that flourished during the formation of the preceding Waynesburg coal bed, but apparently such is not the case. Over the southern half of the basin, however, the Waynesburg sandstone usually rests directly on the Monongahela with marked unconformity, the Cassville, the Waynesburg coal, and a portion of the underlying shales usually being absent. Unconformities in a series of rocks, such as the Dunkard, probably do not have any very great significance; in fact, they occur at several horizons within the series; but the development of the coarse massive Waynesburg sandstone, often a true conglomerate, over much of the unconformity between Monongahela and Dunkard may be indicative of changed conditions.

The Dunkard series of Ohio may be divided, as in Pennsylvania and West Virginia, into the Washington and the Greene formations. The former begins

* Published with the permission of the State Geologist of Ohio.

with the shales above the Waynesburg coal and ends with the Upper Washington limestone; the latter includes the remainder of the series. This division of the Dunkard series was originally suggested by J. J. Stevenson,¹ although, according to his definition, the Waynesburg sandstone and the underlying plant shales are included with the Monongahela series. The usage herein suggested for Ohio has been approved and used by the United States Geological Survey in the various Pennsylvania folios.² Each of the so-called Dunkard formations is made up of a great number of more or less distinct divisions, which are often traceable over very considerable areas and most of which have been given definite locality names by the Pennsylvania and West Virginia Geological Surveys. Many of these can be recognized in Ohio and form the basis of such stratigraphic correlations as is possible between the various parts of the Dunkard basin. This division of the Dunkard into two formations is very arbitrary, as the stratigraphical or even the lithological break at the horizon used is not pronounced. It does, however, mark the highest level at which marine or brackish-water fossils were found and probably represents the approximate close of the oscillations between land and marine conditions, and introduces the purely land and fresh-water deposits in the Dunkard basin.

The Washington formation varies in thickness from 230 feet in Belmont County to more than 300 feet in Washington County, while to the south of Marietta, where the massive sandstones are well developed, this division of the Dunkard includes 286 feet above the Washington coal, hence is probably at least 386 feet in thickness. The Greene formation reaches a maximum thickness of about 250 feet, although it is usually much thinner and often is a mere capping to the higher hills. Because of its limited distribution and its position in the hills, its character is less perfectly known than is that of the lower formation. Much of the topography of the region which it occupies consists of quite gentle slopes, which are covered with a deep soil and often well sodded.

From Marietta southward the dividing line between the Washington and Greene formations is hard to trace because of the absence of the Washington limestones, but it may still be continued by use of the Jollytown coal horizon or the base of the Jollytown sandstone, which probably marks the break between the two divisions. This sandstone stratum may be picked up at various places to the south of Marietta and has occasionally been quarried for grindstones.

A large part of the Dunkard of Ohio is to be classed as "Red Beds," although the Monongahela series and even the Conemaugh are not without their red shales, which in the Monongahela are often so like those of the Dunkard as to make them easily confused if it were not for other well defined strata associated therewith. There are but few really red sandstones, and those are usually only coated red on the outside or weathered surface, in the Ohio Dunkard. The red is thus almost confined to the shales. In the northern part of the Dunkard covered area the red beds are to be found chiefly in the Greene formation, but to the southward most of the shale in the whole series is red. In the main, these shales, sandstones, limestones, and beds of coal represent land and swamp or fresh-water deposits, but the presence of gypsum

¹ Second Geol. Survey Penna., Rept. K, 1875 (1876), pp. 34-56.

² See U. S. Geol. Survey Folios Nos. 121, 180, etcetera.

in certain of the shales and sandstones, and again marine or brackish-water fossils in other beds, indicates that these conditions at times gave place to others of a very different character.

The Dunkard series as a whole is not very fossiliferous; in fact, it is almost as barren of the identifiable traces of life as it is of the workable coal seams, which originally suggested the term "Upper Barren Measures" for this deposit. In addition to the occasional plant fragment that may be found in almost any part of the series, there are certain rather well defined horizons in the Ohio Dunkard which have yielded important fossils. Plants are, of course, of first importance. Their remains are occasionally to be found in the roof shales of any of the coal seams or even in beds of argillaceous shale and sandstone. Almost any outcrop of limestone may be found to contain small fresh-water gastropods and ostracodes. The Middle and Upper Washington limestones often contain fish plates and teeth, some of which are referable to sharks, and are therefore probably marine. A *Lingula* occurs in the shales associated with the Washington coal. The lowest shales of the series are sometimes a black carbonaceous mass associated with a hard limestone, and these beds contain scales, teeth, and coprolites, all of which are probably fish remains. The most important find of the whole fossil collection, however, was made in the red shales of the Washington formation in the vicinity of Elba and Marietta. At the former of these places, near the base of the Dunkard, amphibian coprolites were found in relative abundance. These are remarkably similar to those found in the Permian of the Western States. At the latter place, during the past summer, fragments of a neural spine of *Edaphosaurus* were found in the sandstones associated with the red shales just above the Lower Marietta sandstone. The remains of this reptile have never before been found in the United States outside of Oklahoma, Texas, and New Mexico. The importance of this find must be very evident, since it agrees with the earlier conclusions drawn from identifications of the Dunkard flora and proves the age of the Dunkard to be identical with the Permian or Permocarboniferous of Texas. After having seen the whole vertebrate collection, Dr. S. W. Williston says that "of the fishes I recognize teeth like those of *Diplodus* from the Texas Permian, but this type runs through the Pennsylvanian and is not characteristic. The Elasmobranch spine is unlike any that I have seen in Texas. The coprolites can not be distinguished from those commonly found in Texas and New Mexico. . . . The *Edaphosaurus* spine is unquestionable, small as it is. The range of the family in Texas is both Wichita and Clear Fork. It occurs in New Mexico in the El Cobre beds, which the accumulated evidence now places as the equivalent of the lower Texas beds (Wichita). . . . In Europe *Edaphosaurus* occurs in the uppermost Carboniferous of Kuonova and the Rothliegende of Saxony."³

In view of this evidence of the Vertebrate fossils, there can be no doubt that the lower portion of the Dunkard series is the equivalent of the lower Texas beds (Wichita) which overlies the Cisco, and that in all probability both beds are Permian.

Presented in abstract extemporaneously.

Brief remarks were made by Prof. A. C. Lane.

³ Letter of December 15, 1915.

*SILURIAN SYSTEM OF MARYLAND*¹

BY C. K. SWARTZ AND W. F. PROUTY

(Abstract)

The authors have been engaged in the study of the Silurian system of Maryland for several years and this paper presented a synopsis of their chief results.

The various formations that constitute the Silurian system of Maryland were discussed, including their lithology, faunas, and stratigraphic features. The formations were correlated with those of the same age in New York and other parts of the Appalachian basin. The problem presented by the Silurian Red Beds was considered briefly and certain analogies between them and the Catskill formation were pointed out.

Presented in abstract extemporaneously by the senior author.

DISCUSSION

MR. G. W. STOSE: The stratigraphy presented by Mr. Swartz in the diagrams is no doubt more accurate and final than that in my report on the eastern part of the area about Hancock, Maryland. The interfingering of red sediments toward the east, extending to lower horizons than the base of the Wills Creek, is of special interest and will require careful study before final conclusions of nomenclature and correlation are reached.

In regard to the faunal relations of the Kiefer sandstone, I do not feel competent to speak; but Mr. Ulrich, who was responsible for having it made a member of the McKenzie formation in my report, has, in a recent table, placed it as a member of the Clinton formation, thus agreeing with Mr. Swartz.

MR. GEORGE H. CHADWICK: Objection is made to the inclusion of the Rochester in the Clinton. The distinctive Rochester fauna is widespread and the facts presented in this paper emphasize the undesirability of submerging it in the very different assemblage lying beneath it, against the plain evidence that the Rochester fauna endures with little modification into the Lockport.

HOMOCLINE AND MONOCLINE

BY REGINALD A. DALY

CONTENTS

	Page
Definition of "homocline".....	89
Different definitions of "monocline".....	90
Favored definition of "monocline".....	91
Summary.....	92

DEFINITION OF "HOMOCLINE"

Workers in deformed strata have long felt the need of a general name for a mass of bedded rocks all of which dip in the same direction. Such a mass may be a tilted fault-block, a series of tilted fault-blocks, a combination of

¹ Presented by permission of the Director of the Indiana Geological Survey.

overthrust blocks, one limb of anticline or syncline, an appressed fold, a series of appressed folds, or a local, single flexure in a pile of otherwise horizontal beds. In many an instance the observer is long delayed in reaching a conclusion as to which of these or other categories a group of inclined beds is to be finally referred; after years of research he may not be able to make this decision. Yet in his field notes and reconnaissance report, if not in his final publication, he should have a broad term by which to refer to the structural unit here described.

Feeling this need during years of reconnaissance and detailed studies in the Canadian Cordillera and elsewhere, the writer has suggested the name "homocline" to cover the case.¹ A "homocline" is thus "any block or mass of bedded rocks all dipping in the same direction." According to one of its several definitions, "monocline" has exactly the meaning given to "homocline," and it is necessary to show reason for preferring the new term.

DIFFERENT DEFINITIONS OF "MONOCLINE"

"Monoclinal" was invented by W. B. and H. D. Rogers, who wrote: "We propose the term *monoclinal* to signify a sameness in the direction of dip, and shall term a mountain or valley in which such sameness prevails a *monoclinal* mountain, or *monoclinal* valley."² Since sameness of dip, rather than the singleness of dip direction, is emphasized, it appears that "homoclinal" would have been etymologically preferable. Though the originators of the word, W. B. and H. D. Rogers made very little use of it, and then almost, if not quite, exclusively in the adjective form. Some authors have since employed "monoclinal" as a noun, equivalent to the proposed "homocline," signifying merely a series of beds dipping in the same direction; but "monocline" in this sense has been comparatively little used.

In 1865 Page used "monoclinal" as a noun, meaning a one-limb flexure, or the same as "monoclinal fold," Powell's name for well known flexures of the Colorado plateaus.³ In 1880 Dutton adopted this usage.⁴ In 1882 Dutton used "monocline" with the same meaning, and also in 1882 A. Geikie published what seems to be the first formal definition of "monocline." He expressed it in the following words:

"Curvature occasionally shows itself among horizontal or gently inclined strata in the form of an abrupt inclination, and then an immediate resumption of the previous flat or gently sloping character. The strata are thus bent up and continue on the other side of the fold at a higher level. Such bends are called monoclines, or monoclinal folds, because they present only one fold or one-half of a fold, instead of the two in an arch or trough."⁵

The Dutton-Geikie usage represents a considerable narrowing of meaning when compared with the "monocline" implied in the Rogers' use of the adjective "monoclinal." The same narrower definition of "monocline" is to be

¹ R. A. Daly: Memoir 68, Geol. Survey of Canada, 1915, p. 53.

² W. B. and H. D. Rogers: Reports of the Association of American Geologists and Naturalists, 1840-1842, Boston, 1843, p. 485.

³ D. Page: Handbook of Geological Terms, Edinburgh and London, 1865, p. 312.

J. W. Powell: Colorado River of the West, Washington, 1875, fig. 67, opp. p. 183.

⁴ C. E. Dutton: Report on the geology of the high plateaus of Utah, 1880, p. 26.

⁵ A. Geikie: Textbook of Geology, London, 1882, p. 515.

found in all three of the succeeding editions of A. Geikie's textbook, in J. Geikie's *Structural and Field Geology* (1905), in the *Standard Dictionary* (1895), in Webster's *New International Dictionary* (1910), Murray's *New English Dictionary* (1908), in Norton's *Elements of Geology* (n. d., 1905?), Scott's *Introduction to Geology* (1911), in Blackwelder and Barrows' *Elements of Geology* (n. d., 1911?), and in Tarr and Martin's *College Physiography* (1914).

Several authorities define the adjective "monoclinal" in the sense of W. B. and H. D. Rogers—"dipping in one direction"—while defining "monocline" in the way adopted by Dutton and Geikie. Examples are to be seen in the *Century Dictionary* (1895) and in Murray's *New English Dictionary* (1908). J. D. Dana at first defined both "monoclinal" and "monocline" in the Rogers sense, but in the last edition of his *Manual of Geology* restricted the meaning of "monocline" as Dutton and Geikie had done.*

Among recent writers who have retained both "monoclinal" and "monocline" in the broad Rogers sense are: R. S. Tarr (*Elementary Geology*, 1897), Chamberlin and Salisbury (*Geology*, Volume I, second edition, 1906), Anderson and Pack (*Bulletin 603*, United States Geological Survey, 1915), and R. H. Johnson (*Science*, volume 42, 1915, page 450).

FAVORED DEFINITION OF "MONOCLINE"

It is safe to conclude that the majority of influential authorities writing since 1880 have favored the Dutton-Geikie definition of "monocline." So far as the writer has been able to find out, Dutton had the priority in the use of this word in geology, while A. Geikie seems to be the first to have given it formal definition. Therein lies one reason for retaining this usage.

A second and stronger reason is found in the need for the word in just the sense implied by Dutton and expressed by Geikie. For the observer working in Arizona, Dutton's "monocline" is as necessary as "anticline" is for the worker in Pennsylvania. In general, the geologists who are interested in a simple, ultimately complete system of names for all types of folds may well favor the Dutton-Geikie usage. The use of the noun in that sense makes obviously desirable a similar narrowing of the adjective "monoclinal." This seems to be a compelling argument for disregarding priority in connection with "monoclinal" and for re-defining it so as to cover one instead of many kinds of structure in bedded rocks.

French and German authors have long used the word "flexure" (*Flexur*) to signify "monocline" in the Dutton-Geikie sense. This usage was introduced by Suess and followed by Richthofen, de Margerie and Heim, Reyer, Neumayr, Credner, de Lapparent, and Haug. It is, however, impossible for writers of English to limit "flexure" in this way. The word has too long meant "folding" in general, or simply "fold," as in the writings of the Geikies, Prestwich, Jukes Brown, W. B. and H. D. Rogers, Dana, Dutton, Willis, and others. Suess proposed the restriction of meaning because the monoclinical fold (a result of tension and radially acting force) was thought to be genetically and thoroughly contrasted with anticlines and synclines (results of compression and tangential

* J. D. Dana: *Textbook of Geology*, 1864, p. 42; *New Textbook of Geology*, 4th ed., 1883(?) ; *Manual of Geology*, 4th ed., 1895, p. 102.

force). Yet some structures, correctly described as synclines and anticlines because of their arrangement of dips, are probably due to forces acting radially and not directly parallel to the earth's surface. In any case, it seems better, as far as possible, to keep all these elementary terms purely empirical in meaning and free from theoretical connotation.

SUMMARY

Thus simplicity, the demands of a logical system, the active need of field geologists, and a certain degree of priority suggest that Dutton and Geikie should be followed in their definition of "monocline." It may be noted that de Margerie and Heim recommended the noun "monoclinal" as a synonym for the continental word "Flexur," or "flexure."¹ Hence there appears to be no reason why French and German writers should not aid in internationalizing the nomenclature here advocated.

If "monocline" and "monoclinal" be fixed in the Dutton-Geikie sense, the broad, useful concept denoted by the "monoclinal" of W. R. and H. D. Rogers needs a new name. As already observed, that term is readily found by following literally the Rogers recipe for its manufacture. They intended to name a body of strata showing throughout dip in the *same* direction and for that "homocline" (or "homoclinal"; adjective, "homoclinal") is the appropriate word. Its use would have the advantage of rescuing "monocline" from its threatened fate of meaning two different things, and therefore meaning neither without cumbersome explanations. In stratigraphic, physiographic, economic, and even philosophical geology "homocline" should be distinctly useful.

Presented in abstract extemporaneously.

DISCUSSION

MR. G. W. STOSE: I welcome the suggestion of the term homocline. The need of a term for this structure has long been felt in the United States Geological Survey, where confusion has been avoided in editing by using *monoclinal* for the homocline as distinct from *monocline* or monoclinial fold. The term homocline will not only be useful in referring to rocks having the same dip, but whose structure is not known, as mentioned by Professor Daly, but also for the beds between an anticline and syncline, to which neither the term limb of the anticline nor limb of the syncline is not appropriate.

PROF. WILLIAM H. HOBBS: I wish to add my indorsement of Professor Daly's position and express the belief that the use of homoclinal in the sense of uniformity of dip will prevent much misunderstanding. A region where such a term might have been used to advantage is that of the Taconic shales in eastern New York State. Almost by accident I came on much compressed anticlinal cores in a number of localities which showed the wide zone of westerly dipping shales to be isoclinal, though before regarded as a single limb of a fold.

¹ E. de Margerie and A. Heim: Les dislocations de l'écorce terrestre. Zürich, 1888, p. 26.

OCCURRENCE OF INTRAFORMATIONAL CONGLOMERATE AND BRECCIA

BY F. V. EMERSON¹*(Abstract)*

Areas of round clay pebbles and angular masses of stratified clay appear in the sands and sandy clays of Eocene age near Shreveport, Louisiana. The rounded pebbles are of sandy clay, soft when wet, but fairly firm when dry, and of diameters up to 3 inches. Many pebbles are incrustated with iron oxides, forming concretionary masses. The clay fragments are mainly tabular, but occasionally there are large masses of stratified and often well-jointed clay which stand at all angles. The matrix is a rudely stratified sandy clay, in places strongly cross-bedded.

The rounded pebbles indicate weak wave action in waters of moderate depth and the clay breccias with their disordered arrangements show sliding and slumping.

Presented in abstract extemporaneously.

Brief remarks were made by Messrs. A. C. Lane and A. P. Coleman.

KEWEENAW FAULT

BY ALFRED C. LANE

CONTENTS

	Page
Summary.....	93
History.....	93
The primitive Keweenaw fault.....	95
Primitive erosion.....	96
Relation of Keweenawan formation to Cambrian and Precambrian.....	97
The Paleozoic invasions.....	97
The Appalachian revolution.....	99
Later history.....	100

SUMMARY

The Keweenaw fault began as a block-fault in an interior basin like those of the Great Basin region and was also a line of volcanic activity (figure 2).

The Upper Keweenawan were deposited as playa beds in the trough *d* and shade into the "Eastern" Upper Cambrian sandstone, and that perhaps into the "Trenton" limestone. Thrice later—in Richmond, Niagara, and Mid-Devonian times—the sea covered the region (figure 4), but retired at the time of the Appalachian Revolution, when an overthrust fault took place (figures 6 and 7). Erosion reduced the land to a peneplain in Keweenawan time and exposes sometimes the block-fault, sometimes the thrust-fault.

HISTORY

The great Keweenaw fault, which bounds the south side of one of the greatest copper-producing regions, has been an important one not only eco-

¹ Introduced by A. P. Brigham.

nomically and structurally, but in the history of geology. It is not necessary to review the literature recently summarized in United States Geological Society Monograph 52 and my own report on the Keweenaw series. I may, however, recall the names of Jackson and Foster and Whitney, Agassiz, Pumpelly and Wadsworth, Hubbard, Van Hise and Leith among those who have written on it. Very recently an important paper on Limestone Mountain, by Case and W. I. Robinson,¹ has appeared which has a bearing on the subject, as does also the work of Thwaites in Wisconsin.² I have also occasion to refer to the recent work of Allen and Barrett.³ I have left until the last *honoris causa* the Monographic Bulletin 23, by Irving and Chamberlin, issued just 30 years ago.⁴ It would be strange if 30 years of such active progress in research as the past 30 had not produced more facts and fuller

FIGURE 1.—“Ideal Sketch of the primitive Keweenaw Fault”

As represented in figure 24, page 113, of Bulletin 23, U. S. Geological Survey

light on the subject in a region which has been in the meantime the seat of much diamond-drilling and study.

In this paper I briefly compare three diagrams from Bulletin 23 with three others more nearly in accord with the facts as I now know them. Figures 1, 3, and 5 are figures 24, 25, and 26 of Bulletin 23, not pretending to be sections, but diagrams to explain the view of the authors. Figures 2, 4, and 6 are equally diagrams, not sections, for it would be impossible to get Limestone Mountain into a section with the Keweenaw fault on a sufficiently large scale. The bases are the corresponding figures of the set 1, 3, and 5 modified to

¹ Journal of Geology, vol. xxiii, 1915, p. 256.

² Sandstones of the Wisconsin coast of Lake Superior, Belt No. 25 of Wis. Geol. and N. H. Survey.

³ Contributions to Precambrian geology. Mich. Geol. and Biol. Survey, Pub. 18, 1915.

⁴ U. S. Geol. Survey Bull. 23. “Observations on the junction between the Eastern sandstone and the Keweenaw series on Keweenaw Point, Lake Superior.

represent more nearly the geological structure. Figures 7 and 8 with 6 correspond to 5.

THE PRIMITIVE KEWEENAW FAULT

We⁵ agree that the faulting began in Keweenaw time. In figure 2, by changes from figure 1, I have tried to show (1) that *b* of the Keweenawan lies on the underlying Huronian unconformably, the beds in contact on the two sides

FIGURE 2.—Ideal Sketch of the same Fault as Figure 1, supposing it to be a Block-fault

differing many thousands of feet in horizon. This is well marked around Sunday Lake, as noticed by H. L. Smyth, Rose, and others. In fact, according to Allen's recent discoveries, the whole Upper Huronian was eroded away along much of the Gogebic range. (2) That this fault-line acted as a channel or conduit for volcanic activity, so that intrusive pipes (*h*) and sills (*g*) are

FIGURE 3.—"Ideal Sketch of the Keweenaw Fault, after the Deposition of the Eastern Sandstone and before the secondary faulting"

As represented in figure 25, loc. cit.

found near it—for example, Mount Bohemia gabbro, the Torch Lake quartz porphyry, Indiana Mine porphyries and gabbro, Berglund intrusives, etcetera—and the more viscous felsic lavas are also found near it. One period of eruption of rhyolites culminates before the deposition of conglomerate 8, (*c*) of

⁵ "We" refers to the authors of Bulletin 23 and myself.

this figure. (3) Therefore I have changed the hade of the fault from that of an overthrust to the steep hade of a block-fault. There is some direct evidence of this. It is certainly not as flat as a later overthrust fault (Old Colony and Torch Lake and Oneco sections). A nearly vertical hade would be also a more natural channel for volcanic activities. Many overthrust faults are devoid of them.

PRIMITIVE EROSION

We agree, as shown in figures 3 and 4, that this primitive fault suffered erosion, and that the Eastern or Jacobsville sandstone may have lapped on both sides of it. But the study of land formations, which has much advanced in the last 20 years, would lead one to infer that the valley at *d* was being filled as it was formed, and that there is no such sharp gap in dip or strike or time of deposition between the Upper Keweenaw and the Cambrian sandstone, as is fairly inferable from figure 3. In fact, my own figure 4 does not do justice to my conception of the continuity of sedimentation that takes

FIGURE 4.—A Stage corresponding to Figure 3

Showing the section after the deposition of the four Paleozoic overlaps

place in a block-fault valley or graben. The Keweenaw is for me largely the land facies (the Old Red Sandstone facies) of the Middle and Early Cambrian (Waucobic). We know now definitely through Thwaites and the Wisconsin Survey that the Upper Keweenaw is indistinguishable in lithology, dip, or strike from the Cambrian sandstones, for the "Western" Apostle Islands sandstones that have been hitherto called Cambrian by every one he classes as Upper Keweenaw, and the line between them and the Mississippi Valley fossiliferous Upper Cambrian sandstone is drawn in a region where there are no outcrops. However, Limestone Mountain at *j* has fortunately been left us (figure 8) to show that the "Eastern" sandstone is really directly beneath the Trenton. The Jacobsville or Eastern sandstone shows both in drill cores (as at the Oneco mine) and exposures fragments of the Keweenaw amygdaloids, that indicate clearly that at the deposition there was yet a little of the uplifted fault-block exposed to attack. The association of zeolites with copper shows the cooperation of thermal waters, and I therefore believe that much, if not all, of the copper deposition took place during this period of early erosion.

RELATION OF KEWEENAW FORMATION TO CAMBRIAN AND PRECAMBRIAN

Before the deposition of the Keweenaw the Huronian was uplifted and suffered deep erosion, so deep that, according to Allen,⁶ over a large part of the Gogebic Range the Neo-Huronian, which he calls the Copps formation, is removed. Consequently apparently Van Hise mistook the Animikie, or Middle Huronian, for the Upper Huronian. However great the pre-Keweenaw erosion may be, however, all agree that it existed. It seems to have gone so far and so nearly was a peneplain produced, in the later stages of which conglomerates must be finer and often absent, that coarse basement conglomerates of the Keweenaw are not often found. This is important, as indicating the importance of the gap separating the Keweenaw from the Huronian. The black shales of the Neo-Huronian also might indicate the fine-grained deposits of advanced baseleveling. The Keweenaw formation seems, however, to have been formed when the Huronian was raised above water in a desert something like our Great Basin. The dolomitic marls, the micaceous shales, the arkose sandstones, the slight chemical assorting or leaching of the sediments, as well as the abundance of salt, as I believe, largely connate waters, all favor this conclusion to which, I think, Leith came before myself. I have been especially interested lately in studying angular clay pebbles that I find in the Keweenaw drill cores, and take to be broken up, dried crusts on the top of desert clays. This desert basin formation, with accompanying lava flows, like those of the Snake River Valley of Oregon, lasted until late into the Cambrian; for the Upper Cambrian, Jacobsville, or Eastern sandstone, still is arkose, with as much as three-eighths of the sand grains feldspar, still micaceous, and still has saline, though not *so* saline waters. They become distinctly fresher in the upper part. These Upper Cambrian sandstones also contain pebbles of the Keweenaw, showing that the escarpment of the Keweenaw fault was neither peneplained nor covered at the time of the Upper Cambrian. So I think it is pardonable if I am more confirmed in the belief that the Keweenaw is more largely Cambrian than Precambrian—more parallel to the Triassic, with which it was once confused—than to the Permian. I emphasize the fact that the Keweenaw occurred after rather than before the great Precambrian peneplanation, because Lawrence Martin has acutely suggested this as a test whether the Keweenaw is really Cambrian. The line of contact of the base of the Keweenaw on the various Huronian formations is, as Allen's map shows, fairly smooth, except where disturbed by faulting that also affects the Keweenaw.

THE PALEOZOIC INVASIONS

By the time the sea invaded the valley *d*, however, the fault-scarp was pretty well worn down, as figure 3 indicates, and though I have indicated the Eastern sandstone in figure 4 as not entirely covering the range, it may finally have done so, though even near Limestone Mountain (*j*) there are some small pebbles which may be Keweenaw. Such pebbles certainly come near the Keweenaw fault in strata which one would think are not much below. I

⁶ Loc. cit., chapter III, pp. 58-59.

FIGURE 5.—*“Ideal Sketch of the Keweenaw Fault, after the secondary faulting”*
Loc. cit., figure 26, page 114

FIGURE 6.—*“Ideal Sketch of the Keweenaw Fault just before the secondary Thrust faulting”*

FIGURE 7.—*“Ideal Sketch of the Keweenaw Fault just after the secondary Thrust faulting”*

wanted, however, to indicate the still greater expansion and greater distance from shoreline of the succeeding limestones.

During the Paleozoic we have no evidences of great disturbances; but four times, as Case and Robinson show, the sea advanced over the region, quite possibly when at periods of world-wide quiescence and peneplanation the oceans were so filled as to raise the sealevels at about the same time when, as four chief salients in Schuchert's curve indicate, there was wide-spread submergence of the North American continent. No unconformity and but little disconformity has yet been recognized in these strata in Michigan. They are indicated in figure 4 as practically conformable except near where Limestone Mountain was to be. I have indicated the Richmond as a mere filled channel in the Trenton, because it seems to have been less widely distributed in Michigan.

There may well have been, we should expect that there were, later Carboniferous invasions, all traces of which have been removed. There is little Devonian left.

We have then:

1. "Trenton"—Ordovician—Decorah—Black River..... T.
2. Cincinnati R.
3. "Niagara"—Upper Silurian—Lockport N.
4. Mid-Devonian—probably Hamilton—compare the overlap at Milwaukee, Wisconsin D.

The first and third I had recognized years ago; the second and fourth Robinson has just recognized.

THE APPALACHIAN REVOLUTION

Some time after the Mid-Devonian the faulting, folding, and disturbances of the "Eastern" sandstone took place, which on the whole are very slight, but which left us Limestone Mountain (*j*) as a fortunate remnant. I take it that the further disturbances along the Keweenaw fault, illustrated by figure 5, from Irving and Chamberlin, and 6 and 7, were at about the same time. They were certainly later than the deposition of the Eastern sandstone, and had there been much disturbance earlier than the Devonian I think it would have been registered at Limestone Mountain. Moreover, the earlier Paleozoic is generally quiescent in the Great Lakes region; but the Appalachian Revolution may well have had some effect even here. I have indicated this effect in two stages. In figure 6, I suppose, a continuation of the block-faulting. This is indicated by flatter dips, and even sometimes southerly dips, near the fault along the range (New Baltic and South Lake mines). Though such southerly dips may be produced by simple thrust, it is easier at least to draw them by supposing a monoclinic fold or faulting, such as shown, to precede the thrust. Moreover, the consolidation of the beds *b-d* with a greater thickness down to the Archean than on the side of *h* would naturally lead to some folding or slump faulting on that side. Either with or without this preliminary bending it is easy to see that the same tangential compression that has bowed Lake Superior and produced the thrust-faulting and folding described by Thwaites, striking on an uneven wall, with soft sandstone on one side and

hard trap on the other, would shear off the top and cause it to override the sandstone, as shown in figure 7. That the copper deposition was largely prior to this folding is shown by numerous facts. A copper-bearing lode is cut off by the underthrust Eastern sandstone. The Algomah mine, the Lake, and South Lake may be conceived of as working on one bedded lode folded so as to be exposed three times.

LATER HISTORY

But all this time erosion was going on, and theoretically near this fault there must have been land deposits of Permo-Triassic age, some remnant of which may some day be found like the fragments of Cretaceous found in the Mesabi Range.

During Cretaceous time the peneplain must have been somewhat higher than the present surface. And sometime in the Tertiary, the down-cutting

FIGURE 8.—Ideal Sketch of a Cross-section through the southern Part of the Keweenaw (Copper) Range (ck), Limestone Mountain (j), Silver Mountain (i) to the Huron Mountains.

streams reaching harder rock through the mantles of paleozoics, became more and more adjusted to the structure, though here and there transverse streams persisted in valleys now occupied by Portage Lake, the Flint Steel, Fire Steel, and Ontonagon. This Tertiary uplift presumably culminated in the Quaternary when this region was overridden by the Keewatin ice, and later was about the limit of the westward spread of the Labrador ice. At first the bed-rock conditions are somewhat as in figure 8. Near the fault-line we find both thrust and vertical faults as well as slump faults, the strata disturbed and broken. The thrust-faults have a flatter hade than the strata and the thrust-faults a steeper. Both tend to cut out certain parts of the section.

The present surface level relative to the fault systems is sometimes higher, sometimes lower, so that in places we find the Keweenaw formation overlying the Eastern sandstone, as at *k* of figure 8, and again upturned against it, as would be the case did the present surface pass just above *m* of figure 8.

Presented in abstract extemporeaneously.

SOME STRUCTURAL FEATURES IN THE GREEN MOUNTAIN BELT OF ROCKS

BY C. E. GORDON

(Abstract)

The Taconic, or, more broadly speaking, the Green Mountain belt of rocks, is notably, if not preeminently, a region of fracture. The great displacements are genetically of the reversed type, although in later adjustments there may have been some normal faulting along the planes of earlier reversed faults. These great fracture lines follow the general trend of the Green Mountain axis, but often cross it. An east-west trend of the foliations of the ancient gneisses, which has been observed at certain places, may be genetically associated with these transverse breaks if, as seems likely, the lines of weakness in these basal gneisses coincide with their structural features.

Erosion has been extensive enough to expose the faulted relations of the gneisses and the younger rocks. A reasonable restoration of the Precambrian floor on the basis of this faulting gives a different impression of the topography of the land over which the Cambrian sea transgressed from what the present topographic relations, viewed without recognition of the extensive fracture of the basal Precambrian floor, might convey.

Presented by title in the absence of the author.

FAULTING IN NORTH-CENTRAL KENTUCKY

BY ARTHUR M. MILLER

Faults which are all of the normal type, with prominent drag zones on their downthrow sides, are by no means uncommon in the Blue-grass region of the State, though, on account of the length of time which has elapsed since they were formed and the great thickness of residual soil covering, they are not always easy to detect in the uplands.

They vary in length from a few hundred yards to nearly 50 miles, and in throw from 5 or 10 to 350 feet.

They nearly always occur in pairs—a primary and a secondary fault—the latter evidently consequent on the former.

The adjustment of tension strains accompanying a bowing up of the Cincinnati Geanticline might well result in slippings along tension fissures. Such would constitute the “primary” faults. The dropping down with tilting in of the strata toward these faults would tend to develop parallel tension cracks at certain distances away from the primary cracks. A further slipping and tilting of the strata with the continued bowing up of the anticline would widen and deepen the developing secondary fissures until they, too, would become planes of slipping, constituting “secondary” faults, and the wedges of strata between the primary and secondary would become “fault-blocks.”

This type of faulting is well illustrated in the West Hickman Creek fault strip, which stretches from near Paris, in Bourbon County, to near Union Mills, in Jessamine County, a distance of 25 miles.

The maximum throw near the middle of the strip of the west (primary) fault is about 150 feet. Nearly its whole extent is in the uplands of the Central Blue-grass region, where it displaces Eden shale on Lexington lime-

stone. It has formed with its secondary member a nearly continuous narrow strip of relatively poor Eden shale land, sharply contrasted with the excellent Lexington limestone typical blue-grass land on either side of it. This linear tract of beech-tree, sobby, "soapstone" land—from an eighth to a mile wide, penetrating the heart of the Blue-grass—the farmers of the region, in accordance with a tradition handed down from the early settlers, have been accustomed to account for on the theory that it is an "old buffalo trail." It passes 1 mile east of Lexington, and at this point was chosen as a site for the main city reservoir, the clay bottom it has furnished being excellent for holding water.

A very similar poor-land strip stretches from near Great Crossings, in Scott County, past Stamping Ground, in the same county, to near Peaks Mill in Franklin, a distance of about 12 miles.

The two bounding faults of this strip are the Kissinger (primary) and the Switzer (secondary). The amount of the throws and the strata displaced are the same as in the West Hickman, except that toward the western end Maysville (Upper Cincinnati) has been preserved by being brought down on a level with the Eden.

The most prominent of these faults is that of the Kentucky River. It stretches from near Levee, in Montgomery, to near Burdetts Knob, in Garrard County, a distance of 50 miles. The maximum throw in the middle of the stretch is about 350 feet. It here displaces in the uplands Garrard sandstone and Lower Maysville on Lexington limestone, and at the bottom of the gorge of the Kentucky River Upper Lexington on Lower Highbridge (Camp Nelson). The general trend is north 70° east.

Another is the Glencairn fault, crossing the Lexington and Eastern Railroad at Glencairn Station. It stretches from near Campton, in Wolfe County, to near Irvine, in Estill County, a distance of about 25 miles. The maximum throw is about 150 feet. Through most of its extent it displaces at the surface Lower Pennsylvanian on Upper Mississippian.

The Kentucky River fault is so called because from Boonesboro, in Madison County, to Camp Nelson, in Jessamine County, a distance by the windings of the river of 41 miles, the general course of the river and the fault coincides. In this stretch the river in its meanderings crosses the fault nine times, flowing in a narrow picturesque gorge of hard Highbridge limestone, whenever it is on the northwest (relative upthrow) side of the fault, and in a wider valley, presenting less pronounced scenic features, when it is on the southeast (downthrow) side, where it is bedded in the softer limestones and shales of the Lexington or Eden.

While the general southwest diversion of the river in this course is in evident relation to the fault, the minor bends, or the meanders proper, bear no relation to it.

At the eastern ends the Kentucky River and Glencairn faults die out in monoclines which involve at the surface Mississippian and Pennsylvanian, and underneath strata at least as low as the Silurian, for it has resulted in the accumulation at these two ends of the Ragland and Campton oil pools respectively.

The Ragland was discovered accidentally as the result of wild-cat drilling, the Campton in response to advice given by the writer, based on what he

suspected was the relation in the former instance between the oil and the monoclinical structure and the monoclinical structure and the fault. Later a shallow oil pool was developed at the southwestern end of the Glencairn fault, near Irvine.

The oil horizon in the first two instances appears to be a magnesian limestone of approximately Clinton (Brassfield) age. In the Irvine wells the oil appears to come from the Columbus (Devonian) limestone.

In the Lexington limestone of the Blue-grass region are barite veins which have been rather extensively worked in recent years. They are in evident relation to the faults, being fillings of solution cavities, which are widened cracks usually extending outward from the faults at approximately right angles. The walls show no signs of any considerable vertical displacements, but there is some evidence of horizontal movements.

All this fissuring and faulting appears to be of the same age, which is post-Pennsylvanian, as Pennsylvanian strata are traversed and displaced by some of them.

It is also evidently pre-Tertiary. The latter conclusion is reached from a study of the relations of the Kentucky River to its fault.

The Kentucky appears to be a superimposed river which formerly crossed a baseleveled Cincinnati arch in a northwesterly direction from the present site of Boonesboro to where Elkhorn Creek now empties into it. This would carry it by the present site of Lexington.

As the river during a subsequent period of rejuvenation cut its way deeper into the strata, it came more and more under the influence of the fault, the effect of which was to cause it to veer in its course to the southwest.

The excellent illustrations of intrenched meanders in this portion of its course, as in other portions, would seem to indicate that the diversion was before any of the present meanders were acquired.

These meanders are supposed to date from a period of Tertiary baseleveling, hence the Kentucky River fault, and apparently all the other faulting as well, is pre-Tertiary.

We have independent evidence that the Cincinnati Geanticline was first formed in late Silurian or early Devonian time (researches of Foerste). It appears to have received a second bowing up after the aggradation period of the Pennsylvanian, which followed the complete peneplanation just before the latter period was ushered in.

Provisionally we may assign this faulting to the post-Pennsylvanian revolution.

Though now only the eastern end of the faults traverse and displace Pennsylvanian strata, there is strong evidence from outliers scattered out over the Blue-grass region that the strata from the Ordovician up to and inclusive of the Pennsylvanian—except very uppermost Silurian and lowermost Devonian—once went over the Cincinnati arch.

The recentness of the strata preserved in these faulted outliers is generally proportional to the amount of the throw in the faults and the distance they are situated from the axis of the arch.

One very instructive example is Burdett's Knob, near the southern end of the Kentucky River fault, almost on the summit of the central anticlinal dome.

Here is preserved a little patch of Lower Mississippian.

One can hardly avoid the conclusion that the Mississippian went over the arch; and if the Mississippian was thoroughly baseleveled before the deposition of the Pennsylvanian, it is hard to escape from the inference that the latter must also have stretched across.

Presented in abstract extemporaneously.

DISCUSSION

Prof. FRANK R. VAN HORN: In Professor Miller's discussion of the faults, with mention of the barite veins, he omitted to state that the veins contain small amounts of galena and sphalerite. It is probable that the Lexington veins are similar in origin of those of southern Illinois and western Kentucky, in Caldwell and Crittenden counties, where the same mineral association was found, except that fluorite is the gangue mineral instead of barite. In this latter region the veins were worked at first for the lead, but lately for the fluorite, which was used as a flux.

MECHANICS OF INTRUSION OF THE BLACK HILLS PRECAMBRIAN GRANITE

BY SIDNEY PAIGE

(Abstract)

A consideration of the Black Hills granite intrusion may throw light on the mechanics of intrusion in general.

Field relations indicate that the Black Hills granite came into its present position in the main by physical distension of the invaded rock body, under great load. The schists were deformed by the advance of the magma; were forced into closely appressed recumbent folds. The schistosity produced by this folding lent itself to further injection by the granite through great numbers of parallel dikes and by intricate lit-par-lit intrusion. The harder rocks—the quartzites—were distended and broken apart by the upward movement and the magma flowed in between the blocks.

It is believed the magma was intruded at a relatively low temperature; that it contained considerable water, and acted as a relatively mobile mass.

The relation of dikes to schist layers, the fact that lit-par-lit injection tends to neutralize chemical and physical differences between magma and invaded rock, the probable low temperature of the granite, and the fact that positive evidence of much assimilation is lacking—all support the belief that this process was but a corollary of physical displacement, and not a primary process by which the forward movement of the magma was accomplished. The possibility that large blocks of roof have been engulfed in the magma, and that this process was an important one, must remain an open question.

Presented in abstract extemporaneously.

DISCUSSION

Prof. JOSEPH BARRELL: Mr. Paige has shown in his discussion of the field relations of the Black Hills granite that they indicate intrusion in the main, by physical distension of the invaded rock body, under great load.

This may seem at first sight to be an argument against the validity of other modes of intrusion in other regions, more especially against the validity of the stoping hypothesis. Although, as Daly has said, the stoping hypothesis was postulated independently by a number of geologists, it has been so largely developed by his work that he is deservedly regarded as its chief founder and expositor. As one of those, however, who have contributed papers in which it is held that the rise of certain batholiths was by stoping, it seems appropriate for me to speak of the relations of the two modes of intrusion. The invasion of the Black Hills granite by crustal distension seems a logical conclusion, but at the same time does not controvert the evidence that certain other bodies have risen to their final boundaries by a process of stoping. In this statement I am not taking exception to Mr. Paige's conclusions. In fact, we have discussed this topic and I think we agree; but where a more or less novel view, such as the stoping hypothesis was ten years ago, is advocated as an explanation of certain examples, the idea is apt to arise among readers that the expounders wish to give it a universality of application and eliminate competing hypotheses.

* What, then, is the adjustment of factors which in one case may lead to lateral displacement as a chief mode of invasion, in another case to overhead stoping, permitting a passive rise? The rise of magmas from abyssal depths would appear to result from the unbalancing of a hydrostatic equilibrium. The liquid column is lighter than the surrounding rocks. As the liquid arises above that datum plane where the liquid and solid are under the same pressure, the internal pressure at any level is diminished by the weight of the column of liquid below; the pressure in the surrounding rocks is diminished by the greater weight of the rock column between the level and the datum plane below. A bursting and intruding pressure is consequently developed. In the zone of flowage the thick cover would permit this internal pressure to act laterally, pushing the walls aside. Injection into the roof in steep foliation planes also implies a lateral distension. More or less doming of the cover is of course also to be postulated.

When a large magmatic body has advanced, however, comparatively near to the surface, a lateral distension of the walls or cover becomes subordinate, because the line of least resistance is now for the magma to dome the cover upward, to produce intrusion fractures, and to intrude in distinct sheets and dikes, rather than to produce a *lit-par-lit* injection. The mechanical conditions making stoping a dominant process are then found especially in the zone of fracture; those making for injection, mashing, and crystallization with lateral distension of the wall rocks prevail in the zone of flow. The Black Hills granite belongs to the Precambrian. In these ancient batholiths crustal distension by the invading magmas appears to have been a dominant process, though stoping even here may have participated. The far younger batholiths of the Cordillera were intruded to high levels and with more or less absence of compressive forces. In these stoping appears to be a dominant process in the higher stages of their invasion. If we could restore the great depths eroded from the Archeozoic and look at the former higher levels, or if we could look deeper into the crust to observe the relations of post-Jurassic batholiths yet concealed, the distinction in dominant modes of intrusion evident between the older and younger periods of igneous activity might be found to have largely disappeared.

Mr. CHARLES E. DECKER: Having heard that an unconformity had been found in the Precambrian in the Black Hills, I should like to ask Mr. Paige about the conglomerate he mentions in his paper. Is it a basal conglomerate, and is there evidence of an unconformity beneath it?

Brief remarks were made by Prof. R. A. Daly.

Mr. PAIGE replied: I am gratified that both Professor Daly and Professor Barrell agree that the Black Hills granite may have come to place in such a manner as I have indicated. The idea that the variation in the mechanics of igneous intrusion is a function of depth and temperature has long appealed to me as reasonable: that magmas and the rocks that they invade must act differently at varying depths seems a logical deduction from what we know of the earth's temperature and pressure gradient and from what we have observed in the field. The idea, carried to its ultimate analysis, demands that there should be a region at very great depths where, due to intricate injection and prolonged impregnation by the magma, sediments will break down and lose their identity as such, taking on all the characteristics of igneous rocks.

PRECAMBRIAN STRUCTURE OF THE BLACK HILLS, SOUTH DAKOTA

BY SIDNEY PAIGE

(Abstract)

A study of the Precambrian rocks of the Black Hills reveals a great series of slates and schists, for the most part monotonously alike, striking in a north-west direction, and having steep dips, generally, except in the extreme south-west, to the east. At the south intrusions of granite break through the strata, and around the principal mass of granite about Harney Peak a schistosity is developed parallel with the granite contact.

A closer study shows that the persistent eastward dips represent both schistosity and bedding, the two for the most part parallel, and that the series is compressed into a number of great folds which comprise innumerable minor isoclinal folds.

And, finally, a sufficient number of individual beds can be traced to locate the position and nature of the greater axes of folding and to detect the presence of two important faults.

The nature of the granite intrusion is such as to lead to the belief that the mass exerted an important influence on the folding, and the position of one of the faults was the determining factor in the localization of the great Homestake ore body.

Presented in abstract extemporaneously.

TITLES AND ABSTRACTS OF PAPERS PRESENTED BEFORE THE AFTERNOON
SESSION AND DISCUSSIONS THEREON

The Society convened at 2 o'clock p. m., with President Coleman in the chair.

The Society proceeded immediately to the consideration of scientific papers.

RECTILINEAR FEATURES IN THE EASTERN CATSKILLS

BY GEORGE H. CHADWICK

(Abstract)

It seems to have escaped notice that there is in the eastern Catskill Mountains a series of right-line valleys gridironed across the ranges in a direction 15° east of north similar in character to those features in the Adirondacks that have been interpreted as due to block-faulting; but no proof is at hand that such faults exist in the horizontal red beds of the Catskills. The valleys indicated are, however, parallel with the mural front of these mountains facing the Hudson Valley and with the strike ridges of folded Helderberg limestones in that valley, a few miles distant. The field evidence suggests that the cause of these features is a close spacing of joints along zones at intervals.

Presented in abstract extemporaneously.

DISCUSSION

MR. JOHN L. RICH: Mr. Chadwick has presented a very interesting paper, and with his primary thesis I am heartily in accord. With regard to the straight mural front of the Catskills, which is interpreted as probably a product of glacial erosion, I wish to suggest the alternative hypothesis that, since it parallels the folding in the Hudson Valley to the east, it may be rather a structural feature representing the eastern limit of the flat-lying beds.

The speaker suggests that the roughly accordant, flat-topped summits of the higher Catskills may be remnants of a peneplain, possibly Jurassic. Without specifically opposing this proposition, I wish to give expression to my increasing distrust of the verity in regions of horizontal rocks, particularly where some beds are distinctly more resistant than others, of peneplains for which the only evidence are small flat areas on the summits and accordant altitudes.

DR. W. J. MILLER: In the eastern Adirondacks there are numerous normal faults, proved to be such by most of the familiar criteria for the recognition of faults. In many other cases, however, there are prominent zones of excessive jointing, often showing more or less evidence of faulting by the slickenside, fault breccia, etcetera. In other cases such evidence of actual displacement is lacking. Many times it is difficult to decide whether a given zone is due simply to excessive jointing or whether the jointing has accompanied actual faulting in the zone.

Doctor CHADWICK replied to Mr. Rich as follows: The peneplains bevel across the rock layers, and consequently are independent of the harder layers except for short intervals.

Doctor CHADWICK replied to Doctor Miller as follows: There is no question of the existence of faults in the eastern Adirondacks, but it is unsafe to assume that all similar rectilinear valleys are of fault origin, and particularly in the northwestern Adirondack region, where the work of Doctor Martin fails to find faults, although the physiographic features are similar.

PHYSIOGRAPHIC EVIDENCE OF RECENT SUBSIDENCE ON THE COAST OF MAINE

BY CHARLES A. DAVIS

(Abstract)

For several years the author has been collecting evidence which seemed to him to prove a recent subsidence of the North Atlantic coast, still in progress.

Some of these evidences have already been presented before this Society. During the summer of 1915 some attention has been given to the physiographic evidence of subsidence presented by certain beaches in southwestern Maine, where an ancient beach was found almost buried in a salt marsh, which clearly must have been formed subsequent to the beach. The unaltered crest of this old beach is several feet lower than that of the modern beach of exactly the same type. Other beaches with multiple ridges in the same region give noteworthy confirmation of the evidence furnished by the one mentioned.

Some other physiographic evidence of subsidence now in progress was presented, and this was shown to be fully in harmony with other types of evidence already presented in former papers.

Presented in abstract extemporaneously.

PHYSIOGRAPHIC NOTES ON THE WHITE MOUNTAINS

BY DOUGLAS W. JOHNSON

(Abstract)

The paper presents certain observations made during a brief visit to the White Mountains in the summer of 1914.

The origin of the felsenmeer or block moraine, the date of cirque formation, and the position of the New England peneplain with reference to the mountains are briefly treated.

Presented by title in the absence of the author.

*POSITION OF THE NEW ENGLAND UPLAND IN THE WHITE MOUNTAINS*BY ARMIN K. LOBECK¹*(Abstract)*

An attempt to trace the New England peneplain from the base of Mount Monadnock to the White Mountains has led to the unqualified conclusion that in the latter region the peneplain lies near the base of the mountains, and that the mountains represent monadnocks rising far above the badly dissected upland surface. In addition to the field evidence, a study of the topographic sheets east of the mountains and Hitchcock's Atlas of New Hampshire, by means of projected sections, decisively supports the conclusion.

Presented in abstract extemporaneously.

¹ Introduced by Douglas W. Johnson.

*STUDY OF RIPPLE-MARKS*BY WALTER A. BUCHER¹*(Abstract)*

The mechanics of the problem; mathematical laws; results of new experiments; significance in stratigraphy; field observations in Kentucky and Ohio.

Presented by title in the absence of the author.

DEAD LAKE OF THE CHIPOLA RIVER, FLORIDA

BY E. H. SELLARDS

(Abstract)

The Chipola River in Florida affords a good illustration of a tributary stream ponded by the deposition of sediment in the valley of the main stream. This river, which originates entirely within the coastal plain, flows for a considerable part of its course across limestones and is fed very largely by clear water limestone springs. It is therefore a clear-water stream, carrying only a limited amount of sediment. The Apalachicola River, of which the Chipola is a tributary, heads on the contrary in the mountains of northern Georgia and receives a heavy load of sediment, which is deposited, as the river becomes overloaded, near the Gulf. The Chipola enters the Apalachicola about 25 miles, by land, from the Gulf of Mexico. In this lower part of its course the Apalachicola is rapidly aggrading its valley, while the Chipola, owing to the limited amount of sediment which it carries, is building its valley much more slowly. This condition results in flooding the valley of the Chipola River. The lake thus formed, known as the Dead Lake of the Chipola River, derives an added interest from the fact that it has come into existence so recently that the cypress timber of the former river swamp, now mostly dead, is still standing, although the water in the lake has reached a depth of from 10 to 20 feet, while the lake itself is 10 or 12 miles long and from 1 to 2 miles wide. The channel of the river may still be followed in its winding course through the lake.

Presented in abstract extemporaneously.

*TECTONIC LINES IN THE HAWAIIAN ISLANDS*BY SIDNEY POWERS²*(Abstract)*

The alignment of the Hawaiian Islands shows that they have been formed at the points of intersection of two sets of tectonic lines, one set running parallel to the group of islands from northwest to southeast, the other set being cross-fractures from northeast to southwest. On the island of Hawaii the rift lines of Mauna Loa and of Kilauea are quite pronounced structural

¹ Introduced by Nevin M. Fenneman.

² Introduced by John E. Wolff.

features following the second set. In the case of Mauna Loa, one rift starts near Hilo, on the east coast, and the other from Ka Lae, the southernmost point of the island, and at their intersection the sink of Mokuaweoweo has been formed. South of Mauna Loa are the almost parallel rifts of Kilauea: one from the east of Pahala, on the south shore; one from near the Olaa Mill, on the northeast, and a third from the line of pit-craters on the east—all intersecting in the sink of Kilauea.

Extensive faulting, not clearly related to the above tectonic lines, has accompanied the formation of the islands. On Niihau the eastern portion of the original volcano has dropped out of sight, leaving a line of sheer cliffs. Kauai was originally a volcanic doublet, but the volcanic pile on the northwest has subsided, as is seen in the truncated flows of the Napali cliffs. On Oahu a portion of the Kaala Range may have dropped down in blocks, forming the "pali" (cliff), and cinder cones with some lava flows may have risen between the crags, giving rise to part of the present topography. On the north side of Molokai there is a line of cliffs, rising to a height of 1,000 to 3,500 feet, which bounds the southern third of the original volcano. After the collapse of the northern two-thirds, lava flows built up a small peninsula at the foot of the cliffs, and on this peninsula the leper colony is now situated.

On Hawaii the breakdown of a portion of the Kahala dome is marked by cliffs on the northeast shore. The west side of Mauna Loa is scarred by a crescentic-shaped fault from Kealakekua southward, along which there has been subsidence on the seaward side. Later flows of lava have run over the cliffs and obscured the fault-line. The crater of Mohokea forms a crescentic depression on the south side of Mauna Loa, but the age of this crater, with respect to Mauna Loa, is uncertain. The south side of the dome of Kilauea, near the sea, from Kalapana nearly to Punaluu, has dropped down in a series of steps over which the later lava flows have poured from far back in geologic time until the last flow of 1868. The last movement along this shore was a subsidence of from 4 to 7 feet in 1868.

Presented by title in the absence of the author.

*BANDED GLACIAL SLATES OF PERMOCARBONIFEROUS AGE, SHOWING
POSSIBLE SEASONAL VARIATIONS IN DEPOSITION*

BY ROBERT W. SAYLES¹

(Abstract)

Above the Squantum tillite, which I described last year, there is a series of banded slates about 800 feet in thickness. The manner of transition from the tillite to the slate, with no unconformity, makes it certain that the slate was the result of deposition in waters from the melting of the glacier which formed the tillite. The first of the transition beds are conglomerates alternating with sandstones. Then come sandstones alternating with slates. Here and there it is plain that the ice readvanced over these beds, plowing them up and often dragging upward into the mass thus formed irregular lumps of the clay. Two beds of tillite in the slate itself show the close association of the slate with

¹ Introduced by J. B. Woodworth.

the main tillite. One of these beds is about 50 feet from the main tillite formation and the other about 150 feet. Prof. John E. Wolff has made a microscopical examination of the tillite and slate and finds that they have the same mineralogical composition. As one goes upward in the banded slate it is seen that the layers of sandstone become thinner and thinner, then disappear entirely, and finally the alternating layers are shown only in slate of dark and light bands, the dark bands containing more heavy basic material than the light. The cause of this disappearance of the sandstone layers can be explained by the gradual withdrawal of the glacier and the consequent slackening of the currents, which would be strong enough to carry sand only in the neighborhood of the ice. The lower pebbly members of the transition beds show irregularities of deposition, especially in lenticular forms, due to the inconstant conditions of streams coming from a glacier. A regularity of alternation in deposition, however, becomes evident after the first 50 feet or so of the transition beds have been passed, where the layers indicate deeper and quieter water, and thus more uniformity in deposition. The thin individual layers of sandstone and slate now show through hundreds of feet such regularity in thickness and interval that a regularly recurring cause must be sought.

Professors B. K. Emerson and Gerard De Geer have described this same regularity of layers of sand and clay in the glacial clays of the Connecticut Valley and Sweden respectively and have ascribed the banding to yearly deposition. They believe that the layers of coarser material indicate summer, when the streams from the melting ice were rapid, and the layers of fine material indicate winter, when the streams had less velocity. Neither Emerson nor De Geer could suggest any other period to which such regular alternations could be related. Other American geologists have published this same idea which De Geer conceived in 1878. Among them may be mentioned Dr. Charles P. Berkey, Prof. A. P. Coleman, and Mr. Frank B. Taylor. This explanation may not be generally accepted, but no other cause, so far suggested, accounts for all the facts of the case. When the seasonal units have been accurately determined, they will furnish a basis for estimating the length of time required for the deposition of the formation.

The facts observed in the slate at Squantum resemble so closely those described by De Geer and Emerson that it would seem as if there must be a very strong probability of these being similar cases, in spite of the millions of years which separate the two Glacial periods.

Read in full from manuscript.

DISCUSSION

Prof. W. W. Atwood: This paper appeals to me as particularly interesting. I have seen similarly banded clays associated with the Wisconsin drift and during the past season I found just such banded clays associated with the Eocene till of southwestern Colorado. In examining in the field the slates here described by Mr. Sayles I was convinced that the regularity of the repetition of the bands of sands and clays meant a regularly returning cause, and seasonal changes in stream action seem adequate. This paper suggests that the idea of seasonal changes in deposition so well put forth by De Geer and others may help us in working out a calendar of ancient geologic time and

lead to a fuller appreciation of the significance of varying climatic conditions on sedimentation.

Professor HOBBS: I have been greatly interested in Mr. Sayles' able discussion of this subject and the conclusions which he has reached concerning the annual deposits indicated by the tillite. It was my good fortune to examine with Baron De Geer a number of sections in the thin-banded clay in the delta deposits along the former ice-front of Scandinavia. Checked as these individual bands were by moraines of annual deposition, and worked out as they have been with such infinite care, one could hardly fail to accept the conclusion to which they pointed. The conclusions of Mr. Sayles point the way for many studies in the future which it is certain will be crowned with results. I was interested to note that in some of his sections there was indication of what represents, perhaps, a climatic cycle, something entirely to be expected, though it did not seem to be indicated by the banded clays near Stockholm.

Prof. JOSEPH BARRELL: Doctor Sayles has given a very convincing presentation, showing that the banding in the argillites associated with the Squantum tillite corresponds to the similar banding in Pleistocene glacial clays and is to be interpreted as showing seasonal variations in the Permocarboneous glacial climates. This is both interesting and important, since the biologic evidence of most periods previous to the later Mesozoic shows warmth without marked zonal climates extending into high latitudes.

In the combinations of winter and summer conditions which go to make up the geologic climates, there are four combinations possible, so far as warmth and cold are concerned: *First*, winter may be submerged and summer is dominant throughout the year, with or without a season of drought. This is the present climate of the torrid zone and in a number of past times appears to have been the dominant type even into high latitudes. It means the existence of some condition which carried solar heat received in equatorial latitudes into high latitudes with minimum loss by radiation. *Second*, winter and summer may alternate, but with mild winters, the present climates of the warm temperate latitudes. This was perhaps the commonest type of climate in high latitudes during the Jurassic and Cretaceous periods. *Third*, winter and summer may alternate, but with rigorous winters, leading, with sufficient snow-fall, to glacial conditions. This is the climate typical of high temperate latitudes today, but in the Pleistocene it extended into middle latitudes. It is favored by direct and local absorption of solar heat and its re-radiation, with minor control by a planetary distribution of equatorial warmth. It seems, as Doctor Sayles has here shown, to have been the type also of the Permocarboneous glaciation in eastern Massachusetts. *Fourth*, the climate may be continuous winter, as seen *within* the present ice-sheets in polar latitudes—summer completely submerged, winter continually dominant.

Now, regarding the obliquity of the ecliptic and position of the pole as unchanged through geologic time, the middle latitudes should show the greatest contrast between the total amounts of winter and summer insolation. Doctor Sayles' paper suggests, in accordance with this expectation, that glaciation in all times in the temperate zone has been connected, as at present, with a marked seasonal climate, exhibiting the third of the several combinations listed—a climate distinguished from normal geological climates by the emergence of winter.

Let us go much farther back in time. A similar stratigraphic banding exists in argillites associated with the Huronian tillites at Cobalt, Ontario. I was fortunate in being able to examine this section under the guidance of Messrs. Miller, Knight, and Coleman during the International Geological Excursion in 1913, although they are not responsible for the following statement of observations.

The banding has, I believe, been noted for a long time by the Ontario geologists and may have been discussed in their reports. At the south end of Cobalt Lake occurs a thick bed of argillite delicately banded, indicating rhythmic deposition. The bands are grouped in series which show larger rhythms. If the bands are annual, the rhythmically recurring groups show climatic fluctuations covering periods of years. On the southeast side of Cobalt Lake the argillites are succeeded by a ripple-marked quartzite. The thick till sheet rests on this without obliterating the ripples or crumpling the argillite beds at a greater depth. The boulder conglomerate is faintly bedded with grits and some beds of banded argillite show occasional subangular boulders dropped into them. Many of the boulders are markedly soled and indicate undoubted glacial wear. The whole suggests the presence of quiet water, which at first was far enough from the glacier front to receive only silt and clay. Later it was mantled apparently by an iceberg deposit rather than a true ground till. This implies, however, true till in some region not far distant. At a later stage there may have been an overriding of ice in this same region. In other places at Cobalt what appears to be undoubted tillite containing striated boulders rests directly on the floor of Keewatin rocks, although the final proof, consisting of striations on that floor, has not yet been found. The association of these banded argillites with iceberg deposits and ground moraine indicates, therefore, the existence of summer melting and winter freezing during the Huronian glaciation.

Sederholm also has found in Finland as far back as during Bottnian time a regular alternation of clayey and sandy material which he interprets as an annual stratification, explainable only by assuming a regular change of seasons.

Glacial conditions then seem to be brought about by a marked lowering of the mean annual temperature of the whole globe, accompanied by an accentuation of seasons in middle latitudes. This suggests a less effective spreading out of equatorial warmth at such times and a more direct and rapid absorption and also radiation of solar energy in the temperate zones. Glacial periods are marked by strong seasonal climates, the emergence of winter. This is a fact which, according to its distinctness, must enter into the adjustment of factors bearing on the causes of Glacial periods. The explanation of non-glacial climates must, on the other hand, account for the submergence of winter in middle and high latitudes with the earth's axis permanently situated as it is at present. The explanation of glacial periods and of warm-pole periods are equally important and complementary problems. The presence of marked seasonal climates in temperate latitudes during glacial periods is an important fact which must enter into that explanation.

MR. FRANK B. TAYLOR: Laminated pebbleless clays formed in the preglacial lakes which followed the retreating Wisconsin ice-sheet occur in many places in the Great Lakes region. A deposit of this kind occurs at Bracebridge, in

the highlands east of Georgian Bay. It is nearly 100 feet thick, and if the successive pairs of laminae be regarded as marking annual periods of deposition, this deposit, as I now recall it, would indicate a time of about 2,500 years for its formation. It has seemed to me that laminated clays of this type are particularly characteristic of deposition in association with the retreating ice-sheet, and it is a striking and impressive fact to find precisely similar clays greatly indurated and so closely associated with the ancient tillites.

Mr. JOHN L. RICH: In a bottom of an old pond I once found a deposit of beautifully developed banded clays, in which the bands were marked by fragments of leaves. In this case the banding clearly represented seasonal accumulation, the leaf-bearing partings corresponding to autumn and winter, when the fallen leaves were being carried and deposited by the streams.

Prof. R. B. WOODWORTH said that Mr. Sayles' correlation of the banded shales with the tillite beds at Squantum added another group of water-laid deposits which have been identified among the ancient glacial formations wherein we should expect to find the equivalents of our Pleistocene and modern sand plains, kames and espars, and brick clays; already possible examples of one or another of these have been reported in Brazil and elsewhere.

GEOLOGY OF THE LAKE IDITAROD REGION, ALASKA

BY PHILIP S. SMITH

(Abstract)

The paper describes the areal geology of the Lake Clark-Iditarod region, Alaska. This region is located in southwestern Alaska, extending from the Pacific Mountains to the central part of the Yukon Plateau province. The rocks are dominantly sedimentary strata of Mesozoic age, but some Paleozoic limestones are also exposed. Igneous rocks, both of intrusive and of effusive origin, occur at a number of places, and certain of them seem to have been closely associated with the deposits of commercial value, such as gold and quicksilver. Unconsolidated deposits are widespread and throughout much of the region mantle over and hide underlying bedrock. These deposits are mainly of glacial and glaciofluvial origin, though lacustrine, fluvial, and volcanic ash deposits are also described.

Presented by title in the absence of the author.

CHARACTERISTICS OF THE SOIL AND ITS RELATION TO GEOLOGY

BY C. F. MARBUT

(Abstract)

The soil is made up mainly of rock-waste. Its characteristics are of two kinds: (1) those derived from the rock from which it came, or those that are inherited, and (2) those impressed on it by the forces that have been acting on it since its life history began, or those that are acquired. The latter characteristics are relatively few in number and vary with many factors. The conditions determining their character and strength of expression were discussed.

The relation of geology to soil was brought out by (1) a comparison of the chemical composition of a number of soils developed from the same rocks under varying conditions; (2) a comparison of the physical characteristics of a number of soils developed from the same rocks under varying conditions; (3) a comparison of the physical characteristics of a number of soils developed from different rocks under similar conditions; (4) a comparison of the physical characteristics of a number of soils developed from the same rocks under varying topographic conditions.

Presented in abstract extemporaneously.

GEOLOGIC SIGNIFICANCE AND GENETIC CLASSIFICATION OF ARKOSE DEPOSITS

BY DONALD C. BARTON¹

(Abstract)

Different opinions are held as to the conditions under which arkose has been formed. A study of arkose deposits shows that they are of several distinct types, of which the more important are (a) deposits of great thickness and with fresh feldspars, (b) deposits of moderate size associated with coal and argillaceous beds, and (c) deposits of moderate size, more or less argillaceous and associated with red beds. A survey of the conditions under which arkose could be expected to form shows that the granular disintegration of granite is widespread and is a possibility under practically all climates; that in rigorous climates erosion and deposition of the granitic sand can take place contemporaneously with the disintegration, but that in regions of moist climate the mantle of vegetation must be critically weakened by some change of conditions, such as a marine transgression or increasing aridity, before the products of disintegration and decomposition beneath can be eroded and then accumulated. It therefore seems possible to formulate a genetic classification of arkose deposits.

Presented by title in the absence of the author.

SOME FEATURES OF THE KANSAN DRIFT IN SOUTHERN IOWA

BY GEORGE F. KAY

(Abstract)

In county reports issued by the Iowa Geological Survey and in other publications many of the features of the Kansan drift of southern Iowa have been described, including the original Kansan drift plain, the present topography of the Kansan drift, the tabular divides, the characteristics of the weathered and unweathered zones of the Kansan drift, the gumbo, which is closely related to the Kansan drift, and the fine, loesslike clay overlying the Kansan drift surface, and which has been interpreted by several investigators to be material of eolian origin deposited after a mature topography had been developed on the Kansan drift. The origin of the gumbo has been interpreted

¹ Introduced by R. A. Daly.

differently by different authors, the most recently published view being that of Tilton, who considers the material to have been formed, in the main, during the retreating stages of the Kansan ice. To this gumbo and other materials which he considers to be related in age to the gumbo he has given the name Dallas deposits.

Detailed field studies which are still in progress in southern Iowa seem to warrant the author in making a preliminary statement involving some interpretations which differ from those previously advanced.

(1) The surface of the Kansan drift, after the Kansan ice withdrew, was, according to present evidence, a ground moraine plain, which, from the main divide between the Mississippi and Missouri rivers, sloped gently to the southeast and south toward the Mississippi and to the southwestward toward the Missouri. This drift plain was so situated topographically that weathering agents were very effective, but erosion was slight. As a result of the weathering during an exceedingly long time a grayish, tenacious, thoroughly leached, and non-laminated joint clay, which has been named gumbo, was developed to a maximum thickness of more than 20 feet. This gumbo contains only a few pebbles, which are almost wholly siliceous, and grades downward into yellowish and chocolate-colored Kansan drift from 3 to 7 feet in thickness, in many places with numerous pebbles, few, if any, of which are calcareous. This oxidized but non-calcareous drift, in turn, merges into unleached drift, oxidized yellowish for several feet, below which is the normal unleached and unoxidized dark-grayish to bluish-black Kansan drift. The gumbo is believed, therefore, to be essentially the result of the thorough chemical weathering of the Kansan drift; but, subordinately, other factors, such as the wind, freezing and thawing, burrowing of animals, slope wash, etcetera, have undoubtedly contributed to its formation. The Kansan drift which has been changed to gumbo may have differed somewhat from the normal Kansan drift that lies below the gumbo.

(2) After the gumbo plain had been developed by weathering processes on the Kansan drift plain, diastrophic movements seem to have occurred, the plain having been elevated to such an extent that erosion became effective and valleys began to be cut into the gumbo plain. Erosion of the gumbo plain progressed to such an extent that some valleys were cut to a depth of more than 150 feet before grade was reached and a mature topography was developed. Only remnants of the original gumbo plain remain, the most conspicuous of these being flat, poorly drained areas, known as tabular divides. Where creep and slumping have occurred the gumbo, in places, may be found on slopes at an elevation several feet below the level of the gumbo plain. The tabular divides are more prevalent east of a line drawn north and south through south-central Iowa than west of such a line. In the southwestern part of the State the Kansan gumbo, which is *in situ*, is found only where the divides, which are no longer distinctly tabular, retain the level of the former gumbo plain.

(3) While there is, in places, loess of eolian origin on the Kansan drift of southern Iowa, much of the material which has been described as loess is thought to be not of eolian origin, but to be related more or less closely to the gumbo. The upper few feet of the Kansan gumbo, which is now limited to the tabular divides and divides closely related to tabular divides, is a fine-grained, loesslike, joint clay, in which, if diligent search is made, it is possible

to find a few very small siliceous pebbles similar to those in the normal gumbo, and it is thought that this loesslike clay is the result of changes that have been going on at and near the surface of the gumbo during the great length of time since the normal gumbo was formed. The loesslike clay which is now found as a mantle on the Kansan drift on the slopes and divides that have been brought by erosion considerably below the level of the original gumbo plain is believed to be the product not primarily of wind action, although wind may have been a factor, but chiefly the product of the weathering and concentration of the gumbo and to some extent of the underlying Kansan drift, where erosion has not kept pace with the weathering.

(4) The evidence indicates that the time taken to develop the present topography from the gumbo plain stage, although it represents a great length of time, is short when compared with the time taken to develop the gumbo plain from the Kansan drift. It is thought that the formation of the main part of the gumbo and the development of the present mature topography of the Kansan drift were effected between the close of the Kansan epoch and the advance of the Illinoian ice into Iowa; in other words, during the Yarmouth inter-Glacial epoch. All the evidence indicates that the Yarmouth epoch was an exceedingly long interval of time.

(5) Detailed chemical analyses of gumbo, loesslike clay, etcetera, are now being made in the chemical laboratory of the University of Iowa by Dr. J. N. Pearce. The results of these analyses will go far to strengthen or weaken the interpretations given above from the field evidence.

Presented in abstract extemporaneously.

DISCUSSION

Mr. W. C. ALDEN: I would like to make a few remarks on the *Iowan drift* in connection with Doctor Kay's paper. I have seen a number of exposures of this clay on top of the Kansan drift, which, for want of a better name, we have called "gumbo." It seems to me a very important deposit. When first seen I thought it must be distinct from the underlying till, inasmuch as the till below it is oxidized, though the gumbo itself is mostly gray and unoxidized, excepting at the top. On further examination, however, I do not feel sure of its being distinct, and considerable evidence has been found indicating that it really may be the weathered upper part of the Kansan till which has been thoroughly leached of its soluble material and deoxidized. It is nowhere laminated like a water-laid silt, where I have seen it. It generally contains pebbles scattered sparsely throughout. The pebbles are mostly small and of chert and quartz, with some quartzites and crystallines, the latter badly decomposed. Rarely included boulders are found disintegrating and with feldspars decomposing to kaolin. Thicknesses of 15 to 20 feet of this clay are not uncommon. There is no definite line at the base of the gumbo; it grades into the till below. I am not yet fully convinced that it is the residuum of weathering of the upper part of the till, but there is much to suggest that this is the case. Probably Doctor Kay's investigations will settle the matter. If it is such, I think it indicates a very long time of exposure of the Kansan drift, prior to the Illinoian stage of glaciation, since the gumbo is also found under Illinoian till in Henry County, Illinois. It seems to make the Yarmouth in-

terval much longer than some of us have been inclined to think heretofore, especially since the development of the gumbo appears to have taken place largely, if not wholly, before the dissection of the Kansan drift plain. It seems to be confined to the flat uplands and does not mantle the eroded slopes. In any case it clearly marks a stratigraphic horizon, the top of the Kansan till.

The point to which I wish particularly to call attention is the relation of this super-Kansan gumbo to the *Iowan drift*. For a number of years past there has been considerable question in the minds of some of the geologists as to whether there really is a post-Kansan drift in northeastern Iowa. This was because of difference of interpretation of some of the phenomena. It was finally decided that the Federal Survey should cooperate with the Iowa Geological Survey in a critical review of the evidence in the field. I was assigned to this work with Morris M. Leighton of the Iowa Survey as assistant, and we spent the field seasons of 1914 and 1915 in the investigation. It is a pleasure to be able to announce that we have reached the conclusion that there really is good evidence of the presence of a post-Kansan drift sheet in that area; so that we are in the main in agreement with the late Doctor Calvin in regard to the Iowan drift.

We found the area much less markedly eroded than the Kansan drift area. The topography is of a peculiar type. We have called it a mantled, mature-erosion topography—that is, there are there the large patterns of maturely developed dendritic drainage systems—but most of the minor details have apparently been smoothed out. It is what one would expect if an ice-sheet were to spread over the maturely dissected topography of the Kansan drift area, wearing away the spurs, smoothing out the irregularities, partly filling the valleys, and mantling the whole with a rather thin sheet of till. This is indeed just what seems to have occurred. The slopes are very largely long, low, smooth, and, excepting close to the main streams, almost wholly uncut by lateral drainage lines. The minor valleys are shallow, open swales with smoothly rounded bottoms, where the streams flow in shallow trenches. There is a notable absence of the V-shaped cross-profiles seen in the Kansan area.

The uppermost drift throughout the Iowan drift area is a highly calcareous till. It is more modified by weathering than the Wisconsin, more so indeed than one would infer from some of the published descriptions, yet not nearly so much changed as the Kansan drift; neither is it so much modified as the Illinoian drift. The famous big granite boulders, although not entirely confined to the Iowan drift area, are notably abundant there and seem to be particularly characteristic of this drift deposit.

Now, the particular interest in regard to the super-Kansan gumbo is that it affords a new line of evidence which tends to corroborate the other data. Many new exposures have recently been afforded by grading for country roads, electric interurban railroad lines, and the Chicago, Milwaukee and Saint Paul Railroad; also we made about 250 test borings with an auger 8 feet in length. We have found numerous exposures of the super-Kansan gumbo, or a deposit exactly like it, in the higher parts of the Iowan area, those parts particularly where remnants of the original Kansan drift plain are likely to have been preserved. In several places there is a black carbonaceous bed at the top, apparently an old soil, and overlying this a thin deposit of later till—the Iowan till. Where this till is less than 4 or 5 feet thick, its limestone pebbles and

finer calcareous material have generally been entirely removed by leaching; where the drift is thicker than 4 or 5 feet, it is highly calcareous below the leached zone. At one place the gumbo bed outcropped in a slope about 50 feet below the top of the ridge.

It thus appears that there is clearly a post-Kansan till in northeastern Iowa. It is older than the Wisconsin and seems to be distinctly younger than the Illinoian. The loess, which is largely absent or very thin in the Iowan area, mantles the weathered and eroded surface of the Illinoian drift as it does that of the Kansan. A similar gumbo bed was found in western Iowa between two drift sheets, apparently the Kansan and Nebraskan, but this need not be discussed at this time.

Mr. FRANK LEVERETT stated that in case the gumbo is a derivative from boulder clay microscopical examination should bring to light insoluble minerals, aside from siliceous rocks, that were combined to make up the rocks which are common in the boulder clay, but have not been found preserved in coarse fragments in the gumbo. The interpretation by Professor Kay seems at best but tentative and will need to be tested by such lines of evidence as are now under investigation, so that a year hence we may know better the value of the interpretation. So far as Mr. Leverett's personal observations have gone, there seems lacking the transition zone between boulder clay and gumbo which one would expect to find if the gumbo is a derivative from boulder clay. Usually less than a foot space will take one from typical boulder clay into typical gumbo, and one's first impression is that the gumbo has been deposited on the boulder clay as a distinct bed.

Mr. CHARLES E. DECKER: I have seen examples of the gumbo which Professor Kay describes at the Survey Office in Washington. These samples are very much like a thoroughly leached clay that occurs in connection with the oldest drift in northwestern Pennsylvania at Franklin and Oil City. However, there are larger fragments of rock in the clay at those localities than in the specimens of gumbo.

TRIASSIC ROCKS OF ALASKA

BY GEORGE C. MARTIN

(Abstract)

Triassic rocks, including over 5,000 feet of limestones and shales with highly characteristic marine Upper Triassic (Karnic and Noric) faunas, are now known to be widely distributed in Alaska. The Middle and Lower Triassic are represented by a single known occurrence of marine Middle Triassic shales and possibly also by volcanic and metamorphic beds. The more important Alaskan Triassic sections were described and correlated. Correlations were also suggested with the Triassic occurrences in British Columbia, as well as with the better known Triassic sections in California and other parts of the world. An outline was given of the events of Triassic time in the northwestern part of America.

Presented in abstract extemporaneously.

*LITHOGENESIS AND STRATIGRAPHY OF THE RED BEDS OF SOUTHEASTERN WYOMING*BY S. H. KNIGHT¹*(Abstract)*

The principal object of this paper is to announce the results thus far obtained from a detailed study of the red beds of southern Wyoming. This study is an attempt to determine, first, the origin and source of the sediments of which the red beds are composed; second, the physical conditions which were active during their deposition; and, lastly, their relations to contemporaneous and adjacent formations. The sections thus far studied in detail are exposed along the eastern flank of the Medicine Bow Mountains and along the western flank of the Laramie Mountains. This region affords a good opportunity for a study of this nature, inasmuch as there is a striking change in the lithological character of the lower half of red-bed series, as first brought out by my father, W. C. Knight, and later emphasized by Darton.

The change in lithological character throws considerable light on the direction from which the material was derived and the distance which it has been transported. Along the eastern flank of the Medicine Bow Range the lower half of the red-bed series (Casper formation) consists of about equal amounts of conglomeratic arkose and sandstones, with a very minor development of shale, while the same formation 20 miles to the east, along the western flank of the Laramie Range, exhibits in one section, which may be taken as typical, the following: arkose, 2 per cent; sandstone, 61 per cent; shale, 22 per cent, and limestone, 15 per cent. The total thicknesses of the formations in the two localities are 710 feet and 687 feet respectively. Owing to the size, angularity, and freshness of the feldspars in the conglomeratic arkoses, one may say that the material has not been derived from any great distance; this, together with the fact that these arkoses are practically wanting in the Laramie Range section, leads one to believe that the material had its origin to the west, probably from the region now occupied, in part at least, by the Medicine Bow Mountains. The evidence for the upper half of the series is not so clear, but it is hoped that some petrographical and mechanical analyses now being conducted will throw some new light on the problem.

We may now turn to the physical conditions which influenced the deposition of the red beds in this region. The evidence points to a continental origin for a large part of the series. Torrential, fluvatile, and eolian deposits have been recognized. The evidence for a continental interpretation may be summed up as follows: (1) Lack of continuity in sections measured at short intervals; (2) channeling to a depth of 10 or more feet; in one section seven erosion intervals were noted within 450 feet of the base of the section; (3) the ever-prevailing cross-bedding of both the torrential and eolian type; (4) the character of the material; (5) color; (6) lack of marine fossils. In this connection I wish to call attention to a striking conglomerate 150 feet from the base of the series, in an exposure on Sand Creek, Albany County, Wyoming. This conglomerate has a thickness of 34 feet and consists for the most part of water-worn pebbles and boulders of an igneous rock nature, the maximum

¹ Introduced by A. W. Grabau.

diameter of the larger boulders measuring 10 inches. Embedded near the base of this conglomerate were found a series of sandstone blocks, the largest of which measured 3 feet 8 inches by 7 feet 6 inches in its longest and shortest diameters. Thirty of these blocks were counted in 200 feet of outcrop. The material of which these blocks are composed consists of a fine-grained, evenly bedded, friable, red sandstone. These blocks are very angular in outline and the bedding planes of the various blocks are tipped at all angles. They present a strong contrast to the well rounded, smooth, hard pebbles and boulders with which they are embedded. It is evident from the size and angular outline, together with the friable nature of the sandstone, that these blocks have not been carried laterally any great distance, unless they can be conceived to have been floated or rafted by ice action. A careful search was made for some evidence of ice activities in the conglomerate, but none was found. In seeking for an interpretation which will account for the presence of these blocks in such a position, we have only to turn to some interesting facts recorded higher in the same section. It might be well to note that the lower 500 feet of the section, in which these blocks occur, is composed of heavy beds of a coarse sandy to conglomeratic arkose, interbedded with a number of thinner bands of fine-grained red sandstone, the materials of which are indistinguishable from those comprising the above-mentioned blocks. At a height of 330 feet from the base of the section one of these sandstone members was found to have a uniform thickness of 7 feet. Some 18 feet above this member (348 feet from the base) a second sandstone member, having a maximum thickness of 10 feet, was found to be channeled to a depth of 5 feet. At a distance of 370 feet from the base two large sandstone blocks, having regular bases on the same horizontal plane, were found to be completely inclosed by coarse arkose. The largest of these two blocks measured 26 feet long and 4 feet thick. Lastly, at a height of 416 feet from the base four sandstone blocks, all with regular bases in the same horizontal plane, were found completely inclosed by coarse arkose. The largest of this set measured 2 feet 6 inches in thickness, while the distance between the extreme ends of the two outer blocks was 50 feet. In all cases these sandstone members were found to have irregular upper surfaces, while their lower were approximately straight, and in each set on the same horizontal plane. From the foregoing it is apparent that we are dealing in each case with what was originally a continuous sandstone member, which, with the exception of the case noted at a height of 330 feet, has been channeled previous to the deposition of the overlying arkose members. In the cases noted from 370 and 416 feet from the base this channeling has cut completely through the beds, thus permitting the overlying and underlying arkose members to come together. We may carry this development a step further and assume that during a period of exceptional torrential activity, when coarse conglomerates were being deposited, such sandstone remnants of preexisting beds may have been undermined by the lateral cutting of a stream and tumbled into the river bed, where they would be deposited along with the pebbles and boulders being swept down by the torrent. This occurrence will be described more fully in a subsequent paper.

As to the stratigraphical relationships and age of this red-bed series, two points brought out by this research might be mentioned. First, that all of the limestone members (some ten having been recognized), which tend to replace

the red beds to the east, have yielded marine fossils. These fossils are now being studied in the Paleontological Laboratories at Columbia University. This fauna substantiates the previously recognized Pennsylvanian age of these limestones. In the Red Mountain region, near the Wyoming-Colorado boundary line, a thin red conglomerate was noted 177 feet below the base of the overlying Morrison shale. This conglomerate contains numerous bone fragments, mostly of a water-worn nature; but after a two-day search the writer was rewarded by the finding of a complete limb bone, the nature of which is yet undetermined. It is anticipated that this horizon, which was recognized in two sections 30 miles apart, will yield better material when thoroughly examined.

Presented by title in the absence of the author.

*EXPERIMENT IN THE GRAPHIC PRESENTATION OF THE ECONOMIC GEOLOGY
OF BEDDED DEPOSITS*

BY GEORGE H. ASHLEY

(Abstract)

An attempt to present, by means of charts and maps, all of the known data concerning the resources of an area, so that, without the text, which in the main is confined to an explanation of the charts and maps, the reader may determine for himself the position on or under the surface, probable thickness, etcetera, of any and all beds of coal, clay, limestone, iron, etcetera, at any point in the area, so far as the data on hand warrant any conclusions.

Presented in abstract extemporaneously.

BRECCIATION EFFECTS IN THE SAINT LOUIS LIMESTONE

BY FRANCIS M. VAN TUYL¹

(Abstract)

The Saint Louis limestone, as developed in southeastern Iowa and adjacent parts of Illinois and Missouri, is frequently so badly brecciated and disturbed that all semblance of the original structure is lost. The character and cause of this disturbance has until the present never been fully investigated, although it has been commented on by nearly all geologists who have examined the formation even casually.

Recent study of the Saint Louis has made necessary considerable revision of the formation as defined in previous reports. It is now known that the arenaceo-magnesian limestone sometimes shown at its base and formerly included with it is a distinct formation of the horizon of the Spergen limestone. This has been shown to be both physically and faunally distinct and to be separated from the true Saint Louis by a disconformity.

Moreover, the Pella member, which was formerly regarded as the topmost subdivision of the Saint Louis, is also formationally distinct. This bears a good Sainte Genevieve fauna and is separated from the true Saint Louis be-

¹ Introduced by T. E. Savage.

neath by a disconformity of considerable magnitude. It possesses a characteristic basal sandstone. As at present restricted, therefore, the Saint Louis represents only about one-half of the strata originally referred to it, or about 50 to 60 feet of limestone.

In southeastern Iowa the formation, as revised, is divisible into two well marked portions, which in their typical development are of about equal thickness. These are everywhere separated from each other by a slight disconformity. The lowermost of these is usually dolomitic, either wholly or in part, and the lower portion of the upper one is sometimes so. To the north-westward the upper division soon disappears, but the lower one continues to the north-central part of the State, overlapping older and older formations, until at Humboldt it is found resting on beds of Kinderhook age.

Where undolomitized, both divisions of the Saint Louis consist typically of fine-grained, compact, brittle, gray limestone, which breaks with a subconchoidal fracture. Beds of granular limestones a few feet in thickness, however, appear at some horizons, and the lower portion of the lower division is often shaly or arenaceous, or both. The dolomitic portions, on the other hand, are buff in color, somewhat coarser in grain, and tough. These occur both as interbedded layers in the limestone and as lateral gradations or facies of limestone beds of considerable thickness. That the formation was deposited in shallow seas is clearly indicated by several features hereinafter noted.

The brecciation effects in the formation may be grouped into three main types. First, the disturbed portion may assume the form of a small mound or reef of limestone blocks, usually in a calcareous or sandy matrix, with undisturbed layers lapping up on its flanks and filling in its depressions. These may appear at any horizon in the formation. Second, it may appear along minor faults in the Lower Saint Louis, or be confined largely to one bed of this member, due to differential movements. The latter is accompanied occasionally by the tonguelike extensions of the material forced down into the beds below, especially when the latter are softer. Third, it may embrace the major part, if not the entire formation, over an area of variable but usually limited extent. In this type the disturbance has been produced by mashing on a large scale, and the brecciation is often associated with folds and overthrust faults of small magnitude. The Pella beds are also involved to a slight extent in this disturbance, but in no instances have the underlying formations been found to exhibit signs of this type of deformation.

Regarding the origin of the brecciation, at least three periods of disturbance are believed to have operated in its production. The mounds or reef-like masses of the first type are believed to have been formed during the first period under conditions of violent wave action, possibly induced by local shallowing of the sea during deposition. The presence of local disconformities in the formation and of cross-bedded lime sands supposedly formed by the grinding up of layers already deposited is in favor of this view. Other features which suggest wave action at the time of deposition are contemporaneous erosion phenomena, wave-marks, and cross-bedding.

The brecciation of the second type was formed during the second period of deformation. The importance of this can not be definitely evaluated, since its effects are often overshadowed by the disturbance of the third type. But that this disturbance is distinct from the third is indicated by the fact that

dolomitization intervened between the two. Thus the reefs produced by the first disturbance and the shattered areas and fracture lines produced by the second are often either wholly undolomitized or are very imperfectly altered, while the undisturbed limestone about them is uniformly dolomitized. This proves that dolomitization succeeded the first and second periods of disturbance. But a later disturbance involves both the poorly dolomitized areas and the uniformly dolomitized ones, and a later series of fractures cuts the earlier ones.

Furthermore, the fact that dolomitization never affects the topmost limestone layers of the Saint Louis nor any layers of the Pella indicates that the alteration took place prior to the close of the Saint Louis. Now, since the brecciation of the second type is known to have taken place still earlier, and it is known to be confined to the Lower Saint Louis, it seems probable that this shattering may be related to the uplift which brought this division to a close.

The third period of disturbance was by far the most important. To this is ascribed the extensive mashing and shearing effects and the overthrust faulting on a small scale so common in the formation. This was influenced to a large extent by the effects of the preceding disturbances, thereby obscuring to a large degree the evidence of these. That the deformation which produced this is post-Pella, but pre-Pennsylvanian in age, is indicated by the fact that blocks of Pella limestone have been found sheared down into lower beds, and thus preserved at a locality where Pennsylvanian sandstone generally rests disconformably on these beds of a lower horizon.

The general parallelism of the strike of the faults and of the tilted layers formed at this time with certain anticlines in the region, notably the Benton-sport anticline which trends approximately north 68° west, suggests a common mode of origin of these two types of deformation.

Presented by title in the absence of the author.

VOTE OF THANKS

The following resolution presented by Professor Fairchild was adopted:

"Resolved, That the sincere and hearty thanks of the Society be extended to the members of the local committee who have made the efficient and satisfactory arrangements for this large and very successful meeting, especially Messrs. T. W. Vaughan and A. C. Spencer, in general charge of the local committees, and the chairmen of the subcommittees, Messrs. L. W. Stephenson, P. S. Smith, F. L. Ransome, Whitman Cross, and E. S. Bastin.

"Resolved, That the Society express its thanks to the George Washington University for the use of this building as a place of meeting."

The Society adjourned at 5.20 o'clock p. m.

REGISTER OF THE WASHINGTON MEETING, 1915

FELLOWS

FRANK D. ADAMS	CHARLES R. DRYER
WILLIAM C. ALDEN	HERMAN L. FAIRCHILD
HENRY M. AMI	CLARENCE N. FENNER
GEORGE H. ASHLEY	CASSIUS A. FISHER
WALLACE W. ATWOOD	AUGUST F. FOERSTE
J. AUSTIN BANCROFT	L. C. GLENN
JOSEPH BARRELL	AMADEUS W. GRABAU
FLORENCE BASCOM	JOHN S. GRASTY
R. S. BASSLER	LOUIS C. GRATON
E. S. BASTIN	BAIRD HALBERSTADT
GEORGE F. BECKER	WILLIAM H. HOBBS
CHARLES P. BERKEY	WILLIAM J. HOLLAND
EDWARD W. BERRY	WILLIAM O. HOTCHKISS
S. W. BEYER	ERNEST HOWE
ELIOT BLACKWELDER	ELLSWORTH HUNTINGTON
JOHN M. BOUTWELL	J. P. IDDINGS
ALFRED HULSE BROOKS	JOHN D. IRVING
H. A. BUEHLER	THOMAS A. JAGGAR
CHARLES BUTTS	GEORGE F. KAY
D. D. CAIRNES	FRANK J. KATZ
H. D. CAMPBELL	ARTHUR KEITH
MARIUS R. CAMPBELL	EDWARD M. KINDLE
CHARLES CAMSELL	EDWIN KIRK
STEPHEN R. CAPPS	CYRIL W. KNIGHT
ERMINE C. CASE	F. H. KNOWLTON
GEORGE H. CHADWICK	ADOLPH KNOPF
WILLIAM B. CLARK	HENRY B. KÜMMEL
JOHN M. CLARKE	ALFRED C. LANE
H. F. CLELAND	WILLIS T. LEE
ARTHUR P. COLEMAN	J. VOLNEY LEWIS
ARTHUR J. COLLIER	FREDERIC B. LOOMIS
HENRY P. CUSHING	SAMUEL W. MCCALLIE
REGINALD A. DALY	H. D. MCCASKEY
N. H. DARTON	GEORGE R. MANSFIELD
WILLIAM M. DAVIS	CURTIS F. MARBUT
JOSEPH S. DILLER	FRANÇOIS E. MATTHES
RICHARD E. DODGE	WALTER C. MENDENHALL
JOHN A. DRESSER	GEORGE P. MERRILL

HERBERT E. MERWIN
 ARTHUR M. MILLER
 WILLETT G. MILLER
 WILLIAM JOHN MILLER
 FRED H. MOFFIT
 ELWOOD S. MOORE
 CLEOPHAS C. O'HARRA
 HENRY F. OSBORN
 SIDNEY PAIGE
 FREDERICK B. PECK
 R. A. F. PENROSE
 GEORGE W. PERKINS
 WILLIAM C. PHALEN
 JOSEPH HYDE PRATT
 LOUIS M. PRINDLE
 FREDERICK S. RANSOME
 CHESTER A. REEDS
 WILLIAM NORTH RICE
 GEORGE B. RICHARDSON
 HEINRICH RIES
 RUDOLPH RUEDEMANN
 ROLLIN D. SALISBURY
 FRANK C. SCHRADER
 CHARLES SCHUCHERT
 ALFRED R. SCHULTZ
 WILLIAM B. SCOTT
 E. H. SELLARDS
 EUGENE W. SHAW
 H. W. SHIMER

WILLIAM J. SINCLAIR
 EUGENE A. SMITH
 GEORGE O. SMITH
 J. W. SPENCER
 J. STANLEY-BROWN
 TIMOTHY W. STANTON
 CLINTON R. STAUFFER
 LLOYD W. STEPHENSON
 RALPH W. STONE
 GEORGE W. STOSE
 C. K. SWARTZ
 STEPHEN TABER
 JAMES E. TALMAGE
 FRANK B. TAYLOR
 JOHAN A. UDDEN
 E. O. ULRICH
 FRANK R. VAN HORN
 GILBERT VAN INGEN
 CHARLES D. WALCOTT
 HENRY S. WASHINGTON
 THOMAS L. WATSON
 CARROLL H. WEGEMANN
 STUART WELLER
 LEWIS G. WESTGATE
 DAVID WHITE
 I. C. WHITE
 GEORGE R. WIELAND
 JOHN E. WOLFF
 J. B. WOODWORTH

FELLOWS-ELECT

THOMAS C. BROWN

EDGAR T. WHERRY

In addition to the foregoing, there were registered at the meeting 17 members of the Paleontological Society and 104 visitors.

OFFICERS, CORRESPONDENTS, AND FELLOWS OF THE
GEOLOGICAL SOCIETY OF AMERICA

OFFICERS FOR 1916

President:

JOHN M. CLARKE, Albany, N. Y.

Vice-Presidents:

J. P. IDDIGS, Brinklow, Md.

HARRY FIELDING REID, Baltimore, Md.

RUDOLPH RUEDEMANN, Albany, N. Y.

Secretary:

EDMUND OTIS HOVEY, American Museum of Natural History,
New York, N. Y.

Treasurer:

WM. BULLOCK CLARK, Johns Hopkins University, Baltimore, Md.

Editor:

J. STANLEY-BROWN, 26 Exchange Place, New York, N. Y.

Librarian:

F. R. VAN HORN, Cleveland, Ohio

Councilors:

(Term expires 1916)

R. A. F. PENROSE, JR., Philadelphia, Pa.

W. W. ATWOOD, Cambridge, Mass.

(Term expires 1917)

CHARLES K. LEITH, Madison, Wis.

THOMAS L. WATSON, Charlottesville, Va.

(Term expires 1918)

FRANK B. TAYLOR, Fort Wayne, Ind.

CHARLES P. BERKEY, New York, N. Y.

*MEMBERSHIP, 1916**CORRESPONDENTS*

- CHARLES BARROIS, Lille, France. December, 1909.
 W. C. BRÖGGER, Christiania, Norway. December, 1909.
 GIOVANNI CAPELLINI, Bologna, Italy. December, 1910.
 BARON GERHARD DE GEER, Stockholm, Sweden. December, 1910.
 SIR ARCHIBALD GEIKIE, Hasslemere, England. December, 1909.
 ALBERT HEIM, Zürich, Switzerland. December, 1909.
 EMANUEL KAYSER, Marburg, Germany. December, 1909.
 W. KILLAN, Grenoble, France. December, 1912.
 J. J. H. TEALL, London, England. December, 1912.
 EMIL TIETZE, Vienna, Austria. December, 1910.

FELLOWS

*Indicates Original Fellow (see article III of Constitution)

- CLEVELAND ABBE, JR., U. S. Weather Bureau, Washington, D. C. August, 1899.
 FRANK DAWSON ADAMS, McGill University, Montreal, Canada. Dec., 1889.
 GEORGE I. ADAMS, 17 San T'iao Hutung, Peking, China. December, 1902.
 JOSÉ GUADALUPE AGUILERA, Instituto Geologico, Mexico, Mexico. Aug., 1896.
 WILLIAM CLINTON ALDEN, U. S. Geological Survey, Washington, D. C. December, 1909.
 TRUMAN H. ALDRICH, Birmingham, Ala. May, 1889.
 JOHN A. ALLAN, University of Alberta, Strathcona, Canada. December, 1914.
 R. C. ALLEN, State Geologist, Lansing, Mich. December, 1911.
 HENRY M. AMI, Strathcona Park, Ottawa, Canada. December, 1889.
 FRANK M. ANDERSON, State Mining Bureau, 2604 Ætna St., Berkeley, Cal. June, 1902.
 ROBERT VAN VLECK ANDERSON, 7 Richmond Terrace, Whitehall, S. W., London, England. December, 1911.
 RALPH ARNOLD, 923 Union Oil Building, Los Angeles, Cal. December, 1904.
 GEORGE HALL ASHLEY, U. S. Geological Survey, Washington, D. C. Aug., 1895.
 WALLACE WALTER ATWOOD, Harvard University, Cambridge, Mass. Dec., 1909.
 RUFUS MATHER BAGG, JR., Lawrence College, Appleton, Wis. December, 1896.
 HARRY FOSTER BAIN, 734 Salisbury House, London, E. C., England. Dec., 1895.
 MANLEY BENSON BAKER, School of Mining, Kingston, Ontario. Dec., 1911.
 S. PRENTISS BALDWIN, 2930 Prospect Ave., Cleveland, Ohio. August, 1897.
 SYDNEY H. BALL, 71 Broadway, New York City. December, 1905.
 JOSEPH A. BANCROFT, McGill University, Montreal, Canada. December, 1914.
 ERWIN HINCKLEY BARBOUR, University of Nebraska, Lincoln, Neb. Dec., 1896.
 JOSEPH BARRELL, Yale University, New Haven, Conn. December, 1902.
 GEORGE H. BARTON, Boston Society of Natural History, Boston, Mass. August, 1890.
 FLORENCE BASCOM, U. S. Geological Survey, Washington, D. C. Aug., 1894.
 RAY SMITH BASSLER, U. S. National Museum, Washington, D. C. Dec., 1906.
 EDSON SUNDERLAND BASTIN, U. S. Geological Survey, Washington, D. C. December, 1909.

- WILLIAM S. BAYLEY, University of Illinois, Urbana, Ill. December, 1888.
- *GEORGE F. BECKER, U. S. Geological Survey, Washington, D. C.
- JOSHUA W. BEEDE, Indiana University, Bloomington, Ind. December, 1902.
- ROBERT BELL, Geological Survey, Department of Mines, Ottawa, Canada. May, 1889.
- CHARLES P. BERKEY, Columbia University, New York, N. Y. August, 1901.
- EDWARD WILBER BERRY, Johns Hopkins University, Baltimore, Md. Dec., 1909.
- SAMUEL WALKER BEYER, Iowa Agricultural College, Ames, Iowa. Dec., 1896.
- ARTHUR B. BIBBINS, Goucher College, Baltimore, Md. December, 1903.
- ELIOT BLACKWELDER, University of Wisconsin, Madison, Wis. Dec., 1908.
- JOHN M. BOUTWELL, 1323 De la Vine St., Santa Barbara, Cal. Dec., 1905.
- JOHN ADAMS BOWNOCKER, Ohio State University, Columbus, Ohio. Dec., 1904.
- *JOHN C. BRANNER, Leland Stanford, Jr., University, Stanford University, Cal.
- EDWIN BAYER BRANSON, University of Missouri, Columbia, Mo. Dec., 1911.
- ALBERT PERRY BRIGHAM, Colgate University, Hamilton, N. Y. December, 1893.
- REGINALD W. BROCK, University of British Columbia, Vancouver, B. C. December, 1904.
- ALFRED HULSE BROOKS, U. S. Geological Survey, Washington, D. C. Aug., 1899.
- AMOS P. BROWN, University of Pennsylvania, Philadelphia, Pa. Dec., 1905.
- BARNUM BROWN, American Museum of Natural History, New York, N. Y. December, 1910.
- CHARLES WILSON BROWN, Brown University, Providence, R. I. Dec., 1908.
- THOMAS CLACHAR BROWN, Bryn Mawr College, Bryn Mawr, Pa. Dec., 1915.
- HENRY ANDREW BUEHLER, Rolla, Mo. December, 1909.
- BERT S. BUTLER, U. S. Geological Survey, Washington, D. C. December, 1912.
- G. MONTAGUE BUTLER, College of Mines, Tucson, Arizona. December, 1911.
- CHARLES BUTTS, U. S. Geological Survey, Washington, D. C. December, 1912.
- DE LORME DONALDSON CAIRNES, Geological Survey Branch, Department of Mines, Ottawa, Canada. December, 1912.
- FRED HARVEY HALL CALHOUN, Clemson College, S. C. December, 1909.
- FRANK C. CALKIN, U. S. Geological Survey, Washington, D. C. Dec., 1914.
- HENRY DONALD CAMPBELL, Washington and Lee University, Lexington, Va. May, 1889.
- MARIUS R. CAMPBELL, U. S. Geological Survey, Washington, D. C. Aug., 1892.
- CHARLES CAMSELL, Geological Survey of Canada, Ottawa, Canada. December, 1914.
- STEPHEN REID CAPPS, JR., U. S. Geological Survey, Washington, D. C. Dec., 1911.
- FRANK CARNEY, Granville, Ohio. December, 1908.
- ERMINIE C. CASE, University of Michigan, Ann Arbor, Mich. December, 1901.
- GEORGE HALCOTT CHADWICK, University of Rochester, Rochester, N. Y. December, 1911.
- ROLLIN T. CHAMBERLIN, University of Chicago, Chicago, Ill. December, 1913.
- *T. C. CHAMBERLIN, University of Chicago, Chicago, Ill.
- CLARENCE RAYMOND CLAGHORN, Tacoma, Wash. August, 1891.
- CHARLES H. CLAPP, University of Arizona, Tucson, Arizona. December, 1914.
- FREDERICK G. CLAPP, 120 Broadway, New York, N. Y. December, 1905.
- *WILLIAM BULLOCK CLARK, Johns Hopkins University, Baltimore, Md.
- JOHN MASON CLARKE, Albany, N. Y. December, 1897.

- HERDMAN F. CLELAND, Williams College, Williamstown, Mass. Dec., 1905.
 J. MORGAN CLEMENTS, 20 Broad St., New York City. December, 1894.
 COLLIER COBB, University of North Carolina, Chapel Hill, N. C. Dec., 1894.
 ARTHUR P. COLEMAN, Toronto University, Toronto, Canada. December, 1896.
 GEORGE L. COLLIE, Beloit College, Beloit, Wis. December, 1897.
 ARTHUR J. COLLIER, U. S. Geological Survey, Washington, D. C. June, 1902.
 CHARLES WILFORD COOK, University of Michigan, Ann Arbor, Mich. December, 1915.
 EUGENE COSTE, 1943 11th St., West, Calgary, Alberta, Canada. Dec., 1906.
 ALJA ROBINSON CROOK, State Museum of Natural History, Springfield, Ill. December, 1898.
 *WILLIAM O. CROSBY, Massachusetts Institute of Technology, Boston, Mass.
 WHITMAN CROSS, U. S. Geological Survey, Washington, D. C. May, 1889.
 GARRY E. CULVER, 1104 Wisconsin St., Stevens Point, Wis. December, 1891.
 EDGAR R. CUMINGS, Indiana University, Bloomington, Ind. August, 1901.
 *HENRY P. CUSHING, Western Reserve University, Cleveland, Ohio..
 REGINALD A. DALY, Harvard University, Cambridge, Mass. December, 1905.
 EDWARD SALISBURY DANA, Yale University, New Haven, Conn. Dec., 1908.
 *NELSON H. DARTON, U. S. Geological Survey, Washington, D. C.
 CHARLES ALBERT DAVIS, U. S. Bureau of Mines, Washington, D. C. Dec., 1910.
 *WILLIAM M. DAVIS, Harvard University, Cambridge, Mass.
 ARTHUR LOUIS DAY, Geophysical Laboratory, Carnegie Institution, Washington, D. C. December, 1909.
 DAVID T. DAY, 1333 F St. N. W., Washington, D. C. August, 1891.
 BASHFORD DEAN, Columbia University, New York, N. Y. December, 1910.
 FRANK WILBRIDGE DE WOLF, Urbana, Ill. December, 1909.
 *JOSEPH S. DILLER, U. S. Geological Survey, Washington, D. C.
 EDWARD V. D'INVILLIERS, 518 Walnut St., Philadelphia, Pa. December, 1888.
 RICHARD E. DODGE, Teachers' College, New York, N. Y. August, 1897.
 NOAH FIELDS DRAKE, Fayetteville, Arkansas. December, 1898.
 JOHN ALEXANDER DRESSER, 326 Notre Dame de Grace Ave., Montreal, Canada. December, 1906.
 CHARLES R. DRYER, Oak Knoll, Fort Wayne, Ind. August, 1897.
 *EDWIN T. DUMBLE, 2003 Main St., Houston, Texas.
 ARTHUR S. EAKLE, University of California, Berkeley, Cal. December, 1899.
 CHARLES R. EASTMAN, American Museum of Natural History, New York, N. Y. December, 1895.
 EDWIN C. ECKEL, Munsey Building, Washington, D. C. December, 1905.
 *BENJAMIN K. EMERSON, Amherst College, Amherst, Mass.
 WILLIAM HARVEY EMMONS, University of Minnesota, Minneapolis, Minn. December, 1912.
 JOHN EYERMAN, Oakhurst, Easton, Pa. August, 1891.
 HAROLD W. FAIRBANKS, Berkeley, Cal. August, 1892.
 *HERMAN L. FAIRCHILD, University of Rochester, Rochester, N. Y.
 OLIVER C. FARRINGTON, Field Museum of Natural History, Chicago, Ill. December, 1895.
 NEVIN M. FENNEMAN, University of Cincinnati, Cincinnati, Ohio. Dec., 1904.
 CLARENCE NORMAN FENNER, Geophysical Laboratory, Washington, D. C. December, 1911.

- CASSIUS ASA FISHER, 711 Ideal Building, Denver, Colo. December, 1908.
AUGUST F. FOERSTE, 128 Rockwood Ave., Dayton, Ohio. December, 1899.
WILLIAM EBENEZER FORD, Sheffield Scientific School, New Haven, Conn. December, 1915.
MYRON LESLIE FULLER, 131 State St., Boston, Mass. December, 1898.
HENRY STEWART GANE, Wonalancet, New Hampshire. December, 1896.
JAMES H. GARDNER, 1014 Daniel Building, Tulsa, Oklahoma. December, 1911.
RUSSELL D. GEORGE, University of Colorado, Boulder, Colo. December, 1906.
*GROVE K. GILBERT, U. S. Geological Survey, Washington, D. C.
ADAM CAPEN GILL, Cornell University, Ithaca, N. Y. December, 1888.
L. C. GLENN, Vanderbilt University, Nashville, Tenn. June, 1900.
JAMES WALTER GOLDTHWAIT, Dartmouth College, Hanover, N. H. Dec., 1909.
CHARLES H. GORDON, University Library, University of Tennessee, Knoxville, Tenn. August, 1893.
CLARENCE E. GORDON, Massachusetts Agricultural College, Amherst, Mass. December, 1913.
CHARLES NEWTON GOULD, 408 Terminal Bldg., Oklahoma City, Okla. December, 1904.
AMADEUS W. GRABAU, Columbia University, New York, N. Y. December, 1898.
WALTER GRANGER, American Museum of Natural History, New York, N. Y. December, 1911.
ULYSSES SHERMAN GRANT, Northwestern University, Evanston, Ill. Dec., 1890.
JOHN SHARSHALL GRASTY, University of Virginia, University, Va. Dec., 1911.
LOUIS C. GRATON, Harvard University, Cambridge, Mass. December, 1913.
HERBERT E. GREGORY, Yale University, New Haven, Conn. August, 1901.
GEORGE P. GRIMSLEY, Geological Survey of West Virginia, Martinsburg, W. Va. August, 1895.
LEON S. GRISWOLD, Plymouth, Mass. August, 1902.
FREDERIC P. GULLIVER, 1112 Morris Bldg., Philadelphia, Pa. August, 1895.
WILLIAM F. E. R. GURLEY, University of Chicago, Chicago, Ill. Dec., 1914.
ARNOLD HAGUE, U. S. Geological Survey, Washington, D. C. May, 1889.
BAIRD HALBERSTADT, Pottsville, Pa. December, 1909.
GILBERT D. HARRIS, Cornell University, Ithaca, N. Y. December, 1903.
JOHN BURCHMORE HARRISON, Georgetown, British Guiana. June, 1902.
CHRIS. A. HARTNAGEL, Education Building, Albany, N. Y. December, 1913.
JOHN B. HASTINGS, 1480 High St., Denver, Colo. May, 1889.
*ERASMUTH HAWORTH, University of Kansas, Lawrence, Kans.
RAY VERNON HENNEN, West Virginia Geological Survey, Morgantown, W. Va. December, 1914.
OSCAR H. HERSHEY, Kellogg, Idaho. December, 1909.
RICHARD R. HICE, Beaver, Pa. December, 1903.
*ROBERT T. HILL, Federal Bldg., Los Angeles, Cal.
RICHARD C. HILLS, Denver, Colo. August, 1894.
HENRY HINDS, U. S. Geological Survey, Washington, D. C. December, 1912.
*CHARLES H. HITCHCOCK, 2376 Oahu Ave., Honolulu, Hawaiian Islands.
WILLIAM HERBERT HOBBS, University of Michigan, Ann Arbor, Mich. August, 1891.
*LEVI HOLBROOK, P. O. Box 536, New York, N. Y.
ROY J. HOLDEN, Virginia Polytechnic Institute, Blacksburg, Va. Dec., 1914.

WILLIAM JACOB HOLLAND, Carnegie Museum, Pittsburgh, Pa. December, 1910.
 ARTHUR HOLLICK, Staten Island Association of Arts and Sciences, New Brighton, S. I. August, 1898.

THOMAS C. HOPKINS, Syracuse University, Syracuse, N. Y. December, 1894.

WILLIAM OTIS HOTCHKISS, State Geologist, Madison, Wis. December, 1911.

*EDMUND OTIS HOVEY, American Museum of Natural History, New York, N. Y.

ERNEST HOWE, 77 Rhode Island Ave., Newport, R. I. December, 1903.

GEORGE D. HUBBARD, Oberlin College, Oberlin, Ohio. December, 1914.

LUCIUS L. HUBBARD, Houghton, Mich. December, 1894.

WALTER F. HUNT, University of Michigan, Ann Arbor, Mich. December, 1914.

ELLSWORTH HUNTINGTON, 222 Highland St., Milton, Mass.

LOUIS HUSSAKOF, American Museum of Natural History, New York, N. Y. December, 1910.

JOSEPH P. IDDINGS, Brinklow, Md. May, 1889.

JOHN D. IRVING, Yale University, New Haven, Conn. December, 1905.

A. WENDELL JACKSON, 9 Desbrosses St., New York, N. Y. December, 1888.

ROBERT T. JACKSON, 195 Bay State Road, Boston, Mass. August, 1894.

THOMAS AUGUSTUS JAGGAR, JR., Hawaiian Volcano Observatory, Territory of Hawaii, U. S. A. December, 1906.

MARK S. W. JEFFERSON, Michigan State Normal College, Ypsilanti, Mich. December, 1904.

EDWARD C. JEFFREY, Harvard University, Cambridge, Mass. December, 1914.

ALBERT JOHANNSEN, University of Chicago, Chicago, Ill. December, 1908.

DOUGLAS WILSON JOHNSON, Columbia University, New York, N. Y. Dec., 1906.

ALEXIS A. JULIEN, South Harwich, Mass. May, 1889.

FRANK JAMES KATZ, U. S. Geological Survey, Washington, D. C. Dec., 1912.

GEORGE FREDERICK KAY, State University of Iowa, Iowa City, Iowa. Dec., 1908.

ARTHUR KEITH, U. S. Geological Survey, Washington, D. C. May, 1889.

*JAMES F. KEMP, Columbia University, New York, N. Y.

CHARLES ROLLIN KEYES, 944 Fifth St., Des Moines, Iowa. August, 1890.

EDWARD M. KINDLE, Victoria Memorial Museum, Ottawa, Canada. Dec., 1905.

CHARLES TOWNSEND KIRK, University of New Mexico, Albuquerque, New Mexico. December, 1915.

EDWIN KIRK, U. S. Geological Survey, Washington, D. C. December, 1912.

CYRIL WORKMAN KNIGHT, Toronto, Ontario, Canada. December, 1911.

ADOLPH KNOPF, U. S. Geological Survey, Washington, D. C. December, 1911.

FRANK H. KNOWLTON, U. S. National Museum, Washington, D. C. May, 1889.

EDWARD HENRY KRAUS, University of Michigan, Ann Arbor, Mich. June, 1902.

HENRY B. KÜMMEL, Trenton, N. J. December, 1895.

*GEORGE F. KUNZ, 401 Fifth Ave., New York, N. Y.

GEORGE EDGAR LADD, State College, N. M. August, 1891.

LAWRENCE MORRIS LAMBE, Department of Mines, Ottawa, Canada. Dec., 1911.

HENRY LANDES, University of Washington, University Station, Seattle, Wash. December, 1908.

ALFRED C. LANE, Tufts College, Mass. December, 1889.

ESPER S. LARSEN, JR., U. S. Geological Survey, Washington, D. C. Dec., 1914.

ANDREW C. LAWSON, University of California, Berkeley, Cal. May, 1889.

WILLIS THOMAS LEE, U. S. Geological Survey, Washington, D. C. Dec., 1903.

JAMES H. LEES, Iowa Geological Survey, Des Moines, Iowa. December, 1914.

- CHARLES K. LEITH, University of Wisconsin, Madison, Wis. Dec., 1902.
 ARTHUR G. LEONARD, State University of North Dakota, Grand Forks, N. Dak.
 December, 1901.
 FRANK LEVERETT, Ann Arbor, Mich. August, 1890.
 JOSEPH VOLNEY LEWIS, Rutgers College, New Brunswick, N. J. Dec., 1906.
 WILLIAM LIBBEY, Princeton University, Princeton, N. J. August, 1899.
 WALDEMAR LINDGREN, Massachusetts Institute of Technology, Boston, Mass.
 August, 1890.
 MIGUEL A. R. LISBOA, Irrigation and Water Supply Service, Rio de Janeiro,
 Brazil. December, 1913.
 FREDERICK BREWSTER LOOMIS, Amherst College, Amherst, Mass. Dec., 1909.
 GEORGE DAVIS LOUDERBACK, University of California, Berkeley, Cal. June, 1902.
 ROBERT H. LOUGHRIDGE, University of California, Berkeley, Cal. May, 1889.
 ALBERT P. LOW, Department of Mines, Ottawa, Canada. December, 1905.
 RICHARD SWANN LULL, Yale University, New Haven, Conn. December, 1909.
 SAMUEL WASHINGTON MCCALLIE, Atlanta, Ga. December, 1909.
 HIRAM DEYER MCCASKEY, U. S. Geological Survey, Washington, D. C. De-
 cember, 1904.
 RICHARD G. MCCONNELL, Geological and Natural History Survey of Canada,
 Ottawa, Canada. May, 1889.
 DONALD FRANCIS MACDONALD, U. S. Geological Survey, Washington, D. C.
 December, 1915.
 JAMES RIEMAN MACFARLANE, Woodland Road, Pittsburgh, Pa. August, 1891.
 WILLIAM MCINNES, Geological and Natural History Survey of Canada, Ot-
 tawa, Canada. May, 1889.
 PETER MCKELLAR, Fort William, Ontario, Canada. August, 1890.
 GEORGE ROGERS MANSFIELD, 2067 Park Road N. W., Washington, D. C. De-
 cember, 1909.
 CURTIS F. MARBUT, Bureau of Soils, Washington, D. C. August, 1897.
 VERNON F. MARSTERS, San Juancito, Honduras, C. A. August, 1892.
 GEORGE CURTIS MARTIN, U. S. Geological Survey, Washington, D. C. June, 1902.
 LAWRENCE MARTIN, University of Wisconsin, Madison, Wis. December, 1909.
 EDWARD B. MATHEWS, Johns Hopkins University, Baltimore, Md. Aug., 1895.
 FRANCOIS E. MATTHES, U. S. Geological Survey, Washington, D. C. Decem-
 ber, 1914.
 W. D. MATTHEW, American Museum of Natural History, New York, N. Y.
 December, 1903.
 THOMAS POOLE MAYNARD, 1622 D. Hurt Bldg., Atlanta, Ga. December, 1914.
 P. H. MELL, 165 East 10th St., Atlanta, Ga. December, 1888.
 WALTER C. MENDENHALL, U. S. Geological Survey, Washington, D. C. June,
 1902.
 JOHN C. MERRIAM, University of California, Berkeley, Cal. August, 1895.
 *FREDERICK J. H. MERRILL, 631 Higgins Bldg., Los Angeles, Cal.
 GEORGE P. MERRILL, U. S. National Museum, Washington, D. C. Dec., 1888.
 HERBERT E. MERWIN, Geophysical Laboratory, Washington, D. C. Dec., 1914.
 ARTHUR M. MILLER, State University of Kentucky, Lexington, Ky. Dec., 1897.
 BENJAMIN L. MILLER, Lehigh University, South Bethlehem, Pa. Dec., 1904.
 WILLET G. MILLER, Toronto, Canada. December, 1902.
 WILLIAM JOHN MILLER, Smith College, Northampton, Mass. December, 1909.

- FRED HOWARD MOFFIT, U. S. Geological Survey, Washington, D. C. Dec., 1912.
 G. A. F. MOLENGRAAF, Technical High School, Delft, Holland. December, 1913.
 HENRY MONTGOMERY, University of Toronto, Toronto, Canada. Dec., 1904.
 ELWOOD S. MOORE, Pennsylvania State College, State College, Pa. Dec., 1911.
 MALCOLM JOHN MUNN, Clinton Bldg., Tulsa, Okla. December, 1909.
 *FRANK L. NASON, West Haven, Conn.
 DAVID HALE NEWLAND, Albany, N. Y. December, 1906.
 JOHN F. NEWSOM, Leland Stanford, Jr., University, Stanford University, Cal.
 December, 1899.
 WILLIAM H. NORTON, Cornell College, Mount Vernon, Iowa. December, 1895.
 CHARLES J. NORWOOD, State University, Lexington, Ky. August, 1894.
 IDA HELEN OGILVIE, Barnard College, Columbia University, New York, N. Y.
 December, 1906.
 CLEOPHAS C. O'HARRA, South Dakota School of Mines, Rapid City, S. Dak.
 December, 1904.
 DANIEL WEBSTER OHERN, University of Oklahoma, Norman, Okla. Dec., 1911.
 EZEQUIEL ORDONEZ, 2 a General Prim 43, Mexico, D. F., Mex. August, 1896.
 EDWARD ORTON, Jr., Geological Survey of Ohio, Columbus, Ohio. Dec., 1909.
 HENRY F. OSBORN, American Museum of Natural History, New York, N. Y.
 August, 1894.
 SIDNEY PAIGE, U. S. Geological Survey, Washington, D. C. December, 1911.
 CHARLES PALACHE, Harvard University, Cambridge, Mass. August, 1897.
 WILLIAM A. PARKS, University of Toronto, Toronto, Canada. December, 1906.
 *HORACE B. PATTON, Colorado School of Mines, Golden, Colo.
 FREDERICK B. PECK, Lafayette College, Easton, Pa. August, 1901.
 RICHARD A. F. PENROSE, JR., 460 Bullitt Bldg., Philadelphia, Pa. May, 1889.
 GEORGE H. PERKINS, University of Vermont, Burlington, Vt.; State Geologist.
 June, 1902.
 JOSEPH H. PERRY, 276 Highland St., Worcester, Mass. December, 1888.
 OLAF AUGUST PETERSON, Carnegie Museum, Pittsburgh, Pa. December, 1910.
 WILLIAM CLIFTON PHALEN, U. S. Geological Survey, Washington, D. C. De-
 cember, 1912.
 ALEXANDER H. PHILLIPS, Princeton University, Princeton, N. J. Dec., 1914.
 LOUIS V. PISSON, Yale University, New Haven, Conn. August, 1894.
 JOSEPH E. POGUE, Northwestern University, Evanston, Ill. December, 1911.
 JOSEPH HYDE PRATT, North Carolina Geological Survey, Chapel Hill, N. C.
 December, 1898.
 LOUIS M. PRINDLE, U. S. Geological Survey, Washington, D. C. Dec., 1912.
 *CHARLES S. PROSSER, Ohio State University, Columbus, Ohio.
 WILLIAM FREDERICK PROUTY, University of Alabama, University, Ala. De-
 cember, 1911.
 *RAPHAEL PUMPELLY, Newport, R. I.
 ALBERT HOMER PURDUE, State Geological Survey, Nashville, Tenn. Dec., 1904.
 FREDERICK LESLIE RANSOME, U. S. Geological Survey, Washington, D. C. Au-
 gust, 1895.
 PERCY EDWARD RAYMOND, Museum of Comparative Zoölogy, Cambridge, Mass.
 December, 1907.
 CHESTER A. REEDS, American Museum of Natural History, New York, N. Y.
 December, 1913.

- HARRY FIELDING REID, Johns Hopkins University, Baltimore, Md. Dec., 1892.
 WILLIAM NORTH RICE, Wesleyan University, Middletown, Conn. August, 1890.
 JOHN LYON RICH, University of Illinois, Urbana, Ill. December, 1912.
 CHARLES H. RICHARDSON, Syracuse University, Syracuse, N. Y. Dec., 1899.
 GEORGE BURR RICHARDSON, U. S. Geological Survey, Washington, D. C. December, 1908.
 HEINRICH RIES, Cornell University, Ithaca, N. Y. December, 1893.
 ELMER S. RIGGS, Field Museum of Natural History, Chicago, Ill. Dec., 1911.
 JESSE PERRY ROWE, University of Montana, Missoula, Mont. December, 1911.
 RUDOLPH RUEDEMANN, Albany, N. Y. December, 1905.
 JOHN JOSEPH RUTLEDGE, Experiment Station, Pittsburgh, Pa. Dec., 1911.
 ORESTES H. ST. JOHN, 1141 Twelfth St., San Diego, Cal. May, 1889.
 *ROLLIN D. SALISBURY, University of Chicago, Chicago, Ill.
 FREDERICK W. SARDESON, University of Minnesota, Minneapolis, Minn. December, 1892.
 THOMAS EDMUND SAVAGE, University of Illinois, Urbana, Ill. December, 1907.
 FRANK C. SCHRADER, U. S. Geological Survey, Washington, D. C. Aug., 1901.
 CHARLES SCHUCHERT, Yale University, New Haven, Conn. August, 1895.
 ALFRED REGINALD SCHULTZ, U. S. Geological Survey, Washington, D. C. December, 1912.
 WILLIAM B. SCOTT, Princeton University, Princeton, N. J. August, 1892.
 ARTHUR EDMUND SEAMAN, Michigan College of Mines, Houghton, Mich. December, 1904.
 HENRY M. SEELY, Middlebury College, Middlebury, Vt. May, 1889.
 ELIAS H. SELLARDS, Tallahassee, Fla. December, 1905.
 JOAQUIM CANDIDO DA COSTA SEÑA, State School of Mines, Ouro Preto, Brazil. December, 1908.
 MILLARD K. SHALER, 4 Bishopsgate E. C., London, England. December, 1914.
 GEORGE BURBANK SHATTUCK, Vassar College, Poughkeepsie, N. Y. Aug., 1899.
 EUGENE WESLEY SHAW, U. S. Geological Survey, Washington, D. C. Dec., 1912.
 SOLON SHEDD, State College of Washington, Pullman, Wash. Dec., 1904.
 EDWARD M. SHEPARD, 1403 Benton Ave., Springfield, Mo. August, 1901.
 BOHUMIL SHIMEK, University of Iowa, Iowa City, Iowa. December, 1904.
 HERVEY WOODBURN SHIMER, Massachusetts Institute of Technology, Boston, Mass. December, 1910.
 CLAUDE ELLSWORTH SIEBENTHAL, U. S. Geological Survey, Washington, D. C. December, 1912.
 *FREDERICK W. SIMONDS, University of Texas, Austin, Texas.
 WILLIAM JOHN SINCLAIR, Princeton University, Princeton, N. J. Dec., 1906.
 JOSEPH THEOPHILUS SINGEWALD, Johns Hopkins University, Baltimore, Md. December, 1911.
 EARLE SLOAN, Charleston, S. C. December, 1908.
 BURNETT SMITH, Syracuse University, Skaneateles, N. Y. December, 1911.
 CARL SMITH, U. S. Geological Survey, Washington, D. C. December, 1912.
 *EUGENE A. SMITH, University of Alabama, University, Ala.
 GEORGE OTIS SMITH, U. S. Geological Survey, Washington, D. C. Aug., 1897.
 PHILIP S. SMITH, U. S. Geological Survey, Washington, D. C. Dec., 1909.
 WARREN DU PRÉ SMITH, University of Oregon, Eugene, Oregon. Dec., 1909.
 W. S. TANGIER SMITH, Lodi, Cal. June, 1902.

*JOHN C. SMOCK, Trenton, N. J.

CHARLES H. SMYTH, JR., Princeton University, Princeton, N. J. Aug., 1892.

HENRY L. SMYTH, Harvard University, Cambridge, Mass. August, 1894.

ARTHUR COE SPENCER, U. S. Geological Survey, Washington, D. C. Dec., 1896.

*J. W. SPENCER, 2019 Hillyer Place, Washington, D. C.

FRANK SPRINGER, U. S. National Museum, Washington, D. C. December, 1911.

JOSIAH E. SPURR, Bullitt Bldg., Philadelphia, Pa. December, 1894.

JOSEPH STANLEY-BROWN, 26 Exchange Place, New York, N. Y. August, 1892.

TIMOTHY WILLIAM STANTON, U. S. National Museum, Washington, D. C. August, 1891.

CLINTON RAYMOND STAUFFER, University of Minnesota, Minneapolis, Minn. December, 1911.

LLOYD WILLIAM STEPHENSON, U. S. Geological Survey, Washington, D. C. December, 1911.

*JOHN J. STEVENSON, 215 West 101st St., New York, N. Y.

RALPH WALTER STONE, U. S. Geological Survey, Washington, D. C. Dec., 1912.

GEORGE WILLIS STOSE, U. S. Geological Survey, Washington, D. C. Dec., 1908.

CHARLES KEPHART SWARTZ, Johns Hopkins University, Baltimore, Md. December, 1908.

STEPHEN TABER, University of South Carolina, Columbia, S. C. Dec., 1914.

JOSEPH A. TAFF, 781 Flood Building, San Francisco, Cal. August, 1895.

MIGNON TALBOT, Mount Holyoke College, South Hadley, Mass. Dec., 1913.

JAMES E. TALMAGE, University of Utah, Salt Lake City, Utah. Dec., 1897.

FRANK B. TAYLOR, Fort Wayne, Ind. December, 1895.

*JAMES E. TODD, 1224 Rhode Island St., Lawrence, Kans.

CYRUS FISHER TOLMAN, JR., Leland Stanford, Jr., University, Stanford University, Cal. December, 1909.

ARTHUR C. TROWBRIDGE, State University of Iowa, Iowa City, Iowa. December, 1913.

*HENRY W. TURNER, 209 Alaska Commercial Building, San Francisco, Cal.

WILLIAM H. TWENHOFEL, University of Kansas, Lawrence, Kans. Dec., 1913.

MAYVILLE WILLIAM TWITCHELL, State Geological Survey, Trenton, N. J. December, 1911.

JOSEPH B. TYRRELL, Room 534, Confederation Life Building, Toronto, Canada. May, 1889.

JOHAN A. UDDEN, University of Texas, Austin, Texas. August, 1897.

EDWARD O. ULRICH, U. S. Geological Survey, Washington, D. C. Dec., 1903.

JOSEPH B. UMPLEBY, U. S. Geological Survey, Washington, D. C. Dec., 1913.

*WARREN UPHAM, Minnesota Historical Society, Saint Paul, Minn.

*CHARLES R. VAN HISE, University of Wisconsin, Madison, Wis.

FRANK ROBERTSON VAN HORN, Case School of Applied Science, Cleveland, Ohio. December, 1898.

GILBERT VAN INGEN, Princeton University, Princeton, N. J. December, 1904.

THOMAS WAYLAND VAUGHAN, U. S. Geological Survey, Washington, D. C. August, 1896.

ARTHUR CLIFFORD VEACH, 7 Richmond Terrace, Whitehall, S. W., London, England. December, 1906.

*ANTHONY W. VOGDES, 2425 First St., San Diego, Cal.

- *M. EDWARD WADSWORTH, School of Mines, University of Pittsburgh, Pittsburgh, Pa.
- *CHARLES D. WALCOTT, Smithsonian Institution, Washington, D. C.
- THOMAS L. WALKER, University of Toronto, Toronto, Canada. Dec., 1903.
- CHARLES H. WARREN, Massachusetts Institute of Technology, Boston, Mass. December, 1901.
- HENRY STEPHENS WASHINGTON, Geophysical Laboratory, Washington, D. C. August, 1896.
- THOMAS L. WATSON, University of Virginia, Charlottesville, Va. June, 1900.
- CHARLES E. WEAVER, University of Washington, Seattle, Wash. Dec., 1913.
- WALTER H. WEED, 29 Broadway, New York, N. Y. May, 1889.
- CARROLL HARVEY WEGEMANN, U. S. Geological Survey, Washington, D. C. December, 1912.
- SAMUEL WEIDMAN, Wisconsin Geological and Natural History Survey, Madison, Wis. December, 1903.
- STUART WELLER, University of Chicago, Chicago, Ill. June, 1900.
- LEWIS G. WESTGATE, Ohio Wesleyan University, Delaware, Ohio. Aug., 1894.
- EDGAR THEODORE WHERRY, U. S. National Museum, Washington, D. C. December, 1915.
- DAVID WHITE, U. S. National Museum, Washington, D. C. May, 1889.
- *ISRAEL C. WHITE, Morgantown, W. Va.
- GEORGE REBER WIELAND, Yale University, New Haven, Conn. December, 1910.
- FRANK A. WILDER, North Holston, Smyth County, Va. December, 1905.
- *EDWARD H. WILLIAMS, JR., Woodstock, Vt.
- *HENRY S. WILLIAMS, Cornell University, Ithaca, N. Y.
- IRA A. WILLIAMS, Oregon School of Mines, Corvallis, Ore. December, 1905.
- BAILEY WILLIS, Leland Stanford, Jr., University, Cal. December, 1889.
- ALFRED W. G. WILSON, Department of Mines, Ottawa, Canada. June, 1902.
- ALEXANDER N. WINCHELL, University of Wisconsin, Madison, Wis. Aug., 1901.
- *HORACE VAUGHN WINCHELL, 826 First National Society Bldg., Minneapolis, Minn.
- *ARTHUR WINSLOW, 131 State St., Boston, Mass.
- JOHN E. WOLFF, Harvard University, Cambridge, Mass. December, 1889.
- JOSEPH E. WOODMAN, New York University, New York, N. Y. Dec., 1905.
- ROBERT S. WOODWARD, Carnegie Institution of Washington, Washington, D. C. May, 1889.
- JAY B. WOODWORTH, Harvard University, Cambridge, Mass. December, 1895.
- CHARLES WILL WRIGHT, Ingurtozu, Arbus, Sardinia, Italy. December, 1909.
- FREDERIC E. WRIGHT, Geophysical Laboratory, Carnegie Institution, Washington, D. C. December, 1903.
- *G. FREDERICK WRIGHT, Oberlin Theological Seminary, Oberlin, Ohio.
- GEORGE A. YOUNG, Geological Survey of Canada, Ottawa, Canada. Dec., 1905.

CORRESPONDENTS DECEASED

HERMAN CREDNER. Died July 22, 1913.	EDWARD SUSS. Died April 20, 1914.
A. MICHEL-LÉVY. Died September, 1911.	TH. TSCHERNYSCHEW. Died Jan. 15, 1914.
H. ROSENBUSCH. Died January 20, 1914.	FERDINAND ZIRKEL. Died June 11, 1912.

FELLOWS DECEASED

* Indicates Original Fellow (see article III of Constitution)

- *CHAS. A. ASHBURNER. Died Dec. 24, 1889.
 ALFRED E. BARLOW. Died May 28, 1914.
 CHARLES E. BEECHER. Died Feb. 14, 1904.
 ALBERT S. BICKMORE. Died Aug. 12, 1914.
 WM. PHIPPS BLAKE. Died May 21, 1910.
 AMOS BOWMAN. Died June 18, 1894.
 ERNEST R. BUCKLEY. Died Jan. 19, 1912.
 *SAMUEL CALVIN. Died April 17, 1911.
 FRANKLIN R. CARPENTER. Died April 1, 1910.
 *J. H. CHAPIN. Died March 14, 1892.
 *EDWARD W. CLAYPOLE. Died Aug. 17, 1901.
 *THEO. B. COMSTOCK. Died July 26, 1915.
 GEORGE H. COOK. Died Sept. 22, 1889.
 *EDWARD D. COPE. Died April 12, 1897.
 ANTONIO DEL CASTILLO. Died Oct. 28, 1895.
 *JAMES D. DANA. Died April 14, 1895.
 GEORGE M. DAWSON. Died March 2, 1901.
 SIR J. WM. DAWSON. Died Nov. 19, 1899.
 ORVILLE A. DERBY. Died Nov. 27, 1915.
 CLARENCE E. DUTTON. Died Jan. 4, 1912.
 *WILLIAM B. DWIGHT. Died Aug. 29, 1906.
 *GEORGE H. ELDRIDGE. Died June 29, 1905.
 *SAMUEL F. EMMONS. Died March 28, 1911.
 WM. M. FONTAINE. Died April 29, 1913.
 *ALBERT R. FOOTE. Died October 10, 1895.
 *PERSIFOR FRAZER. Died April 7, 1909.
 *HOMER T. FULLER. Died Aug. 14, 1908.
 N. J. GIROUX. Died November 30, 1891.
 *CHRISTOPHER W. HALL. Died May 10, 1911.
 *JAMES HALL. Died August 7, 1898.
 JOHN B. HATCHER. Died July 3, 1904.
 *ROBERT HAY. Died December 14, 1895.
 C. WILLARD HAYES. Died Feb. 9, 1916.
 *ANGELO HEILPRIN. Died July 17, 1907.
 FRANK A. HILL. Died July 13, 1915.
 *JOSEPH A. HOLMES. Died July 13, 1915.
 DAVID HONEYMAN. Died October 17, 1889.
 *EDWIN E. HOWELL. Died April 16, 1911.
 *HORACE C. HOVEY. Died July 27, 1914.
 THOMAS S. HUNT. Died Feb. 12, 1892.
 *ALPHEUS HYATT. Died Jan. 15, 1902.
 THOMAS M. JACKSON. Died Feb. 3, 1912.
 *JOSEPH F. JAMES. Died March 29, 1897.
 WILBUR C. KNIGHT. Died July 28, 1903.
 RALPH D. LACOE. Died February 5, 1901.
 J. C. K. LAFLAMME. Died July 6, 1910.
 DANIEL W. LANGTON. Died June 21, 1909.
 *JOSEPH LE CONTE. Died July 6, 1901.
 *J. PETER LESLEY. Died June 2, 1903.
 HENRY MCCALLEY. Died Nov. 20, 1904.
 *W J MCGEE. Died September 4, 1912.
 OLIVER MARCY. Died March 19, 1899.
 OTHNIEL C. MARSH. Died March 18, 1899.
 JAMES E. MILLS. Died July 25, 1901.
 *HENRY B. NASON. Died January 17, 1895.
 *PETER NEFF. Died May 11, 1903.
 *JOHN S. NEWBERRY. Died Dec. 7, 1892.
 WILLIAM H. NILES. Died Sept. 12, 1910.
 *EDWARD ORTON. Died October 16, 1899.
 *AMOS O. OSBOEN. Died March, 1911.
 *RICHARD OWEN. Died March 24, 1890.
 SAMUEL L. PENFIELD. Died Aug. 14, 1906.
 DAVID P. PENHALLOW. Died Oct. 20, 1910.
 *FRANKLIN PLATT. Died July 24, 1900.
 WILLIAM H. PETTEE. Died May 26, 1904.
 *JOHN W. POWELL. Died Sept. 23, 1902.
 *ISRAEL C. RUSSELL. Died May 1, 1906.
 *JAMES M. SAFFORD. Died July 3, 1907.
 *CHARLES SCHAEFFER. Died Nov. 23, 1903.
 *NATHANIEL S. SHALER. Died April 10, 1906.
 WILLIAM J. SUTTON. Died May 9, 1915.
 RALPH S. TARR. Died March 21, 1912.
 WILLIAM G. TIGHT. Died Jan. 15, 1910.
 CHARLES WACHSMUTH. Died Feb. 7, 1896.
 THOMAS C. WESTON. Died July 20, 1910.
 THEODORE G. WHITE. Died July 7, 1901.
 *ROBERT P. WHITFIELD. Died April 6, 1910.
 *GEORGE H. WILLIAMS. Died July 12, 1894.
 *J. FRANCIS WILLIAMS. Died Nov. 9, 1891.
 ARTHUR B. WILMOTT. Died May 8, 1914.
 *ALEXANDER WINCHELL. Died Feb. 19, 1891.
 *NEWTON WINCHELL. Died May 1, 1914.
 ALBERT A. WRIGHT. Died April 2, 1905.
 WILLIAM S. YEATES. Died Feb. 19, 1908.

Summary

Correspondents	10
Original Fellows.....	44
Elected Fellows.....	335
<hr/>	
Membership.....	389
Deceased Correspondents.....	6
Deceased Fellows.....	85