they are mingled in privies," and Sir Richard Thorne, his successor, has recently remarked that the midden-privy provides almost every factor "favourable to the setting up and maintenance of those very conditions which seem to be essential to the vitality and multiplication of the typhoid bacillus." The value of these warnings has, moreover been established beyond all question by Professor Delépine, who in a characteristically ingenious manner demonstrated the presence of swarms of Eberth's bacillus in the detritus from between the floor bricks of a privy-pit into which typhoid dejecta had thirteen months before been thrown, the receptacle having been repeatedly subjected to careful disinfecting processes in the interval. To this striking bacteriological proof may be added an abundance of the "evidence of experience" of midden-privy towns in regard to typhoid fever.

In the appended table are set forth the statistics for Stockport during the five years 1893-97. In order to make the inquiry as equitable as possible the privy-pit houses and water-closet houses have, with much labour, been enumerated and classified from the rate-books, and the incidence of typhoid fever on each class compared according to rateable value, which has been taken, partly at the suggestion of Professor Delépine, as a fair indication of the sanitary condition in other respects of the two classes of dwellings.

Table showing for the Five Years 1893-97 the Incidence of Typhoid Fever in Stockport on Privy-pit Houses and Water-closet Houses respectively.

Rateable value.	Privy-pit houses.		Water-closet houses.		Typhoid fever incidence per centum of	
	Average number.	Cases of typhoid fever notified.	Average number.	Cases of typhoid fever notified.	Privy-pit houses.	Water-closet houses.
Under £5	2,811	54	176	1	1.92	0.56
£5 and under £8...	7,771	275	703	3	3.53	0.42
£8 and under £10	1,415	57	746	9	4.02	1.20
£10 and under £12	1,077	42	425	8	3.89	1.88
£12 and under £14	449	16	225	4	3.56	1.77
£14 and under £20	656	30	613	8	4.57	1.30
£20 and under £30	249	13	374	2	5.22	0.53
£30 and upwards...	147	8	369	9*	5.44	2.43*
—	14,575	495	3631	44*	3.39	1.21*

This figure does not include 2 cases reported from the infirmary and 4 from the workhouse hospital.

In 72 instances during the five years infection of a privy-pit with typhoid discharges was followed by one or more cases amongst those who subsequently used the privies. The entire number of such subsequent cases was 116 out of a total of 545 notified, or 21.3 per cent., and as many as 4, 5, 7, and 8 cases were upon occasion traced to the same privy-pit. The details are as follows. In 53 instances 1 subsequent case occurred—viz., in 37 cases within a few weeks, in 10 cases about one year later, in 3 cases about two years later, in 1 case about three years later, and in 2 cases about four years later. In 10 instances 2 subsequent cases occurred—viz., in 7 cases within a few weeks, in 1 case about one year later, and in 2 cases about two years later. In 2 instances 3 subsequent cases occurred—viz., in 1 case within a few weeks and in 1 case about a year later. In 2 instances 4 subsequent cases occurred—viz., 1 within a few weeks and 1 about three years later. In 3 instances 5 subsequent cases occurred—viz., 2 were within a few weeks and the other four years later. In 1 instance 7, and in another 8, subsequent cases occurred within a few weeks. During the same period only 1 subsequent case (and that was three years later) occurred where a water-closet was used. This was in a very large public institution, and it is doubtful whether the water-closet infected by the first sufferer was used by the second.

It is noteworthy that the figures for Stockport show that the incidence of typhoid fever increases with rateable value both in privy-pit and water-closet houses.

Stockport.

A CASE OF
CHRONIC CONSTIPATION ENDING
FATALLY AND ASSOCIATED WITH
ENORMOUS DILATATION OF THE
SIGMOID FLEXURE.

By PEVERELL S. HICHENS, M.A., M.B., B.Ch. Oxon.,
M.R.C.S. Eng., L.R.C.P. Lond.,

ASSISTANT RESIDENT MEDICAL OFFICER TO THE HOSPITAL FOR CONSUMPTION AND DISEASES OF THE CHEST, BROMPTON.

THE patient was a young man, aged twenty years. The account of the case is necessarily rather imperfect, as he was not a patient of any of the physicians at the hospital, but was treated by his own medical man outside. The history of the case showed that from the day of his birth to the day of his death the patient suffered from constipation and his bowels were apparently never opened without recourse to artificial means. For the first week of his life he was exceedingly ill, passed nothing but blood and water by the rectum, and was not expected to live. After the first week he began passing faecal matter, but the motions were never those proper to an infant, consisting almost entirely of scybala, and they were passed with much pain and screaming. The motions, such as they were, were not obtained without previous ½ oz. doses of castor oil, which often had to be repeated two or three times in the course of twelve hours before the bowels acted. As the child grew the bowels were only opened by drugs, and with increased difficulty, and at the age of twelve months the mother began to have recourse to ½ pint enemias of soap and water. Later enemias of a whole pint had to be given and very often had to be repeated two or three times before the bowels were opened. The after-history of the case is merely a repetition. The bowels used only to be opened at intervals of ten days or a fortnight and only after repeated enemias. A very large scybalous motion was then passed, generally succeeded by several loose motions, spread over the next two or three days. The motions could only be passed by the help of gravity—i.e., squatting down over a chamber in the position which, as Dr. Lauder Brunton has pointed out, is the natural and proper one for defecation. The patient very often felt slightly sick before the bowels were opened and was very much collapsed afterwards, so much so that he often had to go to bed for the rest of the day. His abdomen was always greatly distended, so that he could never button the bottom button of his waistcoat or the top button of his trousers. About a month before he died he was taken ill with influenza and on getting better he went away into the country for a holiday. Five days before death he was seized with pains all over his body and slight swelling of the legs, and in consequence he hurried home. He was treated by a medical man who thought that he had a slight touch of rheumatic fever. The night before he died he passed a fairly large motion. On the morning of the next day he was seized with a severe pain over the heart for which the præcordial area was rubbed with liniment, which gave him considerable relief, and shortly afterwards he went to bed feeling fairly comfortable. During the following night he suddenly got out of bed for some unknown reason and fell down dead.

The body was brought to the hospital, and at the necropsy it was found that it was well developed and that rigor mortis was not present. There was very slight oedema over the feet and shins, but enormous general distension of the abdomen. On reflecting the parietes the abdominal cavity was found to be occupied by a tense shining viscus, presenting the appearance of a sac rising out of the pelvis and passing under the ribs, where it doubled on itself and returned to the pelvis again. On closer inspection this proved to be an enormously distended sigmoid flexure. At the point where the sigmoid flexure left the descending colon it turned on itself and passed directly up the left side of the abdomen. The summit of the viscus then passed behind the ribs and xiphisternum with a gentle curvature and descended along the right side of the abdomen to the pelvis to join the rectum. Some little distance above the junction with the rectum the viscus showed a distinct constriction. The remaining abdominal viscera were entirely concealed by the sigmoid flexure, which pushed the liver upwards and backwards, compressed the lungs, and rotated

the heart upwards and outwards. On removing the sigmoid flexure and laying it open it was found to contain an enormous amount of gas and also a large quantity of semi-liquid fæces of the consistence of pea-soup. The total length of the sigmoid flexure when it was opened and laid flat was $22\frac{1}{2}$ inches. Fourteen inches from its upper end was a large cicatrix formed by an almost healed ulcer, probably stercoral in origin, which had caused the constriction above mentioned. The circumference of the flexure above the ulcer was 14 inches, at the ulcer it was $7\frac{3}{4}$ inches, and below the ulcer it was 10 inches. These measurements, of course, represent the circumference of the flexure at its period of least distension, and they also show that the constriction at the site of the ulcer was merely a relative one and could have made no difference, or very little, to the onward passage of the fæces. The walls of the sigmoid flexure were uniformly greatly thickened. This thickening was shown on microscopical examination to be almost entirely due to great hypertrophy of the circular and longitudinal muscular fibres. There was also slight thickening of the submucosa. As regards the rest of the organs, the pharynx, œsophagus, and stomach were quite healthy. The small intestine also showed no lesion and was not at all distended. There was slight general distension of the cæcum and of the ascending, the transverse, and the descending colon. This distension was not at all marked and the large intestine appeared otherwise healthy, as did also the rectum, and there was no sign of a fibrous band, kink, or constriction anywhere except at the ulcer above mentioned. The lungs were small and compressed by the abdominal distension and were slightly œdematous and engorged. The heart was somewhat displaced but was otherwise perfectly sound, as also were the liver and spleen. The cortices of the kidneys were slightly swollen and streaky and on microscopical examination the renal epithelium appeared to be in a state of cloudy swelling.

The disease in this case would appear to come under the heading of so-called "idiopathic dilatation of the colon." Dr. Hale White, in his excellent article on the subject in Allbutt's System of Medicine,¹ mentions that only 12 cases have been recorded in the last forty years, and he provisionally groups them under two heads—those occurring in elderly patients and those occurring in young children due to some congenital inertness of the bowels. Whatever may be the initial cause—whether it was due to a temporary kink, as Rolleston and Haward² suggest, or to the length of the sigmoid in children readily allowing kinking, as Jacobi³ suggests, or to mere constipation and distension with wind or local spasm of the rectum, as Gee⁴ suggests, the attempted remedies in this case must probably have increased the disease. The patient moved in a vicious circle and the sigmoid, already distended, was ever being more distended by the successive enemas administered. So far as could be judged from the post-mortem appearances the only possible treatment which could have had any permanent success would have been a lumbar colotomy.

Brompton.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A NOTE ON PTOMAIN POISONING.

By JOHN CAHILL, M.D. DURH., F.R.C.S. ENG.

THE subjoined instance of multiple poisoning by apparently wholesome food has occurred under my direct observation and may be considered worthy of record.

On Oct. 13th a brace of wild ducks, which appeared to be in very good condition both before and after cooking, formed part of the dinner of a party of five. All five persons partook of the ducks without any ill effect. On the 14th a portion of wild duck, cooked the day before, was eaten at dinner by two persons, appeared very palatable, and produced no ill

effect. On the 15th the bones of the same ducks were placed by the cook in her stock-pot together with the meat for soup. At dinner on that evening there were three persons, of whom one took no soup, one took soup only, and the third took the soup and other courses of the dinner which followed it. Within less than half an hour the two persons who had taken soup at table were seized with sickness and continued to vomit violently for about twenty minutes. The person who had partaken of the rest of the dinner, but not of the soup, was not in any way affected. The cook had tasted the soup by drinking about half a tea-cupful before it went up to table, and she was seized with sickness and vomiting after about the same interval and in the same manner as the two persons upstairs. The soup was obviously the cause of the sickness in these three cases, and there was the following evidence to show that the poisonous element in the soup proceeded from the wild duck, although these had been eaten without harmful result on the two preceding days and had appeared to be perfectly sweet when used as an addition to the soup. When the cook had sent the soup up to table it happened that she gave a bone of duck out of the pot to the cat. A little later, when the three cases of sudden sickness had occurred, inquiry was made as to the effect of this food upon the cat. That animal was found at the back of the house vomiting and evidently in great distress. None of the sufferers appeared the next morning to be any the worse for their experience.

Seville-street, S.W.

SUPPURATION IN AN OLD HERNIAL SAC.

By A. S. NAKASHIAN, M.D.

ON August 27th I was called to see a Turk, twenty-two years of age, who had had right inguinal hernia for ten years. The scrotum was now swollen, tense, and very tender, and the contents were irreducible. Ten days previously he had taken to his bed with chills, fever, and pain in the abdomen. Five days after the beginning of this illness (to use his own language) his "intestines came down and did not go back as usual." It seemed to me like strangulated hernia, but there was no vomiting and the bowels were acting quite regularly. On the second day I operated and evacuated about a pint of fetid pus from the hernial sac which did not contain any intestine. The inguinal canal was firmly closed and there was no communication with the abdominal cavity. I washed out the sac and established free drainage, after which he improved very much for a week. Then he began to get worse, having fever and complaining of abdominal pain. In a few days I detected very marked swelling and dullness in the lower third of the abdomen, which was tender and gave all the signs of containing pus, while the upper two thirds were flat, natural, and not tender. I did not open the abdomen but continued fomentations and tonics with occasional doses of opium to relieve him. By this treatment the swelling and fluid disappeared in eleven days. The man is now going about and transacting his business as formerly.

I think this was a case of tuberculous peritonitis accompanied with tuberculosis of the sac. The interesting features to me were the rapid closure of the sac, the partial peritonitis, and the rapid disappearance of the fluid and inflammation from the abdominal cavity.

Konia, Turkey.

MEMBRANOUS LARYNGITIS, WITH HYPERPYREXIA FROM MALARIAL POISON.

By CHARLES LLOYD WORRALL, L.R.C.P. LOND.

AT the cost of being considered long behind the times I am driven to designate the following as a case of "simple membranous" laryngitis in contra-distinction to diphtheritic laryngitis. This one is compelled to believe owing to the total absence of the usual symptoms of diphtheria. On Sept. 20th I was asked to see a child, two and a half years of age, who had been ailing for about five days, the symptoms notably being a very frequent cough of the characteristic "croupy" type with almost entire loss of voice. On examining the throat a very marked whitish membrane—not of the "wash-leather" type—was seen at the back of the throat implicating the larynx, the tonsils

¹ Allbutt's System of Medicine, vol. iii., pp. 967-972.

² Transactions of the Clinical Society of London, vol. xxix., 1895.

³ Archives of Pediatrics, vol. x.

⁴ St. Bartholomew's Hospital Reports, vol. xx.