Studies in Hepaticae: On *Pallavicinia* *decipiens*¹, Mitten.

BY

J. BRET LAND FARMER, M.A.,

*Fellow of Magdalen College, Oxford; Assistant Professor of Biology at the
Royal College of Science, South Kensington.*

—♦—
With Plates VI and VII.
—♦—

THE genus *Pallavicinia* (Gray), otherwise known as *Blytia*² (Endl.) or *Steetzia* (Lehm.), is one which includes rather more than twenty species, which exhibit, amongst themselves, a considerable range of variation in form. Most of them agree more or less with the type as illustrated by *P. Lyellii*, a plant not unfrequent in this country, and which consists of an elongated thallus, with a prominent mid-rib. Branching may occur dichotomously, or by ventrally-formed shoots. But specimens of *P. Lyellii* not unfrequently depart from the more usual form, in so far as the thallus-wings are often scarcely developed in the posterior region of the plant, which in this part comes thus to consist of little more than mid-rib. It is only as one passes to the more anterior portions nearer the apex, that the lateral expansions attain their full breadth. When however this feature is also accompanied by the formation of a few rapidly succeeding dichotomies in the apical region, the plant assumes an aspect quite foreign to the normal habitus of the species. It is, in these cases, not difficult

¹ This plant was described by Mitten (Journ. Linn. Soc., 1861) as *Steetzia decipiens*.

² In the Synopsis Hepaticarum the generic name is given as *Blyttia*; though Endlicher wrote it as in the text.

to conceive of the posterior almost wingless portion developing as a rhizome, whilst the expanded branch-system would form a 'frond.' This distinction into stipe and 'lamina' is actually met with in some species, e. g. *P. pisiformis*, where, to judge from herbarium specimens, the stipe forms a creeping stem, passing gradually into the 'lamina.' But this morphological differentiation of external form culminates in those species of which *P. decipiens* may be taken as an example.

In this plant there is a well marked creeping subterranean rhizome, from which arise expanded stalked aërial 'fronds' or dichotomous branch-systems. In luxuriant specimens, especially when dried, the general appearance strongly recalls that of some of the simpler Hymenophyllums, but in the fresh, living state, its succulent and turgid character forms a very striking and distinctive feature. It closely approximates in vegetative structure to many species of another genus of Hepatics, *Symphyogyna*, although it is readily distinguishable from them by the character of its fructification.

It is a fact worthy of mention, that within the limits of the genus *Symphyogyna* itself, much the same sort of transition from a simpler to a more complex type of vegetative structure may be traced. This latter genus has however outstripped *Pallavicinia*, in this respect, since it also exhibits considerable diversity in the *directions* of its complexity. Thus, besides exhibiting a gradual advance along lines parallel with those already indicated as occurring in *Pallavicinia*, in other groups of species the thallus becomes deeply indented, so as to become pinnatifid (*S. sinuata*), and so much so as to almost resemble one of the pseudo-foliose genera. It is however interesting to notice that on the same plant branches may arise whose margins are in no way sinuate, but resemble those of any common form in their regular outline; and on the other hand it is perhaps not improbable that in the toothed margins to be met with in some *Pallavicinias*, there is represented a rudimentary stage of that development which gives such a striking appearance to *Symphyogyna sinuata*.

Moreover there are a number of other genera which used to

be united either with *Pallavicinia* or with *Symphyogyna*, on account of their similarity in the vegetative structure of their gametophytes, but which are now separated owing to the important differences between their respective sexual organs or their sporophytes. In fact, similarity in the vegetative structure of the gametophyte, in many of these Hepatic genera at any rate, affords but unsafe ground on which to found any views as to natural affinities and mutual relationships. The existence of parallel developments, as above indicated, is a sufficient example of the soundness of this statement.

Pallavicinia decipiens has been described as occurring on the Horton Plains in Ceylon, and I found it abundantly in this locality, growing on the muddy banks of a stream at an elevation of about 6,200 feet. The specimens from this district, as well as others which I found on the summit of the Knuckles (6,000 ft.) further northwards, were comparatively stunted forms, though they were fruiting freely. At a lower elevation (4,500 ft.) the plant assumes a much more luxuriant form, and it is not uncommon at this elevation on the slopes of Adam's Peak, especially on the paths up from Maskeliya or Ratnapura. Here it grows in large patches and in such profusion as to take complete possession of the soil, to the exclusion of other plants.

Branching.—The plant consists, as has been already said, of an underground stem, and an aërial part consisting of fronds. The latter are formed simply as the ends of rhizome-branches, which, after growing horizontally in the earth for a time, bend upwards, and after emerging from the ground at once begin to flatten out, and branch in such a way as to give rise to the appearances of dichotomously branched fronds (Pl. VI, Fig. 1). At the outer side of the angle made by this upward curving of the underground stem, one or more rhizome-branches are given off, which penetrate the soil, and may either branch in it or at once turn upwards, and in their turn form fresh aërial fronds. But whatever the ultimate fate of the branches, they always arise exogenously throughout the plant.

In the aërial portions the branching which results in the

formation of the frond agrees with that form of dichotomy, so common in Hepatics, in which the original apex ceases to segment, and is replaced by two fresh apices arising right and left of it. This takes place repeatedly, and so the expanded branch-system or frond is formed. When an underground shoot is about to grow up out of the ground, it changes its shape, and from being circular in transverse section, it becomes elliptical, and wings appear on the two edges, which eventually spread out and form the laminae on either side of the mid-rib of the thallus.

The branching in the *subterranean* portion of the plant is very much complicated by the fact that the apical cells of lateral axes do not develop immediately they are differentiated, but usually remain dormant often for a considerable time. These 'eyes,' or resting buds, are constantly met with in sections near the apex of rhizome-branches, and it becomes a matter of considerable difficulty to tell whether the case is one of dichotomy or of monopodial lateral branching. It is certain however that if a dichotomy exists, it does not resemble the ordinary Hepatic type, for there can be no question of the replacement of one apex by two fresh ones; the original apical cell of the shoot is *not* obliterated, but continues to remain active, at any rate until its shoot issues from the soil. The view I have been led to adopt, after examining somewhat large number of preparations, is that the branching in the underground shoots is strictly lateral. The apical cells of the lateral axes are formed at a very early period in the history of the segment cut off from the apical cell; they may even appear as the result of its first division, which cuts off an inner from an outer cell, the latter forming the new apical cell of the lateral axis. These new apical cells never, so far as I have observed them, develop at once, although they sometimes form one or two divisions, after which the large cell, whose size and relative richness in protoplasm mark it out from the surrounding ones, either remains dormant for a time, or, as is still more frequently the case, it does not develop any further at all.

What has been said will show that it is impossible to draw any conclusions as to the presence or absence of adventitious branches (in the true sense of the word) on the rhizome, but structures of this character occur occasionally on the ventral surface of the 'frond,' and commonly beneath a mid-rib near the spot where it forks. I have never seen these adventitious shoots develop, but their rudiments may not seldom be seen, especially when examining sections in which archegonia occur (cf. Pl. VI, Figs. 11-13). As however in the majority of the species of this genus, ventral shoots occur, it is a matter of some interest to meet them also in this aberrant type, although they never, under normal circumstances, pass out of a rudimentary state.

Structure of the Rhizome.—Transverse sections of the rhizome show a clearly marked strand of fibrous cells, whose walls become much thickened and exhibit a number of shallow pits on their walls, at first view resembling ordinary striation. The individual cells are very much elongated, and possess long pointed ends; when examined in sections near the apex, it is seen that the thickening commences in the cells in the middle of the strand, and extends from thence to those at the periphery. Immediately surrounding the strand are the large parenchymatous cells which constitute the chief portion of the rhizome, and which are bounded externally by an epidermis-like layer, the cells of which frequently grow out to form root-hairs (Pl. VI, Fig. 8).

The apex of an underground shoot is occupied by a single cell of peculiar form. The actual determination of its shape is beset with numerous difficulties, but after comparing sections cut in different directions one is led to the conclusion that it resembles a prism whose cross-section is that of an isosceles triangle, whilst the outer (free) and inner surfaces are parallel and they lie in planes at right angles to the long axis of the prism. This is a most unusual form for an apical cell, but the evidence is conclusive as to its occurrence in this plant. Transverse sections of the apex in good preparations always show a *triangular* cell occupying the centre, with

obvious segmentation parallel to its three sides (Pl. VI, Fig. 9): on the other hand longitudinal sections, cut in the plane of the ground, exhibit an *oblong* structure, with apparently oblong segments given off parallel to the sides and inner face (Pl. VI, Figs. 3, 4, 6). Owing to the height of the cell, several consecutive sections can easily be cut through it, both longitudinally and transversely, and the results are consonant with the above formed conclusion as to its actual form.

Sometimes however longitudinal sections exhibit an apical cell of apparently *triangular* outline, similar to that found in the foliose forms (Pl. VI, Fig. 5). But this appearance is due to the section having really traversed the large apical cell in an oblique direction, when of course this appearance would be produced. It is easy to convince oneself of the validity of this explanation by examination of serial sections.

Apart from the appearance as shown by the transverse section, it might be argued that the cell really is a wedge-shaped one, as occurs, according to Leitgeb, in many frondose *Jungermanniae*, e. g. *Blasia pusilla*, and the triangular shape in longitudinal section would, of course, result in such a case if the plane of section happened to be parallel with one of the triangular sides of the wedge, whereas an oblong form would appear in sections passing through planes at right angles to this. But the fact that *transverse* sections of the apex exhibit the triangular form above described, disposes at once of this view, since a wedge-shaped cell with a triangular head could not exist as such, but would be converted into something like a tetrahedron.

The apex of *Pellia epiphylla*, according to Leitgeb's description, accords somewhat with that of *Pallavicinia decipiens*, but in the former the prism is rectangular, and not triangular, in transverse section. Abscission of segments parallel to the innermost face of the cell is however a feature common to both. The form of segmentation prevailing in our plant, as will be seen by comparing Leitgeb's figures¹, differs in important characters from that exhibited by *Pallavicinia*

¹ Leitgeb, Unters. ü. d. Lebermoose, Heft III.

(*Blyttia*) *Lyellii*¹, which was especially studied by him, and this fact affords another example of the marked dissimilarity of form and structure in the vegetative organs, which may exist in even allied species, and the difference described by Leitgeb² as existing between *P. epiphylla* and *P. calycina* in the structure of their respective apices is another case in point.

Division in the lateral segments takes place by means of walls which first cut the segment into an inner and an outer cell, and then, in the ordinary vegetative segments, a further wall parallel to the segment-wall appears. The further divisions were not studied. The apical cells of branches are formed very early, and are delimited by the first wall which appears in the segment, cutting off the inner from the outer cell. It is the latter which becomes the young apical cell of a lateral axis.

The segments given off parallel to the inner face of the apical cell divide chiefly by longitudinal walls, and contribute to form the central strand of the rhizome, which has been described above.

When the dormant resting apical cells of lateral branches are about to become active, walls appear in them which eventually cut out a cell similar to that which has already been described for the mother-apex. But the new walls are somewhat irregular at first (Pl. VI, Fig. 10), and it must be borne in mind that the cell in its dormant condition is in form a rectangular prism, which of course is a consequence of the way in which it originates in the young segment. The new walls then have to cut out a triangular prism, but it frequently happens that this is not effected until several divisions have occurred, and it is to this fact that the somewhat irregular appearance as regards the arrangement of the cells at the new apex is to be attributed.

One result of this lateness in the development of the branches is seen in a feature which was also noticed by Leitgeb in *Pallavicinia* (*Blyttia*) *Lyellii*, and doubtless the explanation is similar in both cases. The axile strand of

¹ Loc. cit.

² Loc. cit.

prosenchymatous pitted cells of the branch never unites with that of the mother-axis, but it ends blindly in the base of the branch. The cells in this region are short, and their cavities are large, and this is of course due to the fact that at this early period, the longitudinal divisions which cause the fibres to be so narrow in the more developed parts of the branch, were much less frequent at the time of its incipient growth, while at the same time transverse divisions were much more abundant than is subsequently the case.

A biological result of this lack of union is seen in the relative independence of the daughter-axes and the ease with which they become severed from the mother-shoot, to form fresh plants.

Structure of the Fronds.—The 'frond' is formed, as has already been stated, by rapidly succeeding dichotomy accompanied by a development of the thallus-wings, which are suppressed on the creeping portions of the plant. The apex exhibits the same structure as in the subterranean parts, though it is more difficult to determine its structure here. The actual apical cell lies at the base of a sinus, caused by the rapid growth of the segments, and it is bathed in the mucilage poured out by the hairs formed for this purpose. The fresh apices which arise as the result of the pseudo-dichotomy in the frond do not behave like those of the creeping stems; on the contrary they develop immediately, and thus the mid-ribs of the different axes are not here disconnected, but form a forked system corresponding to the branches of the frond. The edges of the lamina are toothed, and these teeth are situated as filiform hairs on a multicellular base, but they do not appear to be formed in a definite manner from the segments. The occurrence of adventitious ventral shoots on the fronds has already been alluded to. They *may* have been formed near the apex of the frond, but I do not think this is the case. Moreover, the structure of their apical cells is different from that of growing axes, since they always exhibit a *triangular* outline in longitudinal sections of the frond (Pl. VI, Fig. 12), and this probably

is due to their being in reality wedge-shaped cells. It may here be pointed out that there is not a great dissimilarity between a wedge and a prismatic cell such as has been described for the growing parts of this plant. A triangular prism may be conceived of as a broadened wedge, and the difference, so far as the plant is concerned, lies in the respective positions which such a wedge takes up in the apex. If its cleaving edge is directed inwards, and its head forms the free outer surface, then the ordinary 'wedge-shaped' apical cell arises. If on the other hand, the wedge is situated so that its cleaving edge is directed vertically, its head ventrally, whilst its triangular sides form respectively the inner and outer faces of the cell, then a form of cell arises such as is met with in the growing axes of *Pallavicinia decipiens*. Moreover it is further possible that there is a special significance attaching to the shape of the apical cells occurring on these rudimentary adventitious branches, as pointing to an earlier type from which the triangular prisms of the rest of the shoots have been derived.

Sexual Organs.—The plants, so far as my own specimens show, are strictly dioecious, and the antheridia and archegonia occur on the dorsal surface of the branches of the frond. The latter are grouped over a dichotomy, whilst the former occupy the length of one or more of the final branches of the frond, and are commonly situated on either side of the mid-rib in considerable numbers. A branch of the thallus bearing antheridia differs from the sterile ones inasmuch as it is considerably narrower, and in this feature it recalls the behaviour of the pinnae of many Ferns, where the sporophylls differ from the sterile leaves of the same plant (e. g. species of *Lomaria*).

The antheridia arise as papilliform projections of cells very near the apex, and do not offer any special features of interest in their development. They ultimately form globose stalked bodies each situated in a depression of the frond, and roofed over by a scale which arises behind them, but leaves an opening anteriorly (Pl. VII, Fig. 18), through which the spermatozoids escape.

The archegonia occur in groups upon cushion-like protuberances which arise immediately over a 'vein,' and almost always at a spot where it is just forking. Each archegonial group is enclosed in a circular involucre, whose free margin is toothed or fimbriated.

There are seldom more than two or three archegonial groups formed on a frond, though I have seen as many as seven, and they are nearly always confined to the early branchings of the frond, and never, so far as I have observed, extend to the younger dichotomies.

The archegonia in their development conform to the ordinary Hepatic type and need not therefore be dealt with here, though it may be mentioned that paraphyses, or mucilage-hairs, occur plentifully amongst them. Even before fertilization, a second sheath can be discerned within the involucre, but it is only after fertilization that this develops into the large colesula which is characteristic of the genus. This further growth however takes place immediately upon fertilization, and is the first visible sign of its having been effected. All division in it is limited to an annular zone situated at its base, just where it joins the frond, and nuclear division may be favourably studied in this region.

But besides the effect of promoting the development of the colesula, fertilization also brings about a vast increase in the number and in the growth of cells occupying the base of the archegonial group, as well as in the fertilized archegonium itself. In this way the massive calyptra is formed, within the colesula, and it carries up the numerous barren archegonia on its growing surface. As at the base of the colesula, so also in this tissue, dividing nuclei can be conveniently studied.

Nuclear Division in the Gametophyte.—The nuclei are very small, but they possess the great advantage of containing but few chromosomes, and in all cases when I have been able to estimate their numbers with certainty, there have always been present four chromosomes to each nucleus. Gentian violet and orange used as a double stain are very valuable in researches of this nature, the vividly stained

violet fibrils standing out sharply on a brownish-coloured ground.

No importance can be attached however to results in counting chromosomes, whenever they are as small as is the case in the present instance, unless they can be seen grouped at the equatorial plane, and can be regarded from the *poles* of the spindle. A study of a profile-view may be very misleading, since owing to the minuteness of the bodies, and their closely-aggregated condition, it is extremely difficult to avoid overlooking some of them, or to be in any way certain of the correctness of the results obtained. Seen from the poles, in favourable preparations, the chromosomes are seen to be in the form of loops, and to be four in number (Pl. VII, Fig. 37*a*). The loops split longitudinally in the usual manner, and often in the diaster-stage it becomes a matter of difficulty to feel sure of the actual numbers, even when seen from the pole of the spindle. As the two sets of loops are now moving along the direction of the line of vision, it is obvious that they will present their *ends* to the observer, and thus each individual loop will appear as two dots (Pl. VII, Fig. 37). As the number of the chromosomes has become doubled, previously to moving apart from the equatorial plane, it is clear that at this stage there will be sixteen dots visible at the equator, but that they will not be in the same focus. It only requires a little care however to understand the facts as they thus appear. Beyond the *number* of the chromosomes, the nuclei do not present any especially interesting features, but in the particular just mentioned they acquire considerable interest when compared with the nuclei of the sporophyte.

The Sporophyte.—After fertilization, the oospore grows into a somewhat pear-shaped body within the rapidly enlarging venter of its archegonium, and it soon divides transversely into three cells (Pl. VI, Fig. 15). The lowest cell undergoes very little further division, and contributes in but a slight manner to the structure of the embryo. The middle cell rapidly divides, both transversely and longitudinally, and forms both the stalk and the chief part of the anchor-like

base of the sporogonium. The lining superficial cells of the lower part of the stalk and swollen base are remarkable in the richness of their protoplasmic contents, and in the large size of their nuclei. They perhaps play a part, not merely passive, in the absorption of food from the gametophyte.

The capsular portion of the sporogonium originates from the uppermost cell, but as the sporogenous cells and the wall-layer are not differentiated until somewhat late, it is possible that the upper discs, into which the middle of the three first cells falls, may also contribute a share in the formation of this part of the sporogonium. Divisions in the young capsule are by no means regular, but eventually the tissue which ultimately goes to form the spores and elaters becomes clear, by the relative richness of its cells in protoplasm.

In the young stages of the development of the sporogonium, instances of nuclear division are not unfrequently to be met with, and a striking difference is at once seen between these nuclei and those, already described, of the gametophyte. In the first place the sporophytic nuclei are relatively large, but the important difference lies in the fact that there are not four, but *eight* chromosomes, also of a looped shape, to each nucleus. Even a profile-view awakens a suspicion that the chromosomes, when in the equatorial plane, are more numerous in these nuclei than in the above mentioned gametophytic nuclei, and a view from the pole of the spindle amply confirms this. Fig. 34 illustrates such a view, and it will be seen that there are, in all, thirty-two dots shown, sixteen lightly, and sixteen darkly, shaded.

As a matter of fact this nucleus is in the diaster-stage, and the daughter-chromosomes are beginning to move apart. Those marked more lightly represent the ends of the retiring loops of one nucleus, whilst those shaded darkly belong to the loops of the other one. The chromosomes form U-shaped bodies here, and hence there are in reality only eight chromosomes to each nucleus, the double number (sixteen) being due to the fact that the loops are seen in the direction of the

shanks, which thus appear as dots, and of course twice as many as the actual number of the loops.

This relation between the nuclei of the sporophyte and the gametophyte respectively, affords strong confirmation to Overton's¹ view of the meaning of the halving of the numbers of the chromosomes in the reproductive cells of Phanerogams, when contrasted with the corresponding vegetative nuclei.

As the development of the sporogonium proceeds, the wall becomes several layers of cells in thickness, and the contents of the capsule differentiate into elaters and spore-mother-cells. The former are characterized by their increasingly elongated form, whereas the spore-mother-cells finally become rounded bodies. The elaters are scattered through the sporogenous tissue, and do not form strands, such as occur in *Aneura* or in *Pellia*.

Development of the spores.—The spore-mother-cells are somewhat poor in their cytoplasm, which contains large quantities of oil. A nucleus of large size, enclosed in a special protoplasmic mass, occupies a central position in the cell. Presently the characteristic change which precedes the actual formation of spores in the Hepaticae becomes apparent. The mother-cell becomes tetrahedrally lobed, and the cell-walls, at their inner angles, grow into the cell-cavity towards the nucleus.

This body is surrounded, as has been said, by somewhat dense protoplasm which probably represents an archoplasm, and just before nuclear division it behaves in a most striking manner. Simultaneously it projects out into each lobe of the mother-cell, and thus forms a four-rayed star, with the nucleus occupying the centre. These arms, or spindles together form what I have recently, in a paper communicated to the Royal Society², termed a 'quadripolar spindle.' The spindle-arms or rays stain deeply with nuclear stains, and I am strongly of opinion that the substance from the *nucleolus*

¹ E. Overton, On the reduction of the Chromosomes in the Nuclei of Plants, *Annals of Botany*, vol. VII.

² *Proc. Roy. Soc.*, vol. LIV, No. 330, 1894.

becomes diffused out into them. The objects are too small to admit of this suggestion being verified by observation, but I have already shown¹, and my results have been confirmed and extended by Zimmermann², that the nucleolus may break up and pass out into the cytoplasm during division in pollen-mother-cells, so there is no inherent improbability in the view that a similar state of things obtains here. The chromatic elements of the nucleus next begin to become individualized, and, so far as I have been able to determine, on studying many hundreds of preparations, they do so in an irregular manner. It is possible that the spirem-stage may occur, as it certainly does in the vegetative cells of the sporophyte, but I have never seen it. Four chromatic droplets form the first positive evidence of approaching division which I have seen *in the nucleus*. The quadripolar spindle however has been formed *before* these four chromosomes appear. In several instances I have seen the centre of the nucleus occupied by a large mass of chromatic substance, which was distinctly four-lobed, as though the four bodies, already referred to, arose by the breaking up of an original lump in this way. Fig. 30 is a camera-lucida-drawing of such a case, and I may say that I possess photographs of this actual specimen which exhibit exactly the same appearance as that shown in the drawing.

Shortly after the appearance of the four chromosomes, their arrangement becomes more regular, and their number is increased by division to eight. Their shape is not that of *loops*, as is the case with the vegetative nuclei, but that of *rods*, and it may be remarked that this difference of form is extremely common between the vegetative and reproductive nuclei of the same plant, e.g. in *Lilium Martagon*. The eight chromosomes now point off in pairs as though about to travel to the lobes of the spore-mother-cell, and I believed until recently that only two chromosomes went to each lobe. I have however since found two instances, so clear as to admit of no

¹ J. B. Farmer, On nuclear division in the Pollen-mother-cells of *Lilium Martagon*, Annals of Botany, vol. VII.

² Zimmermann, Beitr. z. Morph. u. Physiol. d. Pflanzenzelle, Bd. II. Heft 1.

doubt whatever, in which, just before the exit from the centre, a further doubling of the chromosomes occurs, so that in reality four of these bodies, and not two, as I previously supposed, go to form the nucleus of each spore. The whole process is very much crowded up, the four-rayed spindle persisting to the end; and even after the exodus of the chromosomes, traces of it can still be seen, converging to the original centre. Moreover, so late is the whole process, that the sculpturing which characterizes the walls of the mature-spore is well advanced before nuclear division is even incepted. In many cells, the retreating chromosomes could be traced, but there nearly always appeared to be only two: this however arises from the fact that I only saw them in profile, and not from the pole, and hence the two nearer ones covered the pair lying below. However in the young nuclei of the spores, whose walls have not as yet been formed to meet in the centre, the four chromosomes could sometimes, though with difficulty, be made out. The archoplasm, which forms the spindle, breaks asunder, and each arm contracts up along with the young nuclei, thus forming a sort of sheath around them, like that which surrounded the original mother-nucleus.

The spores become finally separate by the appearance of membranes which complete the isolation of their contents in the centre of the tetrad, an isolation which is however nearly effected by the earlier ingrowths mentioned above.

Now these features attendant on spore-formation in *Pallavicinia* are so abnormal, or at least depart so widely from our ordinary notions of cell-division, that I deferred publishing them until I should have had the opportunity of comparing it to the process as it obtains in other Hepaticae. I have examined the process in a number of other genera, and am still engaged in extending the observations, which will be brought together in a future communication. In this place I will only refer briefly to these results in so far as they help to elucidate the case of *Pallavicinia*. If the process be followed out in *Aneura multifida*, a form which is abundant in this country, essentially the same phenomena are perceptible,

but the process is, as it were, slowed down. At a certain time, and previously to any visible change in the nucleus, a quadripolar spindle is formed, each of the four rays projecting into one of the four lobes of the spore-mother-cell. At the extremity of the rays may be seen a distinct protoplasmic aggregation which represents a centrosphere, but a centrosome was not observed.

Now this quadripolar spindle is *transient* in *Aneura*, for when the chromosomes become delimited and individualized, the original quadripolar spindle breaks up and is replaced by two *independent* spindles in the equators of each of which there may be counted eleven, or possibly twelve chromosomes. These are excessively small, and it is difficult, and perhaps not very important, to determine their absolute number. Then each spindle acts independently, and the chromosomes (which here too are in the form of short rods) become doubled, and nuclear division takes place as on normal lines.

Aneura pinguis, of which I obtained a good supply when in Ceylon, also exhibits very plainly the quadripolar spindle, which diverges from the relatively large nucleus. In this plant it can be clearly seen that the nucleus, at the stages at which my specimens were fixed, is passing into the spirem-stage. I could not, owing to failure of material, obtain the stages of actual nuclear division in this plant, but the presence of such a spindle is of interest.

In *Lophocolea* I have also seen nuclear division proceeding on the same lines as I have just described for *Aneura multifida*, but I have not seen as yet anything like a quadripolar spindle. It may however ultimately prove to be there, and my failure to recognize it hitherto may be due to its transient existence—a difficulty which I had experienced already in *Aneura*—in *Lophocolea*, as in *Aneura*, and there was the same difference between the reproductive and vegetative nuclei, as regards the shapes of the chromosomes; i. e. in the former they appear as rods, in the latter as loops.

It is obvious, from what has been said, that a serial

modification from normal division to a possibly reduced, and at any rate much compressed, type of karyokinesis is traceable within the genera comprised in the Hepaticae. There are, furthermore points of theoretical interest connected with the modifications just described, but it is best to postpone the discussion of these questions until further investigations have rendered it possible to review and compare the process in a wider range of species than can be done at the present time.

EXPLANATION OF FIGURES IN PLATES VI AND VII.

Illustrating Mr. Farmer's paper on *Pallavicinia decipiens*, Mitten.

Figs. 1-37 refer to *Pallavicinia decipiens*; Figs. 38 and 39 to *Aneura multifida*.

Fig. 1. Barren plant (natural size).

Fig. 2. Fertile frond (enlarged). *a*, involucre, surrounding a group of archegonia: *b*, colesula, from which a sporogonium is issuing.

Fig. 3. Longitudinal section of apex of rhizome, in horizontal plane, with apical cell: *c*, the apical cell of lateral branch.

Fig. 4. Longitudinal section of rhizome-branch which is growing up to form a frond.

Fig. 5. Longitudinal section of rhizome, somewhat oblique, with an apparently triangular apical cell: *c*, apical cell of branch.

Fig. 6. Young branch of rhizome.

Fig. 7. An older ditto, showing the central strand.

Fig. 8. Transverse section of rhizome.

Fig. 9. Transverse section of apex of rhizome: *a*, apical cell: *c*, apical cell of branch (cut below the surface).

Figs. 10, 10 *a*. Young apices of lateral branches forming their apical cells.

Figs. 11, 12, 13. Adventitious buds, from ventral surface of frond beneath archegonial groups.

Fig. 14. Apex of a branch of the frond.

Fig. 15. Group of archegonia, one has been fertilized and contains a three-celled embryo: *i*, involucre: *p*, colesula.

Figs. 16, 17. Young embryos.

Fig. 18. Longitudinal section of antheridial branch (somewhat diagrammatic).

Fig. 19. Spore-mother-cell, with ingrowing cell-walls.

Figs. 20-29. Stages in development of spores.

52 Farmer.—On *Pallavicinia decipiens*, Mitten.

Fig. 30. Spore-mother-cell with four-lobed chromatic mass.

Fig. 31. Nucleus from young sporogonium preparing for division.

Figs. 32, 33. Different views of dividing nuclei from young sporogonium.

Fig. 34. Dividing nucleus in diaster-stage, from the same tissue. Only the *ends* of the looped chromosomes are shown.

Fig. 35. Dividing nucleus in coesula.

Figs. 36, 37, 37 a. Dividing nuclei of cells at the base of the calyptra.

Fig. 38. Spore-mother-cell of *Aneura multifida*, with quadripolar spindle.

Fig. 39. Karyokinesis in spore-mother-cell of the same.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 7.

Fig. 8.

Fig. 9.

Fig. 10.

Fig. 10^a

Fig. 11.

Fig. 12.

J.B. Farmer del.

FARMER.—PALLAVICINIA DECIPIENS, Mitt. & ANEURA MULTIFIDA, Dumort.

Fig. 4.

Fig. 5.

Fig. 6.

Fig. 14.

Fig. 13.

Fig. 15.

Fig. 16.

Fig. 17.

Fig. 18.

Fig. 21.

Fig. 22.

Fig. 23.

Fig. 26.

Fig. 28.

Fig. 30.

Fig. 27.

Fig. 29.

Fig. 31.

Fig.

J. B. Farmer del.

FARMER.—PALLAVICINIA DECIPIENS, Mitt. & ANEURA MULTIFIDA, Du

Fig. 19.

Fig. 20.

Fig. 24.

Fig. 25.

Fig. 32.

Fig. 34.

Fig. 35.

Fig. 36.

Fig. 37.

Fig. 37^a.

Fig. 33.

Fig. 38.

Fig. 39.