aged twenty years, who was hit or kicked in the front of his neck in a football game. When examined it was apparent that the thyroid cartilage had been sprung, as it were, from its seat on the cricoid, and at the same time had suffered a rotation in a vertical direction. The inferior border of the thyroid cartilage was about one-sixth of an inch below its resting place on the cricoid.

The cartilage was replaced by careful digital manipulation, and was maintained in position with adhesive strips placed horizontally. At the end of five weeks the patient had practically recovered.)

Foreign Bodies in the Lungs.—DR. E. FLETCHER INGALS, of Chicago (*Journal of the American Medical Association*, May 27, 1905), reports a case in which he successfully removed a collar-button from the lungs under bronchoscopy.

Tracheotomy Wounds.—DR. MILES F. PORTER, of Fort Wayne, Ind. (*Journal of the American Medical Association*, April 8, 1905), advocates the immediate closure of the trachea with suture as lessening the dangers from pneumonia and infection, shortening the convalescence and reducing the deformity of the scar to the minimum.

PATHOLOGY AND BACTERIOLOGY.

UNDER THE CHARGE OF

WARFIELD T. LONGCOPE, M.D.,

DIRECTOR OF AYER CLINICAL LABORATORY, PENNSYLVANIA HOSPITAL,

ASSISTED BY

G. CANBY ROBINSON, M.D.,

RESIDENT PATHOLOGIST, PENNSYLVANIA HOSPITAL, PHILADELPHIA.

Experiments in Congenital Transmission of Tuberculosis and Congenital Tuberculosis.—The transmission of tuberculosis from the mother through the placenta to the child is known to occur, and FRIEDMANN (*Virch. Arch.*, 1905, Bd. cxxxi. p. 150) has collected 22 cases from the literature in which a placental infection in human beings was proven through anatomical study, while including the observations upon animals upwards of 100 such instances have been recorded. On the other hand, direct evidence to show that tuberculosis can be acquired by the child from the father is almost lacking. Only two cases which suggest this method of transmission have been reported—one by Sarwey, the other by Landouzy. Klebs, however, has stated that "tuberculosis of the father is ten times more dangerous for the child than tuberculosis of the mother."

Bearing upon this question, Friedmann's experiments are of much interest. A series of inoculations of human tubercle bacilli and the