retains its place by a convention which remains valid owing to the force of long-continued custom. If the fluctuations in prices as measured by it became too great, it would have to be discarded as a standard of value.

T. K. R.

Artificial Waterways and Commercial Development (with a History of the Erie Canal). By Dr. A. Barton Hepburn. Pp. ix+115. (New York: The Macmillan Co.; London: Macmillan and Co., Ltd., 1909.) Price 4s. net.

AFTER a long period of effacement, artificial waterways are beginning to regain some amount of public interest and concern. The advent and rapid development of railways during the last century was responsible for their relegation into a background of indifference and neglect, and so long as men's minds were dominated by schemes of rapid locomotion at any cost, it was difficult, and, in fact, impossible, for canals to maintain any footing in competition with a system of transit infinitely more expeditious and direct. But a change is taking place in public feeling. It is being recognised that canals have been at an undue disadvantage, and that, as a means of locomotion, they possess features which merit encouragement and development. Inland water carriage for goods, though slow, is safe and cheap, and canals possess a striking advantage over railways in that, in place of isolated depôts at long intervals, they possess a continuous frontage workable throughout their entire length. On these and other grounds, public interest in canals has been aroused, and a Royal Commission in this country has lately had under consideration the means best adapted for their revival and amelioration.

Dr. Hepburn's book is a timely contribution to the evidence on the subject. Written from an American standpoint, it constitutes an appeal to the citizens of the United States in regard to the development of their artificial waterways. It recites in brief compass the principal historical facts connected with canals throughout the world, and then proceeds to consider in more extended detail the canal system of New York, describing its inception, development, and present condition. Thence the author passes, by a transition natural to a patriotic American, to an account of the Panama Canal, with its vicissitudes and possibilities. The volume closes with fifteen statistical appendices.

Hydrographical Surveying. A Description of Means and Methods employed in constructing Marine Charts. By the late Rear-Admiral Sir William J. L. Wharton, K.C.B. A new edition, revised and enlarged by Rear-Admiral Mostyn Field, F.R.S. Pp. viii+475. (London: John Murray, 1909.) Price 21s. net.

THE late Admiral Wharton's "Hydrographical Surveying," which has been for so many years a standard work and one of the best books for surveyors that has ever been published, has now been brought up to date by his successor, Rear-Admiral Mostyn Field, the present hydrographer to the Admiralty. Admiral Field has endeavoured to alter the text of the former work as little as possible, but at the same time to enlarge it considerably by the addition of new features, including expedients connected with work in the field which have been found useful in practice, in order especially to assist the young surveyor by directing his attention to useful methods of procedure which otherwise he would only pick up as his experience ripened. In addition to these features, all the latest improvements are fully described, such as the use of photography for the reproduction of charts, auto-

matic tide gauges, improved instruments for observing currents and taking deep-sea soundings, and, finally, the usefulness of the Barr and Stroud range-finder for surveying purposes.

The volume as it now appears, brought thoroughly up to date and accompanied by excellent diagrams, cannot fail to be of the utmost value to all surveyors.

H. C. L.

Œuvres complètes de Christian Huyghens publiées par la Société hollandaise des Sciences. Vol. xi., Travaux mathématiques, 1645-1651. Pp. iv+369. (La Haye: Martinus Nijhoff, 1908.)

THIS volume is divided into several parts. The first part deals with Huyghens's early writings (1645-6), and is preceded by an account of a manuscript by van Schooten which formed the basis of Huyghens's first mathematical studies. The writings in question deal, *inter alia*, with elementary geometrical considerations relating to the parabola and funicular polygons. The next portion consists of Huyghens's three books entitled "De iis quæ liquido supernantant" (1650), forming a collection of applications of the principle of Archimedes to floating bodies of simple shapes. A number of geometrical problems dated 1650 follow, and the volume concludes with the "Theoremata de quadratura hyperboles, ellipsis, et circuli ex dato portionum gravitatis centro" (1651). The volume is well got up, and forms an interesting contribution to the history of mathematics.

The General Characters of the Proteins. By Dr. S. B. Schryver. Pp. x+86. (London: Longmans, Green and Co., 1909.) Price 2s. 6d. net.

THIS is another of the series of monographs on biochemistry which are being issued by Messrs. Longmans under the editorship of Drs. Hopkins and Aders Plimmer. The previous monographs have been already noticed in these columns, and two of these dealt with the proteins from the more strictly chemical point of view. Dr. Schryver now adds another chapter to, and by no means exhausts, this large subject. The first section deals with the physical properties of the proteins (solubilities, crystallisation, heat coagulation, rotatory power, electrical conductivity, and so forth); the second with their general chemical characters (tests, distribution of nitrogen, compounds with acids, bases, halogens, &c.); and the third with the precipitin reaction, which is commonly known as the biological test.

The whole is treated in a technical but clear manner; references are given to the authorities quoted, and the booklet will prove a useful addition to the library of the physiologist, and should be found in every laboratory devoted to biochemical research.

W. D. H.

LETTERS TO THE EDITOR.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

The Gravitative Pull upon the Moon.

THE error made by Mr. McLennan in NATURE of May 6, p. 276, is a curious one, which may perhaps be made more often than we are aware of, and therefore is worth correcting.

It is true that gravitational pull and centrifugal force both decrease as square of distance increases, each with its own cause of decrease, so as to remain equal and opposite; but then the two causes of decrease are not to be piled on to one of those forces! That is the error.

Mr. McLennan has done his arithmetic correctly, but the calculation is really extremely simple, as thus:—

The moon's mass is $1/80$ th that of the earth, which is 6×10^{21} tons. At the moon's distance, which is 60 earth's radii, terrestrial gravity is reduced to $1/3600$ th of its value at the earth's surface. Consequently, the weight of the moon, *i.e.* the earth's gravitational pull on it, is equal to the ordinary commercial weight of

$$1/3600 \times 1/80 \times 6 \times 10^{21} \text{ tons.}$$

An alternative, but not a supplementary, way of doing the sum is to say that the moon revolves through the angle 2π in, say, $27\frac{3}{4}$ days, and that therefore the centripetal force necessary to hold it in is

$$m\omega^2 r = \frac{6 \times 10^{21}}{80} \text{ tons} \times \left(\frac{2\pi}{27\frac{3}{4} \text{ days}} \right)^2 \times 60 \times 4000 \text{ miles.}$$

The acceleration part of this is about 13,000 miles per day per day, which is the same as $32/3600$ feet per second per second; and this is obviously in accordance with the law of inverse square.

OLIVER LODGE.

A Direct Estimate of the Minimum Age of Thorianite.

I WISH to record an experiment lately made which affords more direct proof of the great antiquity of radioactive minerals than anything previously attempted.

The principle of the method is to determine (1) the total accumulated helium in the material; (2) the rate of formation in the same material at present.

A sample of thorianite was found to contain 9 c.c. helium per gram.

Four hundred grams of this thorianite was got into solution, and all traces of helium boiled out with scrupulous care. After it had been allowed to stand for seven weeks, the solution was boiled out again. A little nitrogen and nitric oxide were evolved. These were removed by charcoal cooled in liquid air, and the residual helium collected in the capillary of a McLeod gauge. D_2 could be seen in the spectrum, but the volume of the gas was too small to measure. It was certainly less than 2×10^{-6} c.c., perhaps much less.

Thus the annual rate of production of helium per gram is less than 3.7×10^{-8} c.c. The 9 c.c. initially present cannot, therefore, have accumulated in a less time than 240 million years. Experiments on a larger scale, which are in progress, will probably lead to an extension of this estimate.

Prof. Joly, in his interesting book, "Radio-activity and Geology," has brought various objections against the radio-active method of measuring geological age. These will require to be carefully weighed. I may remark, however, that in the present case no uncertainty arises on the ground that the substance may formerly have contained much more uranium or thorium than at present, for thorianite consists almost entirely of the oxides of these elements.

R. J. STRUTT.

Imperial College of Science, May 11.

Sense of Smell in Flies.

By far the most efficient of fly-destroyers with which I am acquainted is a dilute solution of formaldehyde. If two teaspoonfuls of formalin (40 per cent. formaldehyde) be added to a soup-plate filled with water, flies go to it, one after the other, to drink, especially in the early afternoon. Some die in the water; many fall in the immediate neighbourhood of the plate; others succumb on window-sill or floor. As the result of leaving a single plateful of the solution on the kitchen table (I am writing in the south of France) hundreds of dead flies are each day swept up from the floor. Formalin water is free from the gruesome associations of fly-papers and other traps which hold their struggling victims. It may even be turned to ornamental uses. A wire cage placed in the centre of the dish may be crowned with flowers, which flourish equally as well, with some slight but interesting changes in tint, in dilute formalin as in pure water. The solution neither attracts nor repels flies. Two similar dishes placed side by side, the one containing pure water and the other formalin, are visited, so far as one can judge, with equal frequency. It is somewhat strange that so

small a dose proves fatal when taken into the fly's alimentary canal. I find that, to free a room from flies by vaporising formalin, the air must be rendered quite irrespirable by a human being. The room needs to be amply ventilated before one ventures into it.

The interest which attaches to this observation, that flies will drink a solution of formaldehyde, lies in the proof which it affords that the mechanism of their sense of smell is similar to our own. No volatile body the density of which is not greater than that of air is a stimulant of our olfactory membrane. Formaldehyde, H_2CO , has a density of 15 only. Playing in paradoxes, one might say that it undoubtedly has a malignant odour, but we cannot smell it. If the nose be placed close to a vessel containing a dilute solution of formalin a scent is recognisable, but this I take to be due (a chemist will correct me) to impurities present in the commercial product. Yet I find that when I sit within a yard of it my eyes begin to smart. In this respect, however, I am, I know, exceptionally sensitive. I cannot dissect specimens preserved in formalin until after they have been long soaked in water and spirit frequently changed. Once, when conducting a *viva voce* examination with the aid of formalin preparations, I developed so acute and painful, although happily transient, an attack of conjunctivitis as made it impossible for me to attend the examinees' meeting. The fact that so deleterious a volatile body as formaldehyde does not appeal to our sense of smell would seem to confirm the only theory of the physics of olfaction at present plausible, though far from comprehensible, namely, that which attributes to the hairs of the cells of the olfactory membrane the capacity of responding to the alterations in the vibration frequency or amplitude of molecules of air which are caused by the presence amongst them of heavier molecules.

ALEX. HILL.

Mentone.

The Production of Radium from Uranium.

EXPERIMENTS on which I have been engaged for the past six years have until now failed to establish the production of radium from uranium. With carefully purified uranium salts, using considerable quantities, the growth is too small to be detected for the first two or three years, and is less than $1/10,000$ th of what would occur if a direct change of uranium into radium took place. With commercial uranyl nitrate, on the other hand, purified from radium by precipitating barium sulphate in the solution, a distinct though small production of radium was observed in 1905, and subsequently confirmed (*Phil. Mag.*, October, 1908, 632). This is explained by the existence of an intermediate parent of radium in the series with a very long period of life, which has been found by Boltwood and by Rutherford in preparations of actinium, and recently isolated by the former from uranium minerals, and called "ionium."

I have now been able to establish the production of radium in all the solutions of very carefully purified uranyl nitrate prepared by Mr. T. D. Mackenzie and myself (*Phil. Mag.*, August, 1907, 272). Three separate solutions containing 255, 408, and 278 grams of uranium (element), and of age since purification at the present time respectively 3.53, 2.40, and 2.73 years, have been kept under observation. The method of testing has been much improved since formerly, and the error of a single determination as now carried out probably does not exceed 10^{-12} gram of radium. The quantity of radium in the oldest solution is now about 4×10^{-11} gram, which is nearly twice as great as initially. During the past year five measurements of the quantity of radium in this solution have been made, and they show that, within the stated limit of error of observation, the production of radium has proceeded during that period *proportionally to the square of the time*. The tests on the other solution show that in them also the production of radium is now proceeding according to the same law and at a similar rate.

That the initial rate of production of radium from uranium should vary according to the square of the time was deduced mathematically by Rutherford (*Jahr. Rad.*, 1908, v., 164) on the assumption that there was only one intermediate substance of period of life long compared with