greater number of cases the prismatic effect should correct about one-half the manifest error, but in some cases a full correction, especially of a small vertical error, may be given.

Prisms and decentered lenses interfere with some of the visual judgments and therefore are objectionable. For this reason it is better to cure muscle errors of low degree by proper prismatic exercise. Muscle errors of high degree should be treated surgically.

THE GENESIS AND TREATMENT OF THE MYOPIC EYE.*

S. D. RISLEY, M.D.

ATTENDING SURGEON WILLS EYE HOSPITAL.
PHILADELPHIA.

There is no more interesting or important chapter in ophthalmology than that which treats of the etiology, nature and treatment of the myopic eye. The widely differing views which have been expressed regarding each of these points seem to justify a renewed presentation of the subject.

HISTORY.

The observations made in the early part of the century by James Ware regarding the relatively greater prevalence of myopia among the educated portions of the community were verified by the later examinations in the schools by many observers, who demonstrated its steadily increasing percentage as the children advanced in age and school progress. Since this was associated with a nearly equal fall in the percentage of hypermetropic eyes it required but a step to the seemingly logical conclusion that the small hypermetropic eyeball of childhood increased in size with the physiologic growth of the child into emmetropia and from this into the large myopic eyeball; in a word, that the myopic eye was a physiologic evolution. Closely allied to this was the proposition that, since only those eyes which are subjected to the demands of civilized life were observed to undergo this change of refraction, therefore it was the eye of civilization, a species of social evolution, an adaptation to the act of work.

More careful observations of the conditions, however, revealed the fact that these conclusions were untenable, since they must be maintained in face of well-known and characteristic pathologic conditions associated with myopia, which imperiled the integrity of the eye, e. g., posterior staphylomata, choroidal atrophies at the temporal border of the optic nerve and in the region of the fovea, a fluid, i. e., a degenerate vitreous with floating opacities and frequent detachment of the retina, all of which sustained with myopia an intimate and undoubted relation of cause and effect. Increasing experience and painstaking observation have served, after nearly half a century, to maintain and enforce the truth of the early teaching of Donders, to the effect that the myopic eye is a diseased eye.

Accepting the proposition as stated, it is obvious, therefore, that myopia can not be regarded as either a physiologic or sociologic evolution, but is to be viewed as one of the many baneful results of civilized life over the physical well-being of the race. The problem presented for solution is its prevention.

GENESIS OF THE MYOPIC EYE.

The observations of Cohn and others showing the rapidly increasing percentage of the disease during the

educational process led thoughtful observers to inquire into its cause. Since it was evidently in some way closely related to the work in the schools it was ascribed to the bad hygienic environments of the school room, and an era of improvement was inaugurated in the lighting of the school houses, the arrangement and construction of desks and seats, the paper and type used in school books, etc. A few years later a re-examination of the same children under the improved conditions of work, much to the disappointment of these industrious and careful observers, discovered no notable diminution in either the percentage of increase or in the arrest of the progress of the myopia in those already myopic at the first examination. It was obvious, therefore, that some hitherto unnoticed factor was responsible for the genesis of the myopic eye.

The earliest suggestion with which I am familiar of the possible relation between astigmatism and myopia was made by Dr. Green¹ of St. Louis in 1867, and again² in a paper in 1871. He closes the latter paper with the statement that so far as he knew statistics of large numbers of eyes to establish the frequency of astigmatism were wanting. My own statistics collected in the Philadelphia schools, 1878 to 1881, were, so far as I know, the first to establish the truth of Dr. Green's suggestion, of which I was at the time entirely ignorant. In 1880 I published³ a group of cases which, while under observation, had passed from hypermetropic into myopic astigmatism without at any time becoming emmetropic. That is to say, they passed through the turnstile of astigmatism from hypermetropic to myopic refraction. They were all asthenopic, and the increase of refraction was associated with disease of the intra-ocular tunics. This publication was followed in 1881 by the report of the school statistics,⁴ which revealed a group of interesting and important facts bearing on the genesis of the myopic eye.

First. The emmetropic eye not only remained in a nearly uniform percentage through all ages of school life, but enjoyed the highest acuity of vision and was relatively free from pain and disease.

Second. The eyes with a simple hypermetropia stood next to emmetropia in all these respects.

Third. The eyes with hypermetropic refraction, especially where astigmatism was present, manifested symptoms of asthenopia, lowered acuity of vision and pathologic states of the fundus; notably irritative and absorption changes throughout the fundus, granular changes at the fovea and crescents of choroiditis at the temporal margin of the optic disc in all stages of development from a faintly outlined crescent to far advanced atrophies which had before been regarded as physiologic anomalies when seen in hypermetropic eyes, but as pathologic if associated with myopic refraction.

Fourth. It was shown, as had been demonstrated by all observers, that not only did the myopic eye increase in percentage during school life, but that there was a closely corresponding diminution in the percentage of hypermetropia. This group of facts seemed to afford a sufficient demonstration for the conclusion that the myopic eye, with its characteristic pathologic changes in the fundus, was but a later stage of the conditions witnessed in the eyes with hypermetropic astigmatism, hence that the former was recruited from the latter.

Fifth. In the very large percentage of congenitally

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee: Drs. Frank Allport, H. V. Würdemann and J. A. Lippincott.

1. American Journal of the Med. Sciences, January, 1867.
2. Presented to the Amer. Oph. Society.
3. Amer. Jour. of the Med. Sciences.
4. Trans. Med. Soc. State of Pa., 1881. A System of Diseases of the Eye, Norris and Oliver, vol. ii, 1894, Art. "School Hygiene."

defective, that is to say, eyes with hypermetropic astigmatism which were shown to exist in the schools, we have, therefore, a sufficient explanation for the steadily increasing percentage of myopia notwithstanding the improved environments of the school room.

This was essentially a new view of the subject of myopia. The question of heredity resolved itself into a hereditary transmission of the anatomical defect on which the underlying pathologic conditions resulting from eye strain depended and which, in turn, led to the impaired nutrition of the ball and the distention of its coats. The malformation of the eyeball, that is to say, astigmatism, and the faulty binocular balance so frequently observed in astigmatic eyes, doubtless sustain a very important relation to certain distortions of the skull and abnormalities in the shape and size of the bony orbit. We have, therefore, primarily a congenital anatomical anomaly as the starting point or basis for the chain of events we have been considering. The truth of these propositions finds marked confirmation in the daily routine of office work, where we much more frequently find the children of myopic parents with hypermetropic astigmatism than with myopia, which, if carefully corrected, removes the asthenopia and associated conditions which are the starting point for the myopic eye. Indeed, it is, in my experience, extremely rare to find a myopic eye before the school age.

TREATMENT OF MYOPIA.

The treatment of myopia should therefore be conducted with this chain of events in mind. In a word, it should be begun before myopia is established. I have already shown⁵ that in 200,000 eyes corrected by glasses in Philadelphia in twenty years, that the number of eyes with myopic refraction applying for relief fell from 28.43 per cent. in the first period to 16 per cent. in the last, a diminution of approximately 50 per cent. It is apparent, therefore, that our highest duty is in preventing the disease. It was to this end that in 1881 I first urged the importance of examining the eyes of every child at the beginning of the educational process in order to determine if the eyes were fitted to safely undertake the strain of school life. Unfortunately, the myope is with us and requires at our hands the highest skill and solicitous care to ward off its baneful results.

It is obvious from the foregoing conclusions that treatment resolves itself into two factors: First, attention to the pathologic conditions; second, the careful correction of the anomaly of refraction and accommodation and any of the associated abnormalities of ocular balance.

I wish at the outset to urge that no final attempt should be made to prescribe correcting glasses until the fundus conditions upon which the progressive near sight depends shall be brought under control. To this end the following routine treatment has in my own hands been of value. Complete rest from all near work, the steady and persistent use of a mydriatic, protection from light by smoked glasses, the temporary abandonment of the accustomed correcting glasses, the internal administration of the so-called alterative drugs, e. g., potassium iodid and bromid with the bichlorid of mercury, iodid of iron, etc., and the careful attention to the general regimen. This method will usually lead to a more or less speedy relief from the asthenopia and to marked improvement in the condition of the uveal tract. Then, and not before, a careful and painstaking study should be made of the refraction of each eye separately. It is by no means

an easy task to determine accurately the exact static refraction at the macula in high myopia. Therefore, not only patient subjective study with glasses and test cards, etc., should be employed, but the results so obtained should be checked and verified by the objective methods we now have at command.

It is very easy, indeed, to overlook the lower grades of astigmatism, especially if the corneal meridians are heteronomous, a very common condition in myopic eyes. The static refraction of each eye having been determined, we can now for the first time, with any degree of accuracy, study the state of the binocular balance. This should be done with carefully centered correcting glasses, placed in some fixed form of apparatus, the eyes being maintained in the primary position. When studied in this way we soon learn to revise many of the views formerly entertained regarding the mobility of the myopic eye. The great frequency with which a dark ruby glass, or the multiple Maddox rod, or some other suitable device, will reveal a faulty binocular balance, was to me at first a great surprise, but not more so than the promptness with which the asthenopic and irritation symptoms disappeared when this was corrected, either by a glass or by tenotomy in suitable cases. This, however, is to be expected when we consider the essential anatomical basis or origin of both the astigmatism and the faulty attachment of the muscles of the globe.

In the higher grades of astigmatism not only are the corneal meridians likely to be heteronomously placed, but it is comparatively rare not to find an absolute heterophoria due to the faulty attachment of one or more of the ocular muscles.⁶ It is in this fact and to the associated anisometropia rather than in the large ball that we find adequate explanation for the recognized difficulty of binocular vision in many cases of high myopia. It is important in the determination of the ocular balance to avoid deception growing out of the presence of eccentric fixation in one or both eyes caused by small, partial or complete, macular scotomata, so common in high myopia.

I have seen many cases of progressive nearsight in which the tendency to progression, the asthenopia and the choroidal disease were rebellious to all treatment and to the most careful corrections of the anomaly of refraction until the binocular balance was corrected, either by suitable adjustment of prisms or by tenotomy or advancement of the faulty muscles. Many illustrations and instructive examples would be cited if time permitted.

Should full correcting glasses be worn in high myopia? I invariably prescribe full correcting glasses for myopia of 12 D. or less, but if the myopia is higher than 4 or 5 D., under correcting glasses are given for near work, the allotted far point chosen depending upon several factors, e. g., the age of the patient, the power of convergence and the nature and the amount of work required; the condition of the fundus and the tendency to progression of the disease. When the tendency to progression is marked the full correcting glasses are allowed only out of doors, the undercorrecting glass to be worn constantly in the house until the progression is arrested. In many cases bifocal glasses are ordered. If the convergence is weak or painful, a prism, base in, is secured by decentering the glasses or by having the near work glasses ground on a prism with base in, or a somewhat stronger glass is given. If esophoria is present at 6 m. a weaker glass is prescribed, so that all near

5. Archives of Ophthalmology, July, 1894.

6. Abnormalities of Oc. Bal., University Med. Mag., January, 1895. Trans. International Oph. Congress, Edinburgh, 1894.

work is done at the remote point of the glass, and thus all strain taken from the accommodation, which relieves the tendency to undue convergence. In myopia of 15 D. or higher a full correcting glass is rarely desirable or safe and usually is not well born at any age, certainly not when the usually critical fundus changes are present. For such patients I usually prescribe a glass for constant wear, deducting from 3 D. to 5 D., and allow a lorgnette of 3 D. or 5 D. to be placed in front of the under-correcting glasses temporarily when the full correction of the myopia is necessary or desirable to secure accurate distance vision.

RESULTS OF TREATMENT.

What have been the results of these methods of treatment? I have a long list of myopic patients who have been under more or less continuous observation for many years, some for twenty years or more. I have witnessed during these years many instances of separation of the retina, but they have occurred, with but three exceptions, in patients who came for the first time because of this unfortunate accident.

In these exceptions one, an eminent physician with astigmatism = 3 D. in each eye and an additional myopia of 20 D. in the right eye and 18 D. in the left, was knocked down by a bicycle and rose with detachment of the retina in the right eye, which was rapidly followed by opacity of the lens. The left eye became irritable after this accident and slowly developed a small macular atrophy. He had been under my personal observation and care since 1875.

The second exception, a German, aged 49 years at the time of his accident, a head bookkeeper for a large manufacturing firm, had high astigmatism and myopia of 12 D. He had been under treatment for eight years without increase of myopia and had pursued his work without difficulty. After an absence of two years, during the latter part of which he had been in impaired health and had suffered much from headache and weak eyes, he returned with a separation of his retina on the right side, which had come on suddenly the day before after reading late into the night. He was placed in a recumbent posture, atropia instilled and a compressive bandage applied. Under this treatment the detachment was slowly reduced and V. rose to 6/9. This continued for several weeks and he returned to his counting room, but not to work. He started from his home to my office and on the way noted in the cars that he was able to read a printed postal card with the injured eye alone. Reaching the city he ran a short distance and sprang onto a moving street car. The vision at once became blurred, and I found on his arrival at the office that the detachment had recurred.

In the third case detachment had occurred in the left eye when first seen, and in a few months occurred in the right in spite of treatment. Here also there was high astigmatism and additional myopia of 17 D. He was a professor of mathematics, busily engaged in his class work, a hard student, and, as he had a large family dependent upon his salary, refused to abandon his work even temporarily. His choroiditis progressed, a macular hemorrhage occurred and was speedily followed by extensive retinal detachment and blindness. With these exceptions I have seen no retinal detachment occur in my practice among patients under treatment. Macular atrophies which had not already occurred when first seen have been comparatively rare. I have seen no case of myopia progress under my own observation beyond 7 D., and but one of these. Any progression after the

first treatment and correction has been exceptional, and could then usually be ascribed to some imprudence upon the part of the patient, e. g., the neglect of the near work glasses or an inordinate amount of work. A number of cases of increased refraction have occurred after attacks of severe general disease.

These views could be exemplified by many illustrative cases. The following presents many interesting features:

CASE 1.—T. H. P., school boy, aged 8, was brought to me from the interior of Pennsylvania in March, 1879, suffering from weak and red eyes, headache and vision reduced to 1/10 in the right and 1/5 in the left. His mother who accompanied him and a sister aged 6 both had hypermetropic astigmatism. The father was deceased and the state of his refraction unknown. The sister's eyes were studied carefully, revealing a honeycombed choroid and a yellowish crescent of choroiditis at the temporal margin of both nerves. She often complained of headache. The eyes watered in the sunlight and became flushed. Treatment and correction of the refraction error was advised but the advice neglected. She was not seen again until 1898, nineteen years afterward, when she came under treatment with myopic astigmatism and a posterior staphylomata in both eyes.

In the boy the ophthalmoscope revealed an apparent myopia of 4 D. in the right eye and 3 D. in the left. He had a broad yellowish crescent in both eyes and the choroidal stroma was visible in large areas throughout the fundus. After the prolonged use of atropia he received for constant wear —1.5C—1c. ax. 20° in the right and 160° in the left which gave V.=20/LXIV in each eye not improved by a stronger glass. He returned in June, 1881, at the close of his school year with vision reduced with his glasses to 20/160 in the right and 10/200 in the left. The fundus changes had greatly increased. He then received the iodid and bromid of potassium internally and atropia was used three times daily for two weeks. He would then accept no change in the right glass, but vision rose to 20/40. On the left the myopia had increased to 2 D. The astigmatism remained as before. He was not seen again until 1896, fifteen years after. He was then driven from work as a clerk by his bad eyes and headache. He had in the meantime been under advice of a local surgeon, who had given him successively stronger spherical glasses until at the time he was wearing —5 D. sp. before the right and —6 D. sp. before the left, both for distance and for reading. With these V.=6/XV. The astigmatism had been neglected. He had blephorospasm and frequent attacks of violent sick headache. There were large posterior staphylomata, macular disease in both eyes with fluffy fundus and absorption areas. A mydriatic and smoked glasses were once more prescribed together with complete rest, and alteratives internally. These were continued for two weeks, when he selected the following glasses: O.D.—4.C—1.50c. ax. 30° V.=6/VII½, O. S.—5.C—2.00c. ax 150° V.=6/IX. He then returned to his work with an undercorrecting glass for reading and writing, but the headache speedily returned and vision fell in less than a month to 6/XV and 6/XXIV. A mydriatic was again instilled and while waiting in the office V. rose to 6/VII½ and 6/IX respectively. A careful study of the ocular balance showed an esophoria =10° at 6m. and a right hyperphoria of 5°. After sitting with the vertical prism and his correcting glasses the esophoria disappeared. He wore a trial prism for a week and returned without headache and no diminution in his acuity of vision. I then did a tenotomy of the right superior rectus, correcting accurately the vertical deviation. He returned in a month to his clerical work. His eyes remained weak for several months but gradually improved in all respects both in ability for work and in the fundus conditions. There has been no tendency to increase of the myopia and for five years he has pursued the arduous duties of a clerk in a railroad office free from any serious trouble.

Several points are illustrated by this young man's experience. In the first place, the repeated value of the therapeutic treatment by mydriatics, etc., but notwith-

standing this the steady advance of the myopia until both the astigmatism and hyperphoria had been corrected. Second, the evil of ordering, carelessly, successively stronger spherical glasses to improve the distance vision. Third, that in the presence of imperfect binocular balance the most carefully chosen glasses correcting the astigmatism did not secure his safety or comfort; and, finally, the arrest of the disease by correcting the ocular balance and by the careful correction of the anomaly of refraction. Time will not permit any extended discussion of many important points, but one must be noted, viz.: The importance of an under-correcting glass for all near work, and a full correcting glass for distance, both of which shall provide a common far point for each eye.

DISCUSSION.

DR. EDWARD JACKSON, Denver—It is, of course, impossible, and I suppose not desired, to attempt any complete discussion and review of this extremely interesting and valuable paper. I shall only take up, therefore, a few scattered points. With reference to the genesis of high myopia, in all of the cases that I can now recall, seen in early childhood, it has happened that there has been nothing pointing to heredity. I have seen comparatively few cases of high myopia in young children, but it happens that none of them had myopic parents. They did show, however, marked evidence of ill-health, disturbance of nutrition; and it seemed to me that was an important factor in the genesis of their myopia. I am also inclined to think it is important from the fact that the most rapid increase of myopia I have watched has occurred after some temporary severe impairment of the general health and nutrition.

There can be no doubt that the development of myopia during school life is common, and myopia developing in that way has two causes: one, the congenital defects, including astigmatism and possibly malformations of the orbit and attachment of ocular muscles; the other, the close work of school-life, and the use of the eyes under unfavorable conditions. We do not see in young people not exposed to school life the same development of myopia. Near work is one of the elements. Myopia in general may be taken as developing from these congenital defects and near work. In high myopia I believe an additional factor comes in in the use of the eyes under the conditions under which uncorrected myopic eyes must work.

My experience points to the importance of the careful correction of the refraction as a means of preventing the development of high myopia; that if that is attended to the myopia never becomes very high, unless it has already reached pretty nearly the limit before the correction is applied. Generally the progress ceases with full correction. It remains stationary for a time while there may be some change in astigmatism. But if change is once started so that the correction becomes imperfect, the myopia increases.

In the treatment, laying aside the correcting lenses for a time is important if the lenses are not the proper ones. Lenses should be laid aside until the correct ones can be determined. But, as soon as the correct lenses can be ascertained, the patient is better off to wear them. It may be wise to continue the mydriatic in some cases, but the correcting lens should be put on as soon as it can be ascertained.

With reference to the partial correction for near work, under Dr. Risley's teaching I practiced that for some time, but in time gave it up. While I feel that it is necessary in very exceptional cases, I think the better rule is to give the full correction for everything. I think the full correction, by encouraging the child to use the eyes more for seeing distant objects, is an extremely important means of influencing him to adopt those occupations that are not so likely to produce increase in the myopia. I feel about giving a partial correction for near work a good deal like Professor Haab feels about having a small size of magnet. If you give full correction the patient uses the eyes to the best advantage.

I would like to call attention to the literature of the past year on the subject, particularly the series of cases reported from Germany, by Pfalz and Heine before the Heidelberg Congress. Pfalz gave three series of cases, one of undercorrection, one of partial correction for near work, and one of full correction for near work. There were about 18 to 25 cases in each series. In the first two series there was but one case in which the myopia did not increase. In the third series, in which the full correction was worn constantly, the percentage that increased was extremely small. The discussion that followed the reading of these papers is still more remarkable, for not one raised his voice in defense of what we have regarded as the old German teaching, but all endorsed more or less completely the use of the full correction.

DR. RISLEY, in closing—I am obliged to Dr. Jackson for bringing out in the discussion the subject of full correcting glasses in myopia. In my paper I urged the wearing of under-correcting glasses only indoors and then only where the tendency to progression was marked. Notwithstanding the views expressed by Dr. Jackson, my opinion is in favor of using under-correcting glasses for near work and indoors in progressive myopia, otherwise there is almost sure to be headaches, discomfort and increased tendency to progression. I have but rarely seen any increase in the myopia when patients have been faithful to the undercorrecting glasses for near work. One other thing that I possibly did not make quite clear is that glasses should be carefully decentered, or ground on prisms with their bases in, certainly where exophoria is present and in extremely tender eyes where the strain of convergence at the near point is painful.

THE NEED OF CORRECTING AMETROPIA AFTER MIDDLE LIFE.*

C. M. CULVER, A.M., M.D.

ALBANY, N. Y.

The Century Dictionary says that "By a middle-aged man is generally understood a man from the age of 40 to 50." Presbyopia is usually supposed to begin between the ages of 40 and 46, although, with Donders' liberal definition of presbyopia, it may be present at the age of 30 years. Presbyopia, of itself, is properly treated by the wearing of an artificial substitute for the quota of accommodation that has been lost, because of the diminution in the elasticity of the crystalline lens—a diminution incident to advancing age. Convex lenses or an equivalence of their optical power in cases in which lenses need to be used for the correction of ametropia, constitute such substitute for lost accommodative power. This being true, it is primarily essential that concomitant ametropia receive due attention, when glasses for the relief of presbyopia are chosen, the effect of such attention having the form of modification of the strength of the lenses selected.

A common experience of the average oculist may be typified by a hypothesis of a patient whose eyes are hyperopic by one dioptre and whose age is 50: his near-work glasses have collective power, each, of two dioptres: he is fairly robust and wore no glasses till 47 years old, having avoided near-work as much as possible until it became inevitable, when he reluctantly yielded to invincible fate and bought these "No. 18" glasses of his jeweler or grocer. The vendor had a table, gotten at the same time with the sign: "Eyes examined free" and has not dared to go beyond "No. 18" in the case of a customer whose age was not greater than the figures opposite "No. 18," and whose spasmodic accommodation enabled him to read, for the length of time of a "free

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee: Drs. Frank Allport, H. V. Würdemann and J. A. Lippincott.