adhesion. At one point a pronounced ecchymosis was discovered near the mesenteric attachment. This was passed, however, and higher up a knuckle of ileum was found so dark in color that the restoration of its vitality seemed impossible. Accordingly about six inches of this portion of the intestine was resected, including the discolored portion. Safety-pins were used for clamps, the mesenteric vessels were separately ligated, and the cut margin united by a continuous suture. The ends of the ileum were brought together and united by Mannsells's method, except that the margins of the wounds were not painted with Woelfler's solution, nor was the iodoform employed. Before the union had been made the patient partially recovered from the ether, and on account of straining and efforts at vomiting, a quantity of blood and fecal matter escaped into the peritoneal cavity. This accident was attributed to the large size of the safety-pins. To disinfect the parts a large quantity of a fifteen-volume solution of hydrogen dioxide was poured into the cavity and allowed to remain during the completion of the operation. Finally, the cavity was flushed with normal sterilized salt solution, and as much allowed to remain as the cavity would contain. The external wound was closed without drainage. Recovery followed promptly, and by the ninth day the patient was allowed his usual diet. The points of interest to which the author calls attention are the following:

1. Hemorrhage, which in recorded cases, successful and unsuccessful, occurred only once (Croft's case), and this was very slight.
2. The slight amount of shock when the patient was first seen, four hours after the accident, the temperature and pulse being normal.
3. The excision of the contused and perforated portion of the ileum, and the end-to-end union by Mannsells's method, which is comparatively new and has not been employed many times.
4. The use of hydrogen dioxide in full medicinal strength for the purpose of disinfecting the general peritoneal cavity, this being the second successful case in which it has been used.
5. The closing of the abdominal wound leaving the peritoneal cavity full of hot sterilized salt solution, the object of this being to lessen adhesions, to lessen the danger of septic peritonitis, and to aid by osmosis the action of the intestine.
6. The shorter duration of the operation.
7. In the after-treatment the early administration of peptonized food (twenty-two and a half hours).

From a careful study of recorded cases the failure of surgical measures is shown to be due to:

1. Delay, which has been responsible for the largest number of deaths.
2. Hemorrhage.
3. Failure of the suture and septic peritonitis.

Delay in the future will be avoided by a better understanding of the early symptoms by the general practitioner. The important factors in the diagnosis are the history of injury, persistent nausea, hemorrhage, prolonged shock, rise of temperature, increasing rapidity and weakness of the pulse, increased frequency of respiration, rigidity of the abdominal muscles, persistent pain with or without pressure, and the facial expression. Hemorrhage after operation is to be avoided by greater care in tying the mesenteric

vessels, which should be ligated singly and not *en masse*. Failure of the suture will be less common with improved technique, as by Maunsell's method, septic peritonitis will be obviated by the liberal use of hydrogen dioxide, if infection is known to exist, and leaving the abdominal cavity full of bot sterilized salt solution. The author also calls attention to the importance of having an experienced and competent anæsthetizer.

CONCERNING THE ALTERATION IN THE LENGTH OF THE LOWER EXTREMITY IN DISLOCATION OF THE HIP-JOINT.

RIEDINGER (*Deutsche Zeitschrift für Chirurgie*, Bd. xxxvi., S. 102) states that the methods usually employed in determining the amount of lengthening or shortening of the limbs in cases of hip-joint dislocations give erroneous results. He claims that the measurement taken from the anterior superior spine of the ilium will show less alteration than if the measurement be taken from the middle line of the body. It is recommended, therefore, in such cases to take the measurement from the symphysis pubis. The comparative measurements are given in four cases. One of these was a perineal and another an obturator luxation. In these, measurements from the spine indicated shortening, but when taken from the symphysis a lengthening was shown as compared with the sound side.

THE ETIOLOGY OF CHRONIC HERNIA, WITH ESPECIAL REFERENCE TO THE OPERATION FOR RADICAL CURE.

BISHOP (*Lancet*, 1894, vol. i., No. 6), in an interesting article, shows by quotations from a number of eminent authorities the frequent failures that follow the operation of so-called radical cure. The author formulates the prevailing ideas as follows:

1. Hernia is a protrusion of viscera from their normal plane, through the walls which ordinarily contain them.

2. That viscera that are normal, contained within walls of normal strength, do not protrude.

3. If viscera protrude, such a condition is due to (a) an inordinate strength of their attachments (the mesentery); (b) an abnormal attachment of the mesentery to the back of the abdomen (Lockwood); (c) constitutional weakness of the walls and abnormal patulousness of the normal orifices (Mitchell Banks); (d) effects of injuries, operations, sudden strains in lifting, etc.

4. Reducible, incarcerated, obstructed, and strangulated hernia are all varieties of the same condition.

5. The results of operation for radical cure upon all varieties should influence our prognosis as to the prospects of any one.

Particular stress is laid upon the fourth proposition. Excluding congenital hernia, the author suggests that all other cases are naturally divisible into two classes—acute or strangulated and chronic or reducible; and further that these are distinct in causation, pathology, and course, and that their prognosis and treatment should be separately considered. Under chronic hernia are included the incarcerated and obstructed varieties.

Causation. The exciting causes of acute hernia are doubtless those

usually given, viz., strains, as by sudden lifting, etc.; but it is stated that these cannot produce chronic hernia, the causes of which are comparatively slight, and from within, but which act continuously or frequently. The one is unavoidable, but happens seldom; the other is curable or avoidable, but happens frequently.

Pathological appearances. The sac in chronic hernia is slowly produced, and is never tightly distended; it lies in folds at the point of exit from the abdomen. These folds radiate outward and upward over the parietes, and tend to unite when once the efficient cause is removed, so as to form a collar at the level of the ring.

Course. The course of chronic hernia may be divided into three stages:

1. Pre-hernia, or that which precedes any external evidence of hernia. It is peculiar to this variety.

2. The stage of growth, which lasts from the time of appearance to the period when the efficient cause ceases to act.

3. The stage of rest, when the efficient cause is removed and only subordinate or accessory causes remain operative.

In acute hernia there is but one stage which is terminated by operation, unless reprehensible delay is permitted, when there are two—the stage of tension and the stage of gangrene.

The results of treatment should be kept distinct. In the acute variety the state of the intestine is of the first importance; in the chronic variety, the condition of the abdominal wall. In acute hernia operation is a matter of immediate necessity; in the chronic form, one of convenience. In the acute variety the risk to life is great; in the chronic it is practically *nil*. In acute hernia the vitality is already, before surgical interference, greatly lowered, so that the expenditure of a few minutes more or less over the operation may make the difference between life and death to the patient; in the chronic variety, the time expended on the operation is of secondary importance, so that every detail as to technique may be carefully attended to. After operation for the cure of chronic hernia the foundation for relapse is often laid during the first week, when the parts are still in a plastic condition, and when, if the efficient causes are still acting, a return is unavoidable.

If the viscera protrude, it is due to:

1. The inordinate length of the mesentery.
2. Abnormal attachment of the mesentery to the spine.
3. Constitutional weakness of the walls and abnormal patency of the normal orifices.

4. The effects of injuries, operations, strains, etc.

The influence of constitutional weakness, and the effects of injuries, etc., the author meets with the following statements:

1. A careful study of pathological specimens of reducible hernia reveals none of the changes usually to be found in lesions of other parts of the body, due to sudden causes. It reveals, rather, very definite and well-marked differences between herniæ which have been produced suddenly, *i. e.*, strangulated or acute, and those now under consideration. The appearances, as actually seen, are such as are only compatible with the action of a comparatively gentle but prolonged continuous or frequently intermitting force.

2. A superficial inquiry in the case of a reducible hernia will usually elicit

a declaration of injury, but careful cross-questioning will frequently reveal some condition antecedent to the injury mentioned, which is the real cause.

3. Many persons suffering from hernia are men who have enjoyed continuously good health, are descended from healthy parents, are physically stronger than their fellows, and many have shown, by the fact that they have reached a ripe old age in spite of this drawback, that they are of good constitution.

In explaining the formation of a chronic hernia the author calls attention to the following points:

The abdomen is a closed cavity, the wall of which is composed of three layers; of these, however, the cutaneous may be disregarded, inasmuch as it offers no resistance to the outward progress of the hernia. The muscular coat is quite resistant, but is incomplete at certain points. The peritoneal covering is complete, but offers less resistance. As the contents accurately fill the cavity, if the abdominal wall contracts, the internal pressure is increased and there results slight bulging of the peritoneum at the points at which the muscular wall is incomplete. At first the peritoneum resumes its normal condition after the pressure is removed, but finally, after the bulging has occurred a number of times, the elasticity of the parts is diminished and the pouching becomes permanent. In this stage there may be no symptoms or if present are apt to be so slight as not to be noticed. There are, however, a number of exceptions to this rule, illustrations of which are cited by the author.

Attention is called to the fact that the folds which are seen at the neck of a hernial sac, and which radiate for some distance on the parietal peritoneum, are observed only in the acquired reducible form, and are never seen in a congenital or an acute hernia. The effect of bronchitis in causing a constant cough, and of occupations which frequently demand a strain upon the patient, are considered the efficient causes of hernia, which will surely develop once the pouching of the peritoneum has occurred, if the efficient causes remain active. The author has illustrated the views expressed by frequent references to specimens from the Museum of the Royal College of Surgeons and by numerous clinical cases.

In an attempt at radical cure of chronic hernia the choice of operation becomes a matter of great importance. Any plan will probably fail, however, if the cause which originally produced the condition is not removed. To be certain of its previous removal, it is obviously necessary that there should be no doubt as to its identity. This is the task which the author has assumed, and from his study he believes the following conclusions are justifiable:

1. That chronic and acute hernia are absolutely different things, agreeing only in the fact that they are both protrusions of viscera through their normal environments.

2. That they differ in etiology, pathology, and course.

3. That, especially from the point of view of radical cure, it is important to distinguish between them.

4. That in discussing the feasibility of operations for radical cure, and especially the permanency of their results, the etiology of chronic hernia is of immense importance, since no operation for the cure of the results of a cause is almost certain to be useless whilst the cause remains in operation;

and if the cause can be found the patient may be warned to avoid it after the operation has been performed, in order that permanency may be rendered more probable.

5. That in the case of chronic hernia the cause is never one acting suddenly and singly, as overlifting, strains, falls, etc., but always one acting slowly, persistently, gently, habitually (such causes are difficulties in urination and defecation, certain occupations, and, chiefly and most prominently, coughing in all its forms).

6. That the claims of any operation for the radical cure of chronic hernia cannot at present be properly estimated, and that they never can be unless acute and chronic herniæ are absolutely separated, and the true effective causes of them are duly appreciated, other causes being previously carefully eliminated.

ENTERECTOMY BY PAUL'S METHOD.

PAUL reports the following (*British Medical Journal*, 1894, No. 1727): Woman, aged fifty-one years, was admitted to the hospital, having a strangulated femoral hernia with the usual symptoms. After exposing the intestine, an area was found which had already begun to slough, and from which the contents made their appearance. An incision was therefore made in the median line of the abdomen, and after carefully cleansing the hernia it was drawn into the abdominal cavity, care being taken to prevent contact with the peritoneum in the proceeding. After clamping above and below, the injured portion was cut out with scissors, together with the corresponding mesentery, bleeding-points were ligated, and the divided ends united by a bone tube as recommended by the author. The proximal extremity was dilated, and the distal end contracted, so that it was found impossible to invaginate the former into the latter as the author had heretofore done. It was, therefore, necessary to employ a tube of small size ($\frac{3}{8}$ in. $\times$ $1\frac{1}{2}$ in.), fastening this in the distal portion, and by means of the traction thread this was then invaginated into the proximal end—in a direction, therefore, against the current of the contents, and although this method had been found to give good results in the case of experiments on dogs, the author hesitated somewhat about employing it in practice. The patient made a good recovery. During the first twenty-four hours the patient was allowed the yolk of an egg, two ounces of brandy, and some beef essence. This was gradually increased as the condition of the patient warranted.

HORRICKS details in the same journal the following case: A woman, aged thirty-eight, was admitted to the hospital with intestinal obstruction and with severe abdominal pain. Above Poupart's ligament, on the right side, a solid tumor was felt. An exploratory incision was made, which disclosed a tumor implicating a considerable portion of the small intestine. The wound was closed, and the patient recovered from the operation without complication. Three weeks later an operation was performed for the relief of the obstruction. A 3-inch incision was made in the median line rather above the middle. The tumor was drawn out of the incision, and, after emptying and securing the bowel on either side, a V-shaped section of mesentery containing enlarged glands and the affected bowel was excised. After securing the bleeding

vessels a Paul's bone tube was introduced into the lumen having the smaller diameter, and after securing with continuous silk suture it was invaginated into the other open extremity in the manner advised by Paul. The portion of intestine removed measured 39 inches in length; the tumor was found to be a large round-celled sarcoma. The patient made an uncomplicated recovery, and was discharged well at the end of five weeks.

THE DIAGNOSIS AND OPERATIVE TREATMENT OF SOLITARY HYDATID CYSTS OF THE SPLEEN.

AFTER a careful discussion and report of a case operated on by himself, and a study with a summary of sixty-six cases, TRINKLER (*Revue de Chir.*, February, 1894) comes to the following conclusions:

The facts cited speak clearly in favor of operative intervention in the treatment of hydatid cysts of the spleen. This intervention should have for its purpose the discovery of the cyst. Laparotomy, in the full acceptance of that term, should be employed—that is to say, with all the variations proposed by Messrs. Volkmann, Lindemann, Laudau, Säger, etc., accommodating itself to the peculiarities of each cyst.

It is difficult to judge in theory of the merits or demerits of operative procedures, for they all have their good points. There are cases in which Volkmann's operation is apparently less dangerous; nevertheless, when the tumor is small with thick walls and few adhesences, one can, without risking much, operate after the method of Lindemann, Säger, or Pozzi. The last method is the ideal, as the cyst is removed entire.

As a general rule, the incision should be as long as possible; it never does any harm, especially if one desires to employ Volkmann's method.

Possibly one might have to put a couple of stitches more at the top and bottom of the incision at the end of the operation, but it will give the chance to explore all parts of the tumor, to determine its boundaries, its pedicle, and, in short, one will see the actual condition of the organ, a most important factor in planning the operation. As Koenig says, before all one must "see," and this is only possible by a free incision. Consequently the first incision is exploratory and on it depends the choice of operation.

THE ANTISEPTIC POWER OF ICHTHYOL.

FROM interesting bacteriological researches which he has carried on in the hygienic institute of the University of Greifswald, ABEL (*Centralbl. für Bakt. und Parasit.*, 1893, Band xiv., No. 13) deduces the following conclusions concerning its antiseptic and bactericidal power:

1. The ichthyol preparations—ammonium ichthyol and sodium ichthyol—are capable of killing, in weak solutions and short time, the streptococcus pyogenes and erysipelas. The working of the different preparations is just about the same. In suppuration arising from these bacteriological sources, ichthyol can be used with good results, as has been already shown by actual experience.

2. The staphylococcus aureus and albus, the bacillus pyocyaneus, bacilli of typhoid, ozeana, and anthrax, the spirillum cholerae Asiaticæ, possess a