This article was downloaded by: [University of Illinois at Urbana-Champaign]

On: 10 March 2015, At: 16:48

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 6

Publication details, including instructions for authors and subscription information: <http://www.tandfonline.com/loi/tnah12>

LVIII.—On the Galapagos lizards of the genus *Tropidurus*

G.A. Boulenger

Published online: 12 Oct 2009.

To cite this article: G.A. Boulenger (1891) LVIII.—On the Galapagos lizards of the genus *Tropidurus*, *Annals and Magazine of Natural History: Series 6*, 7:42, 501-503, DOI: [10.1080/00222939109460655](https://doi.org/10.1080/00222939109460655)

To link to this article: <http://dx.doi.org/10.1080/00222939109460655>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

drooping, comparatively large; tongue, ocelli, and maxillary palpi invisible; forehead with a projecting fringe of hairy scales; antennæ broken; hind tibiæ with a broad expanding tuft of hairs in the middle.

Type *Isopteryx discoloralis*, Wlk. xxxiv. p. 1315.

[To be continued.]

LVIII.—On the Galapagos Lizards of the Genus *Tropidurus*.
By G. A. BOULENGER.

I HAVE lately reexamined the specimens of *Tropidurus* brought home by Darwin and Commander Cookson, with the view of testing the value of the supposed new species from the Galapagos Islands recently described by Cope* and by Baur†. The specimens known from the Galapagos Islands were referred by Steindachner and myself to two species, *T. Grayi*, Bell, and *T. pacificus*, Stdr., the form with two light dorsal stripes described by Peters in 1871, from Chatham Island, as *Craniopeltis bivittata* being regarded as a variety of *T. Grayi*. It is this very form which has been redescribed, from the same island, by Cope under the name of *Tropidurus lemniscatus*, sp. n., without any reference whatever to Peters's description. What is almost worse is Baur's bold statement, "Ueber *Tropidurus* (*Craniopeltis*) *livittatus* (sic), Peters, dessen Fundort unbekannt ist†, kann ich kein Urtheil abgeben." Is it too much to expect from herpetologists in America that they will look up the literature, and avoid quoting, as if seen by them, works to which they have not referred, as is evidently the case with Dr. Baur? I should also like to know where the latter gentleman has found *T. pacificus* quoted by me from Albemarle. I do not believe in most of the characters set forward by Baur to distinguish different species in the different islands; his statements are to a great extent contradicted by the typical specimens themselves, as when he says that the striped form, from Chatham Island, has 55–61 scales round the middle of the body, and that the specimens with 65 scales

* Proc. U. S. Nat. Mus. xii. p. 145 (1889).

† Biol. Centralbl. x. p. 475 (1890).

‡ Peters states in his original description, "Das einzige mir vorliegende Exemplar . . . stammt von der Galapagosinsel Chatham" (MB. Berl. Ac. 1871, p. 645.)

mentioned by me will prove to belong to the Charles-Island form, which has no stripes, whereas as a matter of fact it is precisely our unique adult striped individual (*T. bivittatus* = *lemniscatus*) which has the 65 scales. Here again we see that Baur has omitted to refer to Bell's original description in the 'Zoology of the 'Beagle,' where *T. Grayi* is figured in a perfectly recognizable manner; otherwise he would not have been embarrassed to guess whether the name *Grayi* was made to apply to the striped form rather than to the spotted one. And finally, whilst I trust nobody will contest the accuracy of the localities given by Commander Cookson, accompanied by dated labels, the specimens from Abingdon Island agree perfectly with Bell's *T. Grayi* and Steindachner's *T. pacificus*, forms held by Dr. Baur to be restricted to Charles and Bindloe respectively.

As the striped form, *T. bivittatus* (*T. Grayi*, spec. b, of my 'Catalogue'), appears to be distinguished constantly, apart from the coloration, by its larger scales on the sides of the neck and by the upper head-scales being more broken up, I think it best to restore it to the rank of a species distinct from *T. Grayi*. But I feel justified in rejecting all Baur's new species as based on mere slight varieties of *T. Grayi* and *T. pacificus*, as will appear from the following synonymy; I also give the number of scales round the middle of the body in the various specimens preserved in the British Museum.

1. *Tropidurus Grayi*.

Leiocephalus Grayi, part., Bell, Zool. 'Beagle.' Rept. p. 24, pl. xiii. fig. 1 (1842).

Tropidurus Grayi, part., Steindachner, Festschr. zool.-bot. Ges. Wien, 1876, p. 310, pl. ii. fig. 1; Bouleng. Cat. Liz. ii. p. 172 (1885).

Tropidurus albemarlensis, indefatigabilis, delanonis, duncanensis, Baur, Biol. Centralbl. x. 1890, p. 478.

1. Darwin's specimen, the type figured by Bell. No doubt from Charles Island. ♂. Sc. 63.

2. Darwin's specimen, probably from the same locality. ♂. Sc. 65.

3. Ditto. ♂. Sc. 61.

4. Ditto. ♂. Sc. 59.

5. Ditto. ♀. Sc. 61.

6. Albemarle, Commander Cookson. ♂. Sc. 57.

7. Abingdon, Commander Cookson. ♂. Sc. 59.

2. *Tropidurus bivittatus*.*Leiocephalus Grayi*, part., Bell, *l. c.**Craniopeltis bivittata*, Peters, Mon. Berl. Ac. 1871, p. 645.*Tropidurus Grayi*, part., Steind. *l. c.*; Bouleng. *l. c.**Tropidurus lemniscatus*, Cope, Proc. U. S. Nat. Mus. xii. p. 145 (1889); Baur, *l. c.*

1. Darwin's specimen, probably from Chatham Island. ♂. Sc. 65.
2. Ditto. Young. Sc. 57.

3. *Tropidurus pacificus*.*Tropidurus pacificus*, Steind. *l. c.* p. 313, pl. ii. figs. 2 and 3; Bouleng. *l. c.* p. 173.*Tropidurus abingdonii*, Baur, *l. c.*

1. Abingdon Island, Commander Cookson. ♂. Sc. 91.
2. Ditto. ♂. Sc. 95.

LIX.—On Anabiosis. By Prof. W. PREYER, of Berlin*.

THE fact of anabiosis, *i. e.* the revivification of perfectly lifeless organisms and their parts, the condition of which differs from ordinary apparent death in the total suspension of the whole of the vital processes, was experimentally established by me more than twenty-five years ago; and since that time I have substantiated it in my lectures and various papers, and have also drawn attention to its great theoretical importance †. I attach especial weight to the proof which I have furnished of the difference between the two antitheses of life, namely:—

- i. Lifeless and capable of living=anabiotic.
- ii. Lifeless and incapable of living=dead.

* Translated from the 'Biologisches Centralblatt,' Bd. xi. no. 1, Feb. 1, 1891, pp. 1-5.

† 'Der Kampf um das Dasein,' Bonn, 1869 (pp. 10, 39); 'Die Erforschung des Lebens,' Jena, 1873 (and 'Tageblatt der 45. Versammlung deutscher Naturforscher und Aertze,' Leipzig, 1872, p. 47); 'Naturwissenschaftliche Thatsachen u. Probleme,' Berlin, 1880; 'Der Hypnotismus,' Berlin, 1882 (p. 282); 'Elemente der allgemeinen Physiologie,' Leipzig, 1883. Cf. "Ueber den Lebensbegriff" in the journal 'Kosmos' (i. Jahrg. ii. Bd. p. 213, Leipzig, 1878), and "Die Wiederbelebung totenstarrer Muskeln" (Amtl. Bericht über die 39. Versammlung deutscher Naturforscher und Aertze in Giessen, 1864, and 'Recueil des travaux de la Société médicale allemande,' Paris, 1865).