undoubtedly most of these were the peripneumonia organisms. But some of the particles I am sure were not organisms, for they were too large and varied too much in size. These particles unquestionably were formed as a result of changes occurring in the medium due to the growth of the microbe. The organisms, not being motile, can not be distinguished clearly from other particles which may be present. However, uniformity in the size of the particles and distinctness in their appearance speak in favor of their being organisms rather than particles of proteid or other material. In dealing, then, with these extremely small organisms, the observations must be controlled most carefully.

Sections of tissue are not suited for the application of this method. In teased specimens, however, one can observe striation in muscle and the course of fibers in brain tissue. I could observe nerve fibers ramifying between cells, and in several instances I was able to see clearly the axis cylinder coming from the nerve cell and to trace it for some distance. The method may be of some value in studying the branching of the nerve fibers.

POSSIBILITIES AND LIMITATIONS.

The possibilities and limitations of the apparatus described here can only be determined by a large number of observations. New methods usually fail to meet all our predictions, yet they often become of service in some manner entirely unforeseen at first. The progress of science depends very largely on the development of new methods of technic, and consequently all such methods should be most thoroughly investigated before being discarded.

There is undoubtedly a large field for the application of this method in studying the structure of that important class of substances known as colloids, both organic and inorganic. This will include such substances as toxins, antitoxins, ferments and the like. The work already done by Raehlman is extremely suggestive, and indicates a most important field for further study.

Emphasis must be placed on the fact that the value of the method is by no means limited to the study of ultramicroscopic particles. One of its greatest uses will be to observe particles or objects, not too small to be seen by our ordinary microscopes, but which, owing to their slight refractive properties, can not now be seen by our present methods. Because of the property of this method of bringing out slight differences of refraction, I am inclined to believe it will replace, for many purposes, at least, the hanging-drop. That we can see objects not seen in the hanging-drop is illustrated nicely by the fact that red blood corpuscles in dilute acetic acid can not be seen either in hanging-drop or under a cover-slip when unstained, but under lateral illumination the stroma of the corpuscles is rendered plainly visible. Now it may be well that there exist important bodies whose refractive powers are too slight to enable them to be seen in hanging-drop, and which can not be stained by our present methods. In such cases this method of observation will be of positive value.

Since the method is applicable only to particles or small bodies in suspension, it apparently can be of no service in determining structure of tissues. The form of larger objects may be observed, but since the method does not magnify, the form of the smaller particles can not be determined.

In respect to the simplicity of the apparatus and to ease of examination of any fluid, this new method surpasses even the familiar hanging-drop.

Finally, I would call attention also to the fact that the few observations reported here were made with an imperfect apparatus.

A much more perfect illuminating system is desired, so as to obtain a smaller and more intense focus of light. This applies particularly to the study of ultramicroscopic particles, for it is generally true that the stronger the light the smaller are the particles which may be seen.

I wish here to express my thanks to Dr. Hektoen for suggestions.

THE DANGERS OF POTASSIC CHLORATE.

PAUL BARTHOLOW, A.B., M.D.

PHILADELPHIA.

Every obscure subject in medical literature should be reviewed from time to time on principles broader and more general than can usually be employed in text-books or special articles; and particularly when such obscurity is caused by a want of precision of language, by errors in the use of terms—faulty or loose terminology—and, what is more serious, by a failure to distinguish between two objects, which, though alike, are chemically and physiologically speaking, separate agencies.

For instance, a frequent and widespread source of error—and consequently of danger—arises from a confusion in the nomenclature. That is to say, potassic chlorate has been confounded, both in medical writings and in prescriptions, with potassic chlorid (KCL). Secondly, by a false analogy, the chlorate has been thought to have properties and effects identical with the chlorid, and not to be any stronger. For the belief that potassic chlorate and potassic chlorid are interchangeable, or at least as alike as twins, it is customary among some writers to blame the pharmacopeias. Our own, as it will be seen, is not liable to this criticism; but the German offends heavily. Thus a high authority¹ says: "In the pharmacopeia by the term potassic chlorate is understood potassic chlorid (KCL), while chemists apply the former name solely to the chlorate." A similar remark is made by Kobert.² On the inelegance of this ambiguous use of scientific terms, Virchow³ comments with severity. In spite of these strictures from such good authority, it is possible to find in medical literature, numerous instances where the term "chlorid" is used where the "chlorate" is meant; and, sometimes in the course of the same paper both terms are employed so loosely that it is frequently difficult to tell from the context which is to be understood.⁴ Finally, not only must the chlorid and chlorate be clearly separated, but both also must be carefully distinguished from the "hypochlorites" (KClO, CaClO₂), often improperly called "chlorids," and in this false character responsible for some cases of poisoning.⁵

Into the error of confusing terms a writer in the *Therapeutische Monatshefte*⁶ falls, while pointing out another blunder. Speaking of the danger of mixing potassic chlorate with organic or combustible substances, he cites as an instance of the non-observance of this rule the formula of a French surgeon where potassic chlorate is prescribed in conjunction with saccharin. The result was a violent detonation.⁷ This mistake, absurd as it

1. Das Chlorsaure Kali, J. von Mering, Berlin, 1885, p. 1.

2. Pharmakotherapie, vol. 1, p. 55.

3. Barbarismen in der med. Sprache.

4. Dictopaeus: Archiv f. path. Anat., vol. xxxiii, p. 510.

5. Dict. de Thér., vol. ii, p. 1.

6. 1894.

7. Rev. de Thér. méd. Chir., 1894, p. 241.

may seem, is natural enough. So, too, an examination of the literature will show that forgetfulness of the explosive power of potassic chlorate has been surpassed by a belief in the harmlessness of the salt. As long ago as 1830, there was a well-established opinion of this kind, an opinion based on experiments as ably conceived and carried out as most of the later ones. Thus, in that year, we find the matter lucidly and with great practical definiteness set forth by Professor Tully. He repeatedly gave, by way of trial, not only to his students but to himself, an ounce of potassic chlorate at a dose. No ill effects beyond slowing of the pulse and moderate intoxication were seen. It is a noteworthy evidence of the accuracy of Tully's experiments⁸ that he first satisfied himself by preliminary tests that he was using true potassic chlorate and not some colorable substitute. In this respect, as well as in other ways, his investigations compare favorably with those carried on in the laboratories of Europe.

Subsequently, Isambert⁹ declared himself in support of the belief in the harmlessness of potassic chlorate. He took it in doses of 20 grams, and observes that it is "d'une innocuité singulière." Again, Chaussier affirms that it may be taken without solicitude (inopinément). Another reports the case of a man who inadvertently took 9 drams without bad results. Such cases were common at one time—a fact remarked by Jacobi, who was one of the first to point out the danger. It was customary to give potassic chlorate in large amounts. It was known not to be very soluble, and for this reason as well as for its postulated safety, it was often administered in saturated solutions.¹⁰ Even at the present day there are to be found some who maintain that the danger is almost a negligible quantity.¹¹ The case of Dr. Fountain, so well known, is an instance not so much of individual rashness as of the singular variability of the action of potassic chlorate. It will be remembered that he died from the effects of taking an ounce, a quantity as we have seen, that was frequently taken without harm. This case, together with one reported in Germany in 1878, were thought the only two recorded.¹²

As an explanation of that variable quantity—the toxic power of a salt—it has hitherto been found useful to employ the term "idiosyncrasy." In a negative way, this hypothesis has been of use in enabling us to describe a latency of poisonous power in a chemical body—a latency exploding into virulence under given circumstances. Latterly this idea has expanded into the notion of the pharmacodynamics of a salt, namely, the power it possesses as an entity, that is, before its molecular structure has been changed. These pharmacodynamic effects, according to Gruetzner, are proportionate—not to the absolute weight of a salt, but to its molecular weight.¹³ It is, therefore, instructive to note that potassic chlorate acts in a state of almost complete molecular integrity, as it passes practically unchanged through the body; and, judging from the way in which it develops a toxic strength, it seems likely that its molecule is endowed with an uncommon potentiality, for its pharmacodynamic effects are often irresistibly sudden and fatal. This sudden explosiveness of poisonous power is to be clearly seen in the cases where the salt has been merely used as a gargle and has caused

death. On this postulate of its latent danger it is possible to rest a good case against potassic chlorate; and on this basis of fact it is possible to reason out the tangle of results protocolled in the "casuistik" of the experimental pathology.

In this field a wide divergence of opinion reigns. Laborde, for example, denies that intravenous injections of potassic chlorate, even in large amounts, have any effects worth mentioning. Stokvis holds that it is of no greater power than common salt, and that it in no way alters the blood elements. This proposition is true of the action of solutions, both isotonic and hyperisotonic.¹³ This view is opposed alike to common sense and to a conclusive clinical experience.¹⁴ Yet it seems to be a fairly well-established opinion among laboratory experts. Thus Riess and Bokai¹⁵ were unable to find the absorption bands of methemoglobin in the spectroscopic analysis of the blood of animals poisoned by potassic chlorate; nor was there any ocular proof of blood changes, such as chocolate discoloration, etc. On the other hand, in cases of poisoning in the human subject, the "striae" of both methemoglobin and hematin have been spectroscopically demonstrated, and what is more, during life.¹⁶ In fact, no one of any practical experience can doubt that potassic chlorate may profoundly alter the composition of the blood. What an abundant clinical experience now shows is, that this alteration of the blood may take place suddenly, and after comparatively small doses. In several instances a small dose, when taken on an empty stomach, has caused death. It shows its utmost power under these conditions, viz., where there is no mechanical interposing agency (such as food, aqueous contents of the stomach) between the salt and the avenues into the blood. It is under these conditions, as any valid observation will show, that potassic chlorate may display a treacherous power.

Importance of the Tongue in Diagnosis.—The old notion that the condition of the stomach lining was reflected in the mucosa of the tongue has been long exploded and even the débris have been scattered to the winds by a communication in the *Gazette des Hôp.* of Sept. 17, 1903, in which Mathieu and Roux discourse on the diagnostic value of the state of the tongue in affections of the digestive tract. They establish that the mucous membrane of the tongue is a false mucosa. It is not formed from the inner layer, but from the outer layer of the blastoderm. In structure it resembles the skin. The "coated tongue" is not due to an actual coating, but is, in effect, a grass plot of long and thick filiform papillae, and the white color is due to a superficial desquamative dermatitis. This dermatitis arises and subsides under the influence of external and internal intoxications, like any cutaneous affection, with the important difference that the tongue is much more intensely sensitive than the skin, owing to its wealth of nerves and vessels. This lingual dermatitis is, therefore, an extremely early and important sign of internal derangement, and thus the tongue retains all its old diagnostic value. The only difference is that the knowledge that this dermatitis may persist after the cause has subsided should suggest that we must not wait for the tongue to clear before instituting nourishing feeding.

8. Boston Med. and Surg. Jour., 1832.

9. Thèse de Paris, p. 14.

10. Küster: Deutsche Zeit. f. prak. Med., 1877, No. 33.

11. Berlin. klin. Woch., 1897.

12. Deutsche med. Woch., 1893, 52; Archiv f. exp. Path., 34, p. 314.

13. Bull. gen. de Théor., 1874, p. 247; Stokvis: Archiv f. exp. Path. u. Pharmak., vol. xxi, p. 169.

14. Marchand: Ibid., xxii, p. 201; Virchow's Archiv, lxxvii, 455.

15. Centralbl. f. Physiol., 1887, pp. 215 and 243.

16. Jacob: Berl. klin. Woch., 1898.