served from disease or extracted when useless, it is dentistry.

So the status of dentistry to-day, as far as our medical brothers are concerned, is not that of a specialty of medicine, principally because we are not graduates in medicine. Are they right? To those who may believe that they are not, it is comforting to feel that though their maternal acknowledgment is so dear and their association so keenly desired, they shall not be permitted to sit sole judges of our relationship.

Other influences, both external and internal, are doing much to drag us from this fireside of art and science. The first, we are led to believe, is the necessary amount of mechanics our profession is obliged to teach and practice; 2, the distinctive degree and curriculum we are attempting to maintain and enforce; 3, the percentage of our profession who are not graduates—many of whom, though, are our bright lights; 4, our determination to be recognized.

The forces and influences which are tending to draw us beneath the maternal roof are, however, such that they will eventually overcome all opposing ones. They are:

1. The addition of medical chairs to dental faculties, and the division of dentistry into the operative and the mechanical dentist.

In all large communities men who make a business of doing mechanical work for one or more practitioners are fast becoming a special department of dentistry. These mechanical dentists in many cities do a very large percentage of all the mechanical work, and in some States the law is inclined to wink at the infractions of these unlicensed.

2. The patent dissatisfaction in which the degree is held, and the disinclination of colleges and universities to settle upon the not unusually adopted degree of doctor of dental surgery, together with the tendency to establish a variety of post-graduate degrees for the purpose of advertising those decorated and the college conferring it.

3. The rapid percentage decrease of the body of non-graduates and the tendency of dental and medical graduates who intend to practice dentistry to acquire both medical and dental degrees.

4. The establishment of the Oral Section of the AMERICAN MEDICAL ASSOCIATION.

The elevation of any particular class of men is, however, dependent upon a demand on the part of their constituents; the people. The greatest advancement of any profession or body, the pulpit, the bar, the medical fraternity, or what not, is the result of enlarged conception on the part of their representatives, and that conception has its stimulus in the public demand.

The forces, then, of successful issue lie in the education of our patients, in the establishment of recognition from them, occasioned by scientific efficiency, and an elevated qualification in the average practitioner.

No amount of recognition from a National, State or local medical association given a body of men undeserving, can be made to stand the test of popular opinion.

We therefore believe the future course of those who are so ably leading, to be not only through the channels of education as it may present itself in the education of dentists in the specialty of medicine,

but that theirs is a plea for text-books and for education of the people.

Complete recognition can only, and will result, in the displacement of the dental degree by the medical for all, with special training for special practice.

DISCUSSION.

DR. R. H. COOL—I am interested in the paper, especially in its advocacy of a wider range of knowledge for the dental practitioner. I take the opposite view, that more knowledge of oral and dental surgery and of its resources by medical men is needed. More and more the best surgeons and best physicians are coming to call in the dentist in oral lesions. As an instance of the value of the dentist's knowledge in these cases, there is no known splint for a broken jaw like that made by an experienced dentist. In antral troubles the surgeon also calls in the aid of the dentist. With regard to instruments, those shown by Dr. Talbot for perforating the antrum produced the impression that they would give a great deal of pain, while our dental engine makes a prettier opening much easier. We can almost say that dentistry is now acknowledged among the leading surgeons as a nucleus for dental information, and I think we are being recognized as a professional body by them.

DR. A. F. McLAIN, Santa Rosa—I agree with Dr. Cool as to the advantages of skilled consultation between surgeons and dentists in diseases of the mouth and jaws. A student at the dental college contributed to a successful result in a case in which there was a fracture of the jaw near the ramus, which may have been comminuted. This student took the case in charge, made an interdental splint for it, and I defy any one to say that the result was not everything that it should be.

DR. M. J. SULLIVAN, San Francisco—I heartily indorse the tone of the paper. I have always believed in a liberal medical education for the dentist. The better the foundation the surer the success, both theoretical and practical.

SOME PATHOLOGIC NOTES ON NEOPLASMS OF THE MAXILLÆ.

Read in the Section on Dental and Oral Surgery, at the Forty-fifth Annual Meeting of the American Medical Association, held at San Francisco, June 6-8, 1894.

BY VIDA A. LATHAM, F.R.M.S., D.D.S.

CHICAGO, ILL.

Among the conditions associated with the pathology of the oral cavity and its contiguous parts, none are perhaps more interesting than those relating to the new formations which may be observed there. A pathologic formation may be simply an excessive production of the normal structure and is a morbid condition only in respect to the amount and arrangement of the ordinary tissues of which it is composed, or of the region in which it is situated. These are known as "homologous" growths and form a large number of all the pathologic formations throughout the body, and include the hyperplasias, exostoses, congenital tumors, and many growths of varying origin, such as the arthritic enlargements, formations following a rheumatic attack, or the chronic inflammatory enlargements due to long continued irritations. The main and most important class of neoplasms are those belonging to the "heterologous" class in which the structural elements are different from the tissues in which they are situated, or with which they are connected. The true distinction is one which depends upon the histologic composition, rather than upon the gross external appearance or upon the clinical peculiarities. The great and only reliable distinction is the microscopic, for upon this depends to a very great extent the correct estimation of the nature and tendency of any pathologic formation, both in relation to its effect upon the tissues with which it lies in direct contact, as well as

its influence upon the well being of the individual. Certain growths are found to consist of a tissue different from any found in the adult body, and therefore, unlike any of the tissues in which they are situated. These are very confusing to the pathologist and they have to be referred to some of the immature forms of the tissues, or to their derivatives. We must, in fact, return to embryologic tissues, for we only find the embryonic types sparingly distributed in adult life or limited to certain parts.

We usually regard growths of mesoblastic origin as of a malignant type, but still the benign forms are found as fibroma and so forth.

Granulation tissue or embryonic connective tissue is not a permanent structure. It only exists in the body of the embryo as a transitory material, the conversion of which produces another and entirely different material. We can easily understand if embryonic tissue re-appears at a time when no process of repair is going on, and accumulates in any great amount, that it may undergo similar changes to those belonging to it in its usual and normal state. It is never a permanent tissue, but is always changing to other and different forms of structure. When a new growth appears it exhibits a tendency to undergo further changes, either extension of disease, or more often a rapid decadence or retrogression of the tissue already formed, and lastly it may be combined with almost any form of tissue in the body.

Text-books on oral surgery and pathology convey erroneous impressions by classifying neoplasms of the oral cavity as ordinary types; but a careful examination with the microscope reveals the fact that they are much more complex and peculiar. It is to this "mixed" class that I find the majority of the tumors of the maxillæ and oral cavity belong, basing this statement on the microscopic examination. These growths are also much more common than is usually thought to be the case, particularly so in the region of the neck, reaching from in front of the submaxillary gland to the parotid, and surrounding the angle of the jaw. Nearly every specimen obtained so far from this region has been of this peculiar type of growth, bearing no resemblance to any fixed type of tissue and having only an ordinary clinical history. The diagnosis made by the operator has in almost every case proved to be erroneous when the growth has been examined microscopically. The commonest varieties perhaps are epithelioma and sarcoma, the latter usually a peculiar mixed form between the round-celled, spindle-celled and osteo-sarcoma. Enchondroma undergoing sarcomatous degeneration is also met with.

Tumors of the salivary glands are almost always confined to the parotid, being very rare in either the submaxillary or sublingual glands.

The commonest type of the parotid tumor is one composed of a mixture of cartilage, myxomatous tissue, a varying quantity of connective tissue and ill-formed glandular acini. Growths of this nature are often called "parotid glandular tumors" or "adenomata," but it is quite rare to find a growth the bulk of which is composed of true gland tissue. The cartilage of these tumors is hyaline and it is by a mucoid degeneration of its stroma, and that of the fibrous tissue of the growth that the myxomatous tissue is produced. Sometimes there is a considerable mixture of sarcomatous tissue; at others the growth is chiefly cartilaginous; and as might be expected

from this variety in structure there is a corresponding difference in the rapidity with which such tumors develop. These growths appear in middle life and are found in either sex. They occur most often in that part which overlaps the lower jaw, are usually encapsulated, and when small adhere but little to surrounding parts. They attain large size if left alone, and may cause death by encroaching upon the pharynx or large vessels. The lymphatic glands are seldom affected, and although the growths may recur locally after removal, they seldom or never become disseminated.

Malignant growths are happily rare in the parotid. About the only growth which occurs in the submaxillary gland is made up of cartilage and even this is rare. According to Bowlby, there are not more than seven or eight cases on record and they are usually enchondromata.

Tumors of the maxillæ possess so many distinct features as to almost require a separate description. These distinct characters depend partly on the proximity of the neighboring mucous cavities, partly also on the presence of the teeth and of the fetal structures from which the latter originate.

Epulis is a loose term often applied to any tumor which appears on the gum, but is best limited to the fibromata. It usually develops from periosteal tissue and is occasionally attached to the periodontium and very often extends into the alveolus. It necessarily compels the extraction of a tooth or teeth and excision of a portion of the alveolar margin. A simple diffuse epulis is usually only a thickening of the gum.

Myeloid sarcomata are often included as myeloid epulides growing within and expanding the jaw bones, presenting rounded, tense and elastic swellings often feeling suspiciously like cysts; and generally found in young people. Other varieties of sarcomata occur within the jaw bones and also within the antrum. The cells are round or oval. The growths being soft and rapid in development, tending to extend all around and to fungate through the skin, affecting the lymph glands and so disseminating.

The distortion of the antrum and its relative structures is too well known to require description. Any tumor growing in this region tends to distend the cavity in this way, but malignant tumors do so in a more typical manner than do the benign growths. Fibro-sarcomata are found in the inferior maxilla (mandible) are not very malignant and when completely removed with the bone seldom recur.

Carcinoma are most often found in the antrum¹ and develop from the epithelioma lining the antral cavity. They are rapid in their course, affect the glands, rapidly disseminate and are usually found in elderly people.

Multilocular cystic tumors, so-called by Mr. Eve,² and known as "cystic epithelioma" stand in the position of a mixed growth—a solid and a true cyst. They are most common in the inferior maxilla and are endosteal or central growths and possibly originate by an ingrowth of the epithelium of the gum and following caries or traumatism of the teeth. They occur at any age but usually between 20 and 40, are slow in growth and may take years to reach any size. The growths are usually not malignant and seem not to affect the glands or disseminate. The

¹ New York Medical Journal, November 10, p. 586, and November 17, p. 612.

² British Medical Journal, Jan. 6, 1883.

bone is usually expanded on its inner side, and is very hard to the touch, sometimes elastic and crepitant. The teeth loosen, fall out, and a sticky discharge comes from the alveoli. Macroscopically the bone shows expansion, the cavities are filled partly with a solid growth of a soft fibrous or fleshy material, partly by fluid, serous or viscid. The septa dividing the cysts are often incomplete and are either fibrous or bony.

Periosteal cysts are situated beneath the periodontium, generally inflammatory in origin and are found in connection with carious teeth or with imperfectly extracted roots. They are very painful.

Antral mucous cysts may also be found and must be differentiated from malignant growths as they produce a similar distension by their slow growth and fungation; and as the bone turns the presence of fluid is soon detected.

Dentigerous cysts and odontomes³ are quite well described in our dental surgeries and pathologies.

Diffuse osseous growths of the antrum are rather peculiar in that they diffuse over the whole area of the antral wall, and do not grow from some definite portion as is the usual mode. Therefore we have no definite tumor, but a general thickening of the whole bone which incloses the antrum and a gradual obliteration of the sinus by the new bony formation. It may be confined to the antral cavity or may extend to the superior maxilla. The growth is slow, occurs in youth and is benign.

Cartilaginous tumors of the superior maxilla are extremely rare and usually combined with sarcomatous elements.

Sarcomata. This group which is mainly due to the exertions of Virchow, includes the fibro-cellular, the mucous tumor, and the myeloid of English surgery. The group, on the whole, nearly corresponds to Paget's recurrent fibroid. The best classification of this group is perhaps a strictly pathologic one after Billroth:

1. Round-celled sarcoma, granulation-sarcoma (including glioma) consists of mesoblastic corpuscles; the inter-cellular substance is homogeneous, striated, reticulate and varies widely in amount.

2. Spindle-celled sarcoma. The cells are usually well-marked, intercellular substance is absent or scanty, homogeneous or fibrous. The most recurrent sarcomata contain this tissue but do not always recur.

3. Giant-celled sarcoma, myeloid. This has the general structural elements of one of the other varieties and has in addition many large multinucleated cells.

4. Net-celled sarcoma—mucous sarcoma. This is not exactly like the myxoma. Myxomata are sarcomata of various kinds and are similar by having a gelatinous appearance. Net-celled sarcomata contain stellate cells with long processes and gelatinous intercellular substance.

5. Alveolar sarcoma are very rare.

6. Melanotic, may be combined with alveolar. Glandular tumors secondary to epithelioma of the lip or tongue become adherent to the lower jaw in the later stages, but must not be confounded with growths originating in the bone itself.

These are, in brief, the most important varieties of neoplasms met with.

With regard to frequency. The superior maxilla seems to be most often involved and usually recovers

itself, on account of its high degree of nutrition. In an article in *Archiv. für Klin. Chirurg.*, (Vol. xviii, 25, 1875) an account is given where the entire jaw was removed fifteen times:

"Twice for carcinoma, five for sarcomatous growths as giant-celled, three as spindle-celled, one as round-celled and one as medullary. Cylindromata twice required partial resection; epitheliomata twice an almost complete resection; enchondroma one, a total resection. Mortality being 15 per cent. According to this proportion carcinoma is more common than medullary sarcoma."

This does not agree with Prof. C. O. Weber's account: "The usual result of carcinoma is a recurrence after about one to three years and usually recurring in the cicatrix."

The diagnostic signs of antral growth are as follows: When springing from the interior of the antrum, the orbital, nasal, palatal and buccal walls of this cavity are expanded, according as the tumor grows upward into the orbit, inward into the nostril, downward into the mouth, or forward upon the cheek, causing a prominence of one side of the face. A tumor of the antrum in many instances grows simultaneously in all these directions, producing exophthalmos, obstruction of the nostril, prominence of the cheek, and bulging of the roof of the mouth. When springing from the anterior surface of the superior maxilla it causes a projection of the cheek, dipping down between the gums and the soft structures of the face. It may extend backward into the interior of the antrum by increase in size, thus secondarily implicating that cavity. When springing from behind the superior maxilla the upper jaw is first pushed bodily forward, but as the antrum becomes secondarily involved, any of the walls of this cavity subsequently become expanded and rendered prominent.

A serious affection of the gums appears in a more innocent garb, apparently a mere ulcer. Dental surgeons are more apt to see it in its earliest stages than are surgeons. A sore appears, perhaps awakened by a carious tooth. It does not heal kindly, but exhibits a tendency to extend. Its borders become hard and thick, its base soon adheres to the bone beneath—this is cancer.

"Bone is never the seat of cancer primarily." This broad statement was first made some years ago by the best German pathologists, and cancers of the superior maxilla were held up as positively disproving any such idea. But now we know there are but two locations in the superior maxilla where cancer is first encountered. The first appears as above—upon the gum; the second in the antrum of Highmore. In each case it starts as an ulcer, eats its way into the bone, and never appears as a distinct tumor. The sore is found upon the gum, its edges thick and firm, base immovable, and this characterizes its malignant infiltrating tendency. Dentists can not have these facts too strongly impressed upon them. It will be noticed that carcinoma of the body of the superior maxilla conceals its true nature at first, and except by a careful person is not recognized until extensive destruction of bone has resulted, possibly an opening made in the oral, nasal or orbital cavity and is soon beyond operative relief.

Butlin has removed the upper jaw within seven weeks of the first signs of the disease, only to find the bone entirely destroyed and sinuses burrowing into all the

³ New York Medical Journal, p. 612, 1894.

surrounding parts. Thus we see the necessity, when malignant disease is suspected, to examine not only the mouth, nose and orbit, but even the throat, and here comes the necessity for broad medical and dental education in the combined efforts of thorough dentists and surgeons. Professor C. Heath gives an excellent suggestion, and a most valuable one, when he says the attachment of any growth to the roots of extracted teeth should excite the suspicion that a serious disease is present in the antrum, and a microscopic examination should be made.

In conclusion, we have seen from the foregoing remarks that neoplastic growths of the maxillæ are then modified forms of the great groups, carcinoma or sarcoma. They are, in fact, more truly complex and diffuse in structure, so much so as to require special grouping; for in structure the histologic-elements are not allied as is usually the case in growths found in other parts of the body. I regret to say that lack of time prevented the subject being brought before you in a more thorough manner, but as their structure is so complex more special study was required. It is hoped this may be completed later and published with photographs and cases. Might I suggest that members of the Association and profession kindly collect such specimens and examine them, or aid by loaning or giving specimens for examination? They would thus enable better and more complete work to be done in this line, and so preserve valuable material. They would confer a great favor upon the essayist, and at the same time help to make understood some of the great questions which occur in oral pathology, especially in the histologic details.

BIBLIOGRAPHY.

- Heath, C., Injuries and Diseases of the Jaws.
 Heath, C., Lectures on Certain Diseases of the Jaws. Jacksonian Prize Essay.
 Butlin, Malignant Diseases.
 Sutton, J. B., Tumors, Innocent and Malignant.
 Bowlby, Surgical Pathology.
 Pepper, Surgical Pathology.
 Butlin, On the Tongue.
 Southam, Regional Surgery.
 Ziegler, Pathologic Anatomy.
 Treves, Manual of Surgery. (Cassell & Co.)
 American System of Surgery.
 American System of Dentistry.
 Garretson's Oral Surgery.
 Colver & Smale, Dental Pathology.
 Blodgett's Dental Pathology.
 Dental Cosmos.
 Journal of Anatomy and Physiology.
 Proc. of the Odontological Society of Great Britain.
 Lancet.
 British Medical Journal.

DISCUSSION.

DR. JACOBS—Some seven or eight years ago I studied the structure of tumors considerably, but I have done very little work in that direction lately. I should not therefore dare to discuss the paper further than to confirm it as far as I know, and to say that Dr. Latham is the hardest worker I ever knew, and I have no doubt of the correctness of everything she says.

SOCIETY PROCEEDINGS.

Medical and Chirurgical Faculty of Maryland.

*Semi-Annual Meeting held at Cumberland, Maryland,
 Nov. 21 and 22, 1894.*

FIRST DAY.

The Medical and Chirurgical Faculty of Maryland convened in session in the council chamber of the city hall at Cumberland, Nov. 21, at 10 A.M. After the Address of Welcome by Dr. J. M. SPEAR, of Cumberland, and the Address by the President, Dr. R. W. JOHNSON, Dr. H. SALZER read a paper on "Distension of the Stomach," in which he spoke of the methods of diagnosis of diseases of the stomach, and especially distension of that organ, by examination of the gastric secretion. He did not always use the stomach

tube, but let the patient vomit the substance taken, and then made an examination.

DR. WM. OSLER thought that the method of Dr. Salzer was more pleasant than the one by the use of the stomach tube. He thought that the examination for movable kidney, as suggested by Dr. Salzer, was important. He generally found signs of dilatation of the stomach in neurasthenic patients, and palpable if not movable kidney, and it may be at times a floating kidney. He believed in the abdominal bandage after pregnancy, and it was important in preventing these troubles.

DR. GEORGE J. PRESTON thought that all those who saw many cases of neurasthenia would be interested in this paper. It is a question how far the nervous condition affects the distension and how far it should be treated. In many cases that he had treated he had noticed diminished knee jerk.

DR. E. M. SCHAEFFER would support what he said about the corset and absence of exercise. He thought the use of the tight corset was potent to do much harm, and exercise would correct many troubles, and he approved of the bicycle in certain cases.

DR. ROBERT L. RANDOLPH read a paper entitled "Some Suggestions as to the Treatment of Wounds Involving at the same time the Lens and the Ciliary Region," and reported two cases in which he did not find it necessary to remove the eye, but took out the lens and one case got well, with some vision.

DR. HARRY FRIEDENWALD read a paper on "Removal of Particles of Steel from the Interior of the Eye with a Magnet," in which he related several cases successfully treated, showed the steel and the magnet, and explained the manner of using it.

DR. WILLIAM H. WELCH then made some

REMARKS ON THE DIAGNOSIS AND TREATMENT OF DIPHTHERIA.

The study of the antitoxins is the order of the day. The evidence that the Klebs-Löffler bacillus is the cause of diphtheria is very conclusive; no one can doubt this now. It is constantly present in every case of genuine diphtheria; isolated in pure culture it is productive of the disease and the disease can be produced experimentally in no other way. We are familiar with the toxic principle of this disease and we know its physiologic but not its chemic properties. This substance has been found in the bodies of those dead from diphtheria. There is probably no organism the study of which has led to such practical results as this. It gives us the diagnosis, prognosis and treatment. The diagnosis can be made with certainty by the demonstration of the specific bacillus. There are many diseases which are like diphtheria in their clinical history, such as membranous croup. Then there are the cases where the disease would not have been recognized on account of the slight symptoms if the bacillus had not revealed the diagnosis, and these light cases are the very ones which spread the disease. Then our increased knowledge of this disease allows us better to understand its symptoms and the question whether it is a local or constitutional disease is cleared up. It develops locally; the bacilli are attached to the membrane in the pillars of the fauces and here the toxins find their way into the body, causing the constitutional symptoms. The prognosis hinges to some extent on the demonstration of associated bacteria. Other bacteria may be associated with the Löffler bacillus in the throat; thus the presence of streptococcus gives an unfavorable prognosis and offer fewer chances for successful treatment. An accurate diagnosis is not only important in order to carry out the treatment, but makes it possible to take sanitary precautions. Of those who are exposed to diphtheria not a small number receive the bacillus without getting the disease, especially in crowded districts. The organism has been found in as many as 50 per cent. of the persons exposed whether they took the disease or not, which shows that there must be a susceptibility, which is a very important point. This susceptibility may be local, that is, susceptibility exists for the local effects and not for the toxins, or the reverse may be true; the local effects may be slight and the general systemic poisoning very severe and in some cases there is no susceptibility at all. Some cases show a thick membrane and little poisoning, and *vice versa*. The method of making the bacteriologic diagnosis is a simple one. As little of the membrane as possible is taken, or some of the secretion from the posterior pillar of the fauces near the larynx. First, cover-slip preparations are made. If it is a case of genuine diphtheria the diagnosis can be made in about 50 per cent. of all cases with no further ex-