and reduced. The head showed a considerable excavation in its center, which was utilized as a recipient for the diaphyseal end. Thus wiring could be dispensed with. No reaction following, the left elbow was corrected five days afterward. The skiagraph represented backward dislocation of the elbow and fracture of the internal epicondyle. There were also indications of much proliferating bone tissue. Reduction of the backward dislocation could not be accomplished before liberating a number of ligamentous fragments. No wiring was necessary after reposition was accomplished. Another skiagraph showed the condition of the joints five weeks after operation. The patient can use both of his arms now, the result being nearly perfect.

Remarks.—In this case no Roentgen examination was made until three months after the accident. If it had been done the treatment would, of course, have been entirely different.

CASE 8.—The skiagraph showed malunion of fracture of humerus five weeks after the injury, which was supposed to have been a dislocation. The head is turned outward, a portion of the articular surfaces being united with the diaphyseal fragment by callus formation. Division was performed by chisel. The result is good.

CASE 9 was one of non-united fracture of radius and ulna, the fragments overlapping six weeks after injury (Fig. 5). Recovery ensued after the fragments had been severed by chisel and wired.

CASE 10.—The skiagraph showed overlapping fragments of the femur of a child of 2 years. Shortening amounted to two inches. Four weeks after the injury the fragments could still be severed by blunt force, the directions of the forcible motions being indicated by the skiagraph.

CASE 11.—A boy, aged 10, sustained a fracture of the lower end of the femur. The lower fragment, including the patella, was displaced backward and became united to the posterior surface of the femur. The shortening four months after the accident amounted to three inches. The patient is now looking forward to an operation.

CASE 12.—Boy, aged 12, sustained a fracture at the upper third of the femur, followed by displacement one-half year ago. It was corrected at the time, the fragments being brought into exact apposition. An effort was made to hold the fragments together by an aluminum tube, which was inserted into the medullary canals of the two fragments (Fig. 6), but the upper half of the tube became loose, the fragments were displaced and greatly irritated so that inflammatory processes followed. For a year a fistula was discharging seropus at the inner margin of the thigh. The nature of the deformity, as well as that of the fistula, was not disclosed until I resorted to the Roentgen method, which showed the overlapping of the bones and the displaced aluminum tube.

MALIGNANT DEGENERATION OF WARTS AND MOLES.*

EDMUND A. BABLER, M.D.

Associate Surgeon, St. Louis Skin and Cancer Hospital; Assistant in Surgery, Washington University.
ST. LOUIS.

The frequency with which warts and moles suddenly increase in size and show other evidence of malignant change does not seem to be appreciated by either the medical profession or the laity. My attention has recently been attracted to this important subject by a series of very impressive cases observed at the St. Louis Skin and Cancer Hospital.

As early as 1860, Collins called attention to the peculiar changes occasionally observed in warts and moles; they suddenly increase in size, soften and appear to ulcerate. Virchow and Billroth were, perhaps, the first to describe cases in which a wart showed sarcomatous change; they designated the pathologic condition plexi-

form sarcoma, and expressed the opinion that warts and moles not infrequently undergo sarcomatous degeneration. In 1892, Unna contended that many, if not most, of these warts and moles are epitheliomata rather than sarcomata, thus concurring with McCarthy, who, in 1880, reported a very interesting case which macroscopically and microscopically corresponded with the sarcomatous growths described by Billroth and Virchow. McCarthy felt confident that the cord-like masses of cells were not of connective tissue origin, but were some form of rapidly growing epithelial structure. Keen has found that growths which arise from warts proper are generally epithelial carcinomata. Several of Keen's cases of malignant disease originating in moles were undoubtedly sarcomata.

In 1903 Eve reviewed the records of the London Hospital for the past twenty years and found that twenty-six of the thirty-three cases of melanosarcoma situated in the skin probably originated in pigmented moles. Eve was convinced that the tendency to the formation of secondary growths in the liver does not appear to be so great in the melanosarcoma of the skin as in the case of primary melanosis of the eye, and that the excretion of melanin in the urine is not so frequent. Wilson and Kalteyer reported a case of melanosarcoma originating in a mole on the forehead, and collected fifty of these cases, in 69 per cent. of which the growth had its origin in a mole or nevus, thereby pointing out the importance of removing warts, moles and nevi as soon as possible lest they become malignant.

It is true that a wart or a mole may never become malignant, but the literature, as well as the cases studied by me at the St. Louis Skin and Cancer Hospital, will convince the most skeptical that an apparently harmless wart or mole may suddenly begin to grow, and often-times quite rapidly, destroying the life of the unfortunate individual.

In some of our cases the more or less constant irritation to which the mole or wart was subjected seemed to be the exciting factor. In one instance the patient attributed the malignant change to the bite of a woodtick. Several of our cases of malignant disease had their origin in a mole which had frequently been accidentally cut by a barber. Most of our cases originated in warts. wart.

It must be admitted that the general practitioner is not always to blame for the delay of excising these apparently harmless blemishes. I have recently observed several cases in which the patient refused to have the growth removed. In our report will be found an instance in which a patient with a mole on his left temple refused operation. He is now eking out a most pitiful existence; the left eye and practically the entire left half of his face have been destroyed by the disease.

The plates submitted with this report emphasize the fact that warts and moles should be removed as soon as possible. I fully concur with Keen, who has contended that the moment warts and moles begin to increase in size they are almost invariably already malignant growths and should be treated as such. Just here I would like to protest against the practice of cauterizing warts on the face; they should be excised.

Cases 1, 2, 6, 8, 9, 11, 12, 16, 17 and 18 occurred in the service of Dr. Lutz, Case 7 in that of Dr. N. B. Carson, and Cases 3, 4, 5, 10, 13, 14 and 15 in the service of Dr. H. G. Mudd. It is a pleasure to acknowledge my indebtedness to Drs. Lutz, Carson and Mudd for the courtesies extended. I am also greatly indebted to Dr.

* From the St. Louis Skin and Cancer Hospital.

Guthrie McConnell for the pathologic reports, and to Dr. Mook and Dr. Wells for the excellent photographs.

REPORT OF CASES.

CASE 1.—History.—Mr. W. E. W., aged 63, was admitted Nov. 19, 1907. Fifteen years ago he noticed a small seed-wart growing on the back of his left hand, but, since it did not cause him any discomfort or did not grow, he gave it only passing consideration for nine or ten years, when seven or eight small, brownish, warty-like growths appeared on the back of his right hand and a scaly, roughened area on the right side of his nose. At this time the wart on the left hand began to grow quite rapidly and he trimmed it frequently. Three years ago the growth on his left hand was the size of a walnut and divided into two equal portions; he was advised to have it removed, but refused. Six months ago it became painful and discharged an offensive, yellowish green fluid. Arsenic was applied and reduced the size of the growth very much, but within a very few weeks the growth had attained its former size. Arsenic was again applied with a similar result. Recently the growth on his left hand has been growing very rapidly and a hard, flat, ulcerated growth has appeared on his left cheek. His left arm is practically useless; he can not flex or extend it very freely.

Fig. 1.—Epitheliomatous degeneration of a wart on the hand.

Examination.—On the dorsal surface of the left hand was a large, cauliflower-like growth covered with a yellowish, greenish, offensive discharge; the lower third of the forearm was swollen; the flexor muscles were markedly contracted, limiting very materially the movements of the forearm; epitrochlear gland not enlarged. On the dorsal surface of the right hand were eight brownish, firm nodules (Fig. 1). On the left malar eminence was a firm, flattened mass, about one inch in length and one-half inch in width, presenting the appearance of an epithelioma (Fig. 1). On the right side of the nose was a scaly-like indurated area as large as a silver dime.

Operation.—The left forearm was removed by disarticulation at the elbow. The rubber tissue drain was removed on the third day. Union was complete on the fourteenth day. The growth on the left malar eminence was also excised. Union was primary, but about three weeks later a small nodule was noted in the upper, outer margin of the line of incision. X-ray exposures have been given each week since the latter finding.

Microscopic Examination.—The specimen from the hand contained many distinct nests of cells of various sizes. The nests were separated by connective tissue which in places was

very markedly increased. The nuclei of the cells varied greatly in size, shape and staining qualities, the large round faintly staining nuclei predominating. Many epithelial pearls were present, with a large amount of keratin formation. In places, near the periphery of the specimen, there was considerable edema. The specimen from the cheek showed numerous nests of cells, the nests being smaller in size than those of the specimen from the hand. The connective tissue was greatly increased, with considerable congestion throughout the specimen. Lymphocytic infiltration was quite marked. Pathologic diagnosis: epithelioma of hand and cheek.

CASE 2.—History.—Mr. J. K., aged 72, was admitted Dec. 17, 1907. Ever since he can remember he has had a mole on the left side of face near the external canthus of eye; six years ago the barber accidentally cut it, but it healed within a short time. A few months thereafter a small scab appeared on the mole and he picked it off with his finger nail; another scab appeared and was removed; after which time the mole became an ulcerating growth; all attempts to heal it were futile. He consulted many prominent surgeons, but was refused an operation. Pastes, salves, powders, x-ray, and other applications were not of any benefit; the disease continued

Fig. 2.—Malignant degeneration of mole on temple.

unabated. During the past six months the destructive process made very rapid progress; patient lost seventy pounds and the pain became of an agonizing character. Morphin, in very large doses, is necessary to secure relief.

Examination.—The left eye and greater portion of left side of face had been destroyed.

Treatment.—Palliative.

Pathologic Diagnosis.—Rodent ulcer.

CASE 3.—History.—Mrs. J. E., aged 54, was admitted Jan. 22, 1906. When thirty years old she noticed a small mole growing on her left temple near the hair margin. She frequently irritated the mole while combing her hair; sometimes the comb scratched off the top so that the mole would bleed. About four years ago the mole began to grow and develop into a malignant ulcer. The destructive process extended gradually to the left ear, which latter it eventually destroyed. A portion of the scalp and left temporal muscles have been destroyed. At present she can not hear very well with left ear; pain is quite severe. At times the pain extends to the orbit and to the parietal region. There is a constant ringing noise in left ear.

Examination.—A large, granular, ulcerated area about four

inches in diameter, the center of which was the middle ear, occupied the left temporal region; the left external ear was destroyed (Fig. 2), also the entire petrous portion of the temporal bone; the dura was in view.

Operation.—Parts curetted and cauterized. Temporary benefit; involvement of brain; death.

Microscopic Examination.—A section, including a portion of the skin of the ear, as well as the ulcerating surface, was examined. The squamous epithelium of the skin showed little change till the immediate neighborhood of the ulcer was reached. The epithelium sent numerous broad projections into the deeper tissues. There was, however, no attempt toward breaking through and infiltrating. The surface epithelium beyond this point became rapidly thinner and disappeared. The portion at the ulcerated end was composed of irregular nests of cells containing long oval vesicular nuclei that stained deeply. No intercellular fibers were present. The cells composing the tumor resembled those of the germinating layer and did not take on any of the changes seen in

Fig. 3.—Malignant degeneration of wart on lower eyelid.

squamous epithelioma. There was also considerable connective tissue hyperplasia at the edge of the ulcer. Round-cell infiltration quite marked.

Pathologic Diagnosis.—Carcinoma (rodent ulcer).

CASE 4.—History.—Mrs. S. O'M., aged 74, was admitted Feb. 25, 1905. At fifty years of age she noticed a small wart on her right lower eyelid. Five years ago the wart began to grow and soon became red and ulcerated and in two years the destructive process involved the entire right half of the lower lid and attacked the eye. One year later she had the lid and eye removed, but within a short time the condition was as bad as before operation. Pain has become a very prominent and distressing symptom. Pastes, salves and powders have been applied without benefit.

Examination.—The right orbital cavity was occupied by a cauliflower-like growth from which exuded an offensive, purulent discharge. Patient was emaciated, pale and weak.

Operation.—The orbital cavity was thoroughly cleaned out. Within a very short time evidences of recurrence were noted (Fig. 3). The disease extended to the brain; patient died a short time ago.

Pathologic Diagnosis.—Epithelioma of orbit and brain.

CASE 5.—History.—Mr. P. G., aged 70, was admitted Aug. 25, 1905. Six years ago he noticed a small wart-like nodule on the left side of his nose. Six months ago it began to grow, extending to cheek. It became lobulated and resembled a bunch of grapes. At present it is growing at a rapid rate.

Examination.—On left side of nose and cheek was a red, lobulated growth about as large as half of a walnut, presenting the appearance of an epithelioma.

Operation.—Growth was excised. Union was primary. Patient has not been seen since May, 1906. No recurrence at that time.

Pathologic Diagnosis.—Epithelioma of nose and cheek.

CASE 6.—History.—Mr. J. L. T., aged 73, was admitted July 11, 1906. He had a small mole on the left side of his nose ever since he can remember. In 1889 a barber accidentally cut off the top of the mole; the wound healed, but about a year

Fig. 4.—Epitheliomatous degeneration of a mole on the neck.

ago a scab appeared and he picked it off. Within a short time another scab appeared and he removed it; this process was repeated at frequent intervals until nine months ago, when the mole became a destructive ulcer which has grown rapidly during the past two months.

Examination.—On the left side of the neck near the angle of the jaw was an indurated ulcer, one and one-half inches in length and one-half of an inch in width, presenting the usual appearance of a rodent ulcer. No palpable glandular involvement. Growth was excised. Primary union.

Microscopic examination showed rodent ulcer.

CASE 7.—History.—Mr. J. P. C., aged 56, was admitted June 26, 1906. For many years he has had a small mole on the right side of his face, but it did not cause him any inconvenience until in 1889, at which time a barber cut it while shaving him. The wound healed within a short time, but a few months later a barber cut it again, after which it did not

heal satisfactorily. A few years ago he noticed that the mole was ulcerated and increasing in size. Pain has not been severe in character.

Examination.—On the right side of the face, over the molar eminence, was situated an irregular, depressed ulcer, worm-eaten in appearance, about as large as a silver half-dollar. It presented the appearance of a rodent ulcer. No palpable glandular enlargement.

Operation.—Growth excised. Primary union. Recovery.

Microscopic Examination.—The neoplastic cells were distributed throughout the specimen in two different arrangements. In places there were quite large, irregular nests of cells containing oval or round nuclei that stained fairly deeply. The peripheral layer of cells was arranged in a distinctly columnar fashion. There was no restricting membrane, the cells lying in direct contact with the adjacent tissue. In other places the cells had a more or less regular arrangement into narrow strands surrounding distinct acini. In many of these openings were found blood cells and granular matter. Some of the acini contained a somewhat granular mass which stained slightly with eosin. Round-cell infiltration was quite marked. Some muscle fibers showed hyaline change. Numerous mitotic figures.

CASE 8.—History.—Mr. McC., aged 74, was admitted Jan. 15, 1907. Ever since he can remember he has had a small mole or birthmark on the left side of his neck, just below the tip of the mastoid process of the temporal bone. Sixteen years ago a woodtick bit him on the mole. He was unable (he thinks) to remove all of the tick, and is under the impression that his present condition is the result of this. Two or three months after he was bitten he noticed that the mole was very tender to touch, and that it was increasing in size. Various preparations were applied; a small ulcer appeared on the mole, which showed a tendency to increase in size and refused to heal kindly. He finally succeeded, however, in healing the ulcer. A few years later he accidentally contused the left side of his neck against the limb of a peach tree; the parts became very sore and ulcerated. The tissues immediately surrounding the mole became indurated and of increased size. He consulted various "quacks" and spent all of his money, but the ulcer continued to get larger and larger. As time went on the growth caused more or less constant and severe pain; pain was especially severe at night. Recently the growth has been increasing more rapidly and has destroyed a large portion of the left side of the neck, posterior to the sternomastoid muscle. He has lost considerable weight.

Examination.—A large, granulated, depressed ulcer with irregular, firm margins occupied the left posterior side of the neck (Fig. 4). The muscle was exposed in the central portion of the ulcer; portions of the muscle showed characteristic malignant change. Portions of the margins of the ulcer were undermined. Cervical (left) glands palpably enlarged. The sternomastoid muscle (left) very firm and infiltrated at its upper portion. General condition of patient quite good.

Treatment.—I gave trypsin injections for a considerable time, but with only temporary benefit; the severe pain became lessened to such an extent that the patient could sleep the entire night. Within a short time, however, the pain returned, but was never as severe as previously. The trypsin treatment did not change the condition of the growth, except that it became clean. Owing to illness in patient's family, he was called home about three months after admittance to hospital. Not traced.

Microscopic Examination.—One-half of the specimen consisted of a mass of quite dense connective tissue covered by a layer of surface epithelium. The other half was made up of a more open connective tissue framework, scattered throughout which were numerous small, irregular nests of cells. These nests were not limited by any membrane and were separated from each other by greater or less amounts of adult connective tissue. There was a very general widespread round-cell infiltration.

Pathologic Diagnosis.—Squamous epithelioma.

CASE 9.—History.—Mr. G. T. W., aged 54, was admitted Oct. 1, 1906. About two or three years ago he noticed a small warty growth on his lower lip. In October, 1905, it began to grow

rapidly, and the physician whom he consulted diagnosed it cancer. Acids, pastes and plasters were applied without benefit. The disease continued unabated. He lost weight; appetite poor; pain more or less constant and quite severe.

Examination.—A large cauliflower-like growth occupied practically the entire lower lip and a part of the chin and inferior maxilla in median line. The mouth was apparently free from involvement. The growth was very firm, except in the ulcerated portions. Patient presented a somewhat cachectic appearance.

Operation.—The lower lip in its entirety and the submental and submaxillary glands were removed, and the diseased bone was thoroughly curetted. Superficial tissues of face were approximated and a new lip was formed. Patient developed a pneumonia and died a few days after the operation.

Microscopic Examination.—The superficial epithelium was very much thickened and the papillæ were large. Lying deeper within the tissues were irregular nests of cells containing large round bubble-like nuclei. The cells in the periphery of these nests showed quite a well-marked columnar arrangement. In the center of many of the nests the cells had become considerably flattened and had undergone keratin formation with loss of the nuclei. Round-cell infiltration was very well marked beneath the epidermis and between the cell nests. Very few mitotic figures were found. The submaxillary gland showed slight inflammatory changes.

CASE 10.—History.—Mr. L. W., aged 58, was admitted Aug. 23, 1906. About seventeen years ago he noticed a small warty-like growth on right half of his lower lip. For years after its appearance it began to grow at a rapid rate and he had it excised. Three years later it returned and in 1901 he had it cauterized. The growth disappeared and did not return until three years later; it then appeared in the form of a small, hard ulcer, which latter soon became of a rapidly destructive character. He tried various so-called "cures," but the disease continued unabated. No loss of weight; appetite good.

Examination.—The entire right half of lower lip had been destroyed. A hard crate-like, ulcerating mass presented itself in this area; the lower teeth, normally covered by the right half of the lower lip, were exposed. The submental glands, as well as the submaxillary glands, were not palpably enlarged. Floor of mouth was apparently free from involvement.

Operation.—Radical excision. Patient recovered. Union was primary. No recurrence up to present time.

Microscopic Examination.—The greater part of the specimen was composed of irregular masses of cells containing rather small round or oval vesicular nuclei. These masses had no limiting membrane and ramified through the tissue in the form of a coarse network with more or less connective tissue in between. In a number of places the tumor cells had a distinct adenomatous arrangement. Narrow strands of about two cells in thickness formed a network enclosing spaces that appeared to be filled with a secretion. There were several areas in which the central portion of the cell masses had undergone a marked degeneration.

CASE 11.—History.—Mrs. M. C., aged 56, was admitted Oct. 21, 1907. Many years ago she noticed that she had a mole on the right side of her nose, but, since it did not cause her any inconvenience, she did not pay any attention to it until five or six years ago, at which time it began to increase in size. Pastes, salves and other "frauds" were applied without benefit. The growth rapidly increased in size. Pain was not severe or constant.

Examination.—On the right side of the nose, at about the center, was a brownish, nodular growth, one inch in length and one-half inch in width; it felt tense and firm and was in the skin.

Operation.—Excision under local anesthesia. Primary union. Subsequent Roentgen ray treatment.

Microscopic Examination.—The superficial epithelium was considerably thickened and showed a very great keratin degeneration. There was some parakeratosis.

CASE 12.—History.—Mr. W. F. R., aged 53, was admitted March 6, 1907. Five years ago he noticed a small, white, wart-like growth on his lower lip near its center. He pulled it out, but within a few weeks it returned. The process was fre-

quently repeated. The substance removed was always of the same consistency, spongy-like and closely resembled a wart. Five or six months after its first appearance he consulted a so-called "cancer doctor," who attempted to treat it with his pastes, etc., but without success. The application employed seemed to accelerate the development of the growth. One year ago the growth involved the entire right half of the lip; he consulted a surgeon who excised the lower lip. Within six months the growth returned (?) and rapidly attained the size of a small closed fist. Another surgeon found the inferior maxilla also involved. Three weeks ago the growth on lip and a portion of the inferior maxilla were removed. At present there are evidences of recurrence. Loss of sixty pounds in weight during the past year.

Examination.—Mouth deformed, due to previous operation. There was constant flow of saliva from mouth, at right angle of the very narrowly contracted mouth was a small nodular growth about as large as a pigeon's egg. Glands not palpably enlarged. Floor of mouth not involved. Patient could scarcely open his mouth.

Treatment.—He had been subjected to Roentgen rays, but without benefit. Injections of trypsin were tried, but so many abscesses followed its use that they were discontinued. The lower lip was freed from the inferior maxilla and the growth was excised under chloroform anesthesia. Patient has not been heard from since he left the hospital. Probable recurrence.

CASE 13.—History.—Mr. J. T., aged 75, was admitted Sept. 4, 1907. About three years ago he noticed a small wart on the left side of his nose, but, since it did not seem to grow and did not cause him any pain, he let it alone; within a very short time it became an ulcer and grew rapidly. Painful at times.

Examination.—On the left side of nose was a hard movable mass about as large as half of a walnut. The edges of the mass were of firmer consistency than its central portion. The growth presented the appearances of an epithelioma.

Treatment.—Excision under local anesthesia. Primary union.

Microscopic Examination.—The specimen was composed of many small cell nests separated from each other by a slight amount of connective tissue. In many places the nests appeared to have fused, thus forming masses with scalloped edges, resembling rosettes. In the nests the peripheral cells were arranged in a quite regular columnar manner, the central cells being massed together irregularly. Many of the cells were undergoing division. Some round-cell infiltration in the connective tissue.

CASE 14.—History.—Mr. L. B. L., aged 78, was admitted May 12, 1906. Eight years ago he noticed a small wart-like growth on his lower right eyelid, but it did not cause him any discomfort or did not seem to increase in size until four years later, at which time a small scab appeared. He pulled this off and the wart bled freely. Within a short time the wart began to ulcerate and discharge a greenish, offensive pus, and the parts became swollen and more or less painful. The destructive process extended until the entire lower lid was destroyed; the upper lid was so swollen and edematous that he could not see out of his right eye. No loss of flesh.

Examination.—The right lower lid was practically destroyed. The tissues adjacent to the base of the lower lid were indurated and infiltrated by a growth presenting the appearance of an epithelioma. The upper lid was edematous and obstructed sight of right eye. Eyesight not destroyed. Lymph glands not enlarged.

Operation.—The lower lid and tissues at base of lid were excised. Within a very short time the eye showed evidences of being affected by the malignant process. The right orbital cavity was cleared out. Shortly thereafter the patient died.

Microscopic Examination of Eyelid and Contents of Orbit.—The first specimen was completely surrounded by squamous epithelium which on one side was quite thin and had almost no papillae. The epithelium increased gradually and extensions dipped deep down into the corium, forming anastomosing strands. The cells varied in size, and many contained numerous minute granules almost black in color. In a few places the cells were flattened and arranged concentrically, but had

not undergone a keratin transformation. No mitoses. The second specimen showed large masses of adult connective tissue separating and surrounding irregular nests of cells. These cells had quite large spindle-shaped nuclei, vesicular in character. There was no limiting membrane, the cells lying free within the tissues. Numerous mitotic figures present. Round-cell infiltration quite well marked. Vessels congested; some interstitial hemorrhage present.

Pathologic Diagnosis.—Rodent ulcer.

CASE 15.—History.—Mr. C. W. N., aged 59, was admitted June 8, 1906. Was never ill until two years ago, when he noticed a small wart growing on penis near urethral opening. A few months later he consulted a physician, who burned off the wart. Contrary to expectations, the wound would not heal. Six months ago he was operated on, a portion of the penis being removed. He has become steadily worse and has lost considerable weight.

Examination.—Patient anemic and looked badly. The entire penis had been destroyed and was replaced by a large, foul sloughing ulcer. The testicles were swollen and the scrotum was edematous and ulcerated in places. The inguinal glands were swollen, tender and sloughing. The condition was pitiful and exemplified the distressing result of the physicians' failure to do a radical operation instead of a partial one.

Treatment.—Patient's condition did not warrant operation.

Pathologic Diagnosis.—Epithelioma of penis.

CASE 16.—History.—Mrs. R., aged 70, was admitted Dec. 10, 1907. Sixteen years ago she noticed a small, white, seed-wart on her upper (right) eyelid. It seemed to remain unchanged in size for twelve years, when it began to grow and shift its site from the outer third to near the inner canthus of the eye. Caustics, pastes, salves, etc., had been applied without benefit. Pain was not a prominent symptom. The growth increased very rapidly in size, a small nodule appearing near the margin of the lid.

Examination.—About half an inch above the inner canthus of right eye was a nodular, slightly ulcerating mass as large as a five-cent piece. The growth presented the usual characteristics of a rodent ulcer. On the border of the lid, near the inner canthus, was a small, hard whitish nodule about as large as a pin-head.

Operation.—Excision. Wound healed by granulation. Exposure to Roentgen rays twice a week. Recovery.

Microscopic Examination.—Specimen showed numerous nests of cells varying in size and shape. These nests were distinctly separated by connective tissue. The nuclei of the cells varied greatly in shape and staining qualities; the oval, deeply staining ones predominating. There was a very large amount of round cell infiltration and of congestion. No attempt at pearl formation.

CASE 17.—History.—Mr. J. J. D., aged 59, was admitted April 28, 1906. About two years ago he accidentally noticed that he had a small wart-like growth on the back of his left hand; it did not cause him any pain or discomfort until about six months later, when it began to grow. Various acids, pastes and other drugs were applied without success. Five months ago the growth became ulcerated and began to grow rapidly. Two weeks before admission he had a physician remove the growth, but when the sutures were removed, a week after operation, the physician found that the parts had not healed; the wound was gaping.

Examination.—On the dorsum of the left hand, between the base of thumb and index finger, was an irregular, exuberant ulcer, the size of a silver quarter. Axillary and epitrochlear glands not palpably enlarged.

Treatment.—X-ray exposures were given twice a week for some months. The parts healed and the patient discontinued his visits to the clinic.

CASE 18.—History.—Mr. G. S., aged 62, was admitted Oct. 14, 1907. Had a wart on his lower lip for twenty-five years, but it did not grow until two months ago. Lower lip had become swollen and the growth was the size of a filbert.

Examination showed patient well nourished and in apparent good health, except for the growth on and swollen condition of his lower lip. Near the center of the free margin of the lip was a scaly growth presenting the characteristics of an

epithelioma. The swollen condition of the lip did not appear to be due to the epitheliomatous growth only. The submental and the submaxillary glands were not palpably enlarged.

Having been so frequently impressed with the fact that a wart may suddenly show malignant changes, and appreciating the importance of early excision, the patient was admitted to have the growth removed. One day later, the entire lower lip was excised and the submental glands removed. The silkworm gut sutures were removed on the fifth day. Union was primary.

Pathologic Report.—The epithelium suddenly changed from the normal to an irregular layer of great thickness. Long finger-like projections of cells extended deep into underlying tissues. Throughout the greater part of the growth, the cell nests penetrated down into the muscular tissue and brought about an atrophy of the muscle fibers. Along the advancing periphery of the new growth there was a very marked infiltration of lymphocytes and plasma cells. The cells found in the nests were large and polygonal, and contained a large round or oval vesicular nucleus. Many of them showed varying degrees of keratin formation, and typical epithelial pearls were very numerous. The cells on the surface of the growth showed marked keratin formation and parakeratosis. On the mucous side of the specimen was found a small salivary gland. On the skin side were seen numerous hair follicles, sebaceous and sweat glands. Lying deeper within the muscular tissues were two bodies, circular in shape, with a surrounding rim of homogeneous substance that stained with eosin. This ring enclosed a space in which were found several round and oval bodies that contained cells and nuclei enclosed within a definite membrane. In between these bodies was a small amount of granular pink-staining material. External to the capsule was a slight amount of round-cell infiltration. These masses lay embedded in the muscular tissue, and the adjacent muscle fibers had been flattened in varying degrees. The appearance was that of an encapsulated trichina. These bodies lay quite close to quite a large artery.

CONCLUSIONS.

1. The laity should be impressed with the danger of permitting warts and moles to grow, and should be urged to have them excised at once.

2. The early excision of apparently harmless warts and moles will save days of suffering; in many instances, the life of the individual will be saved thereby.

3. The moment that warts and moles begin to grow they are almost invariably already malignant growths and should be treated as such.

4826 Delmar Boulevard.

STATIC ELECTRICITY.

ITS PHYSICAL PROPERTIES, MODALITIES, PHYSIOLOGIC EFFECTS AND THERAPEUTIC INDICATIONS.

WILLIAM BENHAM SNOW, M.D.
NEW YORK CITY.

To standardize the therapeutics of static electricity properly a consideration of the type of apparatus and modalities employed is a question of first importance. There are three types of static machine at present in the market—the Holtz, Toepler-Holtz and the Wimshurst—which produce currents of practically the same quality but varying materially, other things being equal, in the potential as measured by spark-gap and shrdlu odhmc discharge when high resistance is placed in the circuit.

My experience and familiarity with static machines leads me to the conclusion that the Holtz machine, having from eight to sixteen revolving plates and being provided with an exciter, is the best type of machine for medical purposes. My reasons for this conclusion are: 1, During the administration of the Morton wave current, the modality of most importance in static thera-

peutics, this type of machine will produce, other things being equal, the longer spark-gap and discharge with the greatest degree of regularity; 2, this type of machine is more easily kept in repair; 3, when having the suitable number of revolving plates, sufficient current is evolved when running at a low rate of speed (300 to 400 per minute) without producing alternating oscillatory currents as is sure to occur with high speed machines.

For the evolution of currents of high frequency, in connection with the static resonator, a high-speed apparatus having a smaller number of revolving plates, in which a unidirectional current is not desired, the same objection does not obtain. The idea of inducing a general purpose current by employing a smaller number of revolving plates at high speed is not practical or scientific, for the quality of the current, as to proportion of potential and amperage, is not under control as it is when derived from a number of revolving plates sufficient to produce the current at a lower rate of speed. With high-speed machines the spark-length is deficient and the current is oscillatory or alternating in character.

At the present time the static machines of the Holtz type having sixteen revolving plates 30 to 36 inches in diameter and constructed with a high degree of insulation to prevent escape of the current to and through the case, and provided with a means of controlling amperage of the current without reducing the potential as measured by the length of the spark gap when administering the wave current, is the type of static machine most cordially to be commended. For most purposes, and for most patients when the employment of the x-ray and high-frequency currents are not taken into consideration, a static machine having eight revolving plates of the above type will meet most indications.

CLASSIFICATION OF STATIC MODALITIES.

Numerous static modalities are employed therapeutically, of which the following may be considered in order of excellence: 1, The static wave current; 2, the static spark; 3, the direct vacuum tube current administered in the same manner as the static wave current; 4, the static brush discharge; 5, the static induced current; 6, static insulation; 7, the static spray. In this classification we are not considering the high-frequency currents—auto-condensation, auto-conduction and the high-frequency effleuve—also produced by the static machine of a quality similar to the high-frequency currents derived from other sources.

The properties of the static current, which render it unique, are: 1, intensity of diffusion; 2, constant or unidirectional; 3, high potential, relative to small amperage; 4, ease of control of current as to periodicity or frequency, by regulation of speed and spark-gap, regulation of amperage and voltage by the same means, and the additional regulation of ground connection by placing in the circuit a variable resistance; 5, it is the only apparatus with which the current can be satisfactorily administered from one side of the machine with the other side grounded and the patient insulated, thereby producing unique physical and physiological effects not derived from other apparatus.

DIFFUSION OF THE STATIC CURRENT.

The remarkable degree of diffusion of the static current is evidenced by the passage of the current through materials which are non-conductors of other currents, as its passage through the length of a long platform with the discharge of a spark several inches in length, or the passage of the current through wooden electrodes—a