of her bowel movements. Almost wholly disbarred from going into society, her misery during this period can readily be imagined.

On November 27, 1900, I operated, assisted by Drs. Charlton Wallace and J. W. D. Maury, her family physician, Dr. Stuart Close being present. The technic employed was one which in its simplicity and application seems hard to improve on. It was devised by Dr. Ristine of Knoxville, Tenn., more than twelve years ago and its technic was published by him some years afterward in the *American Journal of Obstetrics and Diseases of Women*. The details, accompanied by a rough sketch, which may serve better than words to make clear one or two steps, are as follows: The rectum being empty and cleansed, is packed with gauze. A horseshoe-shaped flap of mucous membrane, with its toe upward, is next outlined. The incision for this begins on either side of the vagina at the ends of the torn and retracted sphincter ani. The cut ascends on either side and meets at the toe, placed, in a case as bad as this, just as high as possible behind the cervix uteri—almost touching the cervix—so as to waste no length of flap that it is possible to secure. With a tear only dividing the sphincter, and not the rectum, obviously the horseshoe cut need not ascend so high.

The apron of the flap thus turned down will have its sound, i. e., its mucous membrane surface toward the feces and its raw surface forward. It will be dissected down as far as the upper end of the tear permits. If of proper length its lower or free ends should now hang as low as the anal level behind it. Any excess of length of this apron is a protection, for the time, to the dissected flesh in front against accidental infection, and if it should not finally shrivel, it can be snipped away.

Search is next made for the torn sphincter, the ends of which must be freshened and then brought into mutual apposition by means of tanned or chromicised catgut. The sphincter muscle, even in this case, could still be plainly recognized. By dissecting down and lengthening the horseshoe ends as needed, tissue is at length reached which when seized and pulled on by the forceps can be felt to draw the sphincter on either end. Before apposing these ends we should insert and tighten the continuous suture shown as E and E' in the accompanying sketch. This helps to prevent any "dead space." The curved needle is inserted and brought out but four times in passing this stitch. Following the union of the sphincter the surgeon will probably perform the customary operation for torn perineum.

Before concluding, a moderate section of the sphincter at the rear anal margin should be made (in a case recently torn) to weaken it temporarily—for protection against the tearing out of the important stitches uniting its ends. Perhaps I need not say that in the case under discussion, after twenty-three years of disuse, it was thought unnecessary to weaken the sphincter further in this way. When this is done no stitch is taken; a clot fills the tiny opening made by the slender blade. The bowels are moved the second day and each day thereafter. Electricity and strychnin are of value in aiding the sphincter to recover its function.

I would refer those especially interested to the article by Dr. Charles G. Child, Jr., of this city, in the *Journal of Obstetrics and Gynecology of the British Empire*, July to December, 1905. It will be found that in a few particulars my operation differs from that of Dr. Ristine—the principle, however, is his, as also should be the entire credit of this, one of the very few operations which seem to me perfect in technic.

APPENDECTOMY DURING HERNIOTOMY.

JOHN A. HAWKINS, M.D.

Professor of Rectal Diseases in the Medical Department of the Western University of Pennsylvania; Rectal Surgeon to St. Francis Hospital.
PITTSBURG, PA.

The article by Dr. John A. Wyeth, in *THE JOURNAL*, Nov. 24, 1906, reminded me of two patients on whom I recently operated.

The first patient, a man of 67, had suffered with a large inguinal hernia for many years. On opening the sac I found that it contained the cecum, the adherent appendix and part of the ileum. The appendix, though not inflamed, was removed, and the abdomen closed by the imbrication method of Fowler. Recovery was uneventful.

The second case was that of a woman of 62, who suffered from an inflamed bubonocoele. The sac contained a structure of whose identity I was not altogether certain until I had examined it closely, but which proved to be the appendix. The constricted ring was incised and the hernia repaired by the Bassini method, the appendix being removed in the usual way. There was nothing unusual about the recovery. The appendix was 10 cm. in length, 12 mm. in diameter at the distal end and 4 mm. at its attachment, being club-shaped.

219 Sixth Street.

A CASE OF INFECTION BY STRONGYLOIDES INTESTINALIS.*

J. N. HALL, M.D.

Visiting Physician to the Denver City and County Hospital.
DENVER.

This parasite, associated with Cochin-China diarrhea, has been reported so rarely in the United States that the following case should be recorded. The exact relationship to tropical diarrheas seems not definitely established:

Patient.—A Japanese ship's cook, aged 40, was admitted to my service in the Denver City and County Hospital, Oct. 18, 1906.

History.—We learned that he had visited many ports in Asia and other parts of the world, had been in America seven years, in Colorado three years. He had been sick for ten weeks, although below par for three or four months. He had a slight cough, had probably lost 20 or 30 pounds in weight, and was moderately anemic. He had five to eight stools in the twenty-four hours, mushy in consistence, foul smelling and of clay, slate or dark green color. No blood nor mucus. He complained of constant pain in the abdomen, and in the rectum on defecation. He vomited a few times, especially after eating sweet food.

Examination.—Physical examination of the chest and abdomen was negative. The urine was acid, sp. gr. 1030, a trace of albumin, no sugar, no bile, indican increased. A few small and narrow hyaline and granular casts, occasional blood cells and leucocytes. The examination of the sputum was negative. The blood, two weeks before death, showed: Hemoglobin (Tallquist), 65 per cent.; red cells, 3,200,000; white cells, 4,400. Moderate poikilocytosis and no megaloblasts nor normoblasts were found. A differential count of white cells showed polynuclear neutrophils, 51.5 per cent.; lymphocytes, large and small, 47 per cent.; eosinophiles, .05 per cent. The absence of eosinophilia is to be noted.

Suspecting amebic dysentery, I asked Dr. B. B. Blotz, in charge of the laboratory, to examine the stools. His report is as follows:

"Larvæ of the *Strongyloides intestinalis* were found in great numbers. At least fifteen examinations were made and the parasite was present in each. An attempt was made to obtain the adult by incubating the larvæ for seventy-two hours at

* Reported, with exhibition of specimens, to the Denver City and County Medical Society, No. 6, 1906.