ensuing I discovered that a fine, robust specimen which I had taken from decaying hickory the day previous had its left elytron .25 mm. shorter than its right, this time the difference being much less, but still noticeable. I then examined fifteen other specimens of this species in my collection, with the following result: One with left elytron .25 mm. shorter than right; one with left .20 mm. shorter than right; one with left shorter than right, but the difference hardly appreciable; one (small specimen) with right .20 mm. shorter than left; and one in which the right was so slightly less than the left that the difference could scarcely be seen. The remaining ten showed no appreciable differences in this respect. So of seventeen specimens examined, seven had the elytra unequal in length in a considerable degree, one being especially prominent thereby. And it is noticeable that in five of the seven specimens it was the left elytron that was the shorter, these also being the cases in which the inequality was most prominent. I believe all of my specimens, with one exception, were taken from their cells in the wood, as they are found after having assumed the imago. I have no doubt that if others would examine the specimens of this species in their collections, many more such examples would be found which have been overlooked. It would be interesting to know the result of such examinations. It is probable that the elytra, being organs not of strictly primary value to the insect—elytral invariability in this direction not being absolutely essential when within certain limits—have thus been permitted to vary without the variations being struck out by natural selection. I have not observed this elytral inequality in any other Coleoptera as yet.

A form of this species, which is less robust, I occasionally find; it is slightly narrower in proportion and more delicately marked, but upon sending specimens of the two to Dr. Horn, he informs me that they do not differ appreciably, but are both *oculatus*.

CORRESPONDENCE.

NOTE ON THE HABITAT OF *XYLORYCTES SATYRUS*.

Dear Sir: Mr. W. F. Robinson, of Bridgeton, N. J., sent, Oct. 28, 1884, a specimen of this beetle, with the remark that it is being dug up around the roots of trees there in abundance by his school boys. The collections here contain it not farther to the north than Pennsylvania. Not

being able to find it recorded from N. J., or farther to the north, I should like to hear if its occurrence north of Pa. is known. I may add that Mr. Robinson gave to the collection a very small specimen of *Strategus antaeus* collected by him in Nantucket, Mass., Aug. 24, 1884.

DR. H. A. HAGEN, Cambridge, Mass.

Dear Sir: In reference to your article in the August number, allow me to add as injured by *Pulvinaria innumerabilis*, the following: No. 1, *Acer saccharinum*; 2, *A. rubrum*; 3, *Negundo aceroides*; 4, *Ulmus fulva*; 5, *U. americana*, 6, *Celtis occidentalis*; 7, *Morus rubra*; 8, *Fraxinus sambucifolia*; 9, *Populus balsamifera*; 10, *Juglans cinerea*; 11, *J. nigra*; 12, *Rhus toxicodendron*. *Tilia americana* I have not seen affected here; our cultivated grape (*Vitis*) very seldom, and sparsely, if at all, while our wild grape seems perfectly free from the pest. This insect, better known here as the Maple-slug, has been found by me on all the above, never taking note unless I found it at least six times on the same kind of trees, growing far apart and in quantities as follows: On No. 1, sparsely; 2, soft maple, very full; 3, ash-leaved maple, better known as box-elder, very full, equal to No. 2; 4, 5 and 6 less affected but alike; 7, 8 and 9 sparsely; 10 and 11 least affected, while 12 (being our poison ivy) only when the tree to which this parasitic plant was attached was affected. I had made my observations in Peoria, Tazwell and Livingston Counties, and have noticed in the City of Peoria especially the stone pavements sprinkled with the so-called honey dew in patches here and there, so thick that any passer-by would look up wonderingly unless familiar with the cause. As such articles are of interest to farmers and to growers of shade trees, they will be glad to get such information as experience can give them through some observation. The last visit of this insect in great numbers was about five or six years ago. The question here has frequently been asked with some anxiety: How will they appear next season, etc.? Their natural enemies and wet weather, especially when the rains come frequently during hatching, as well as other causes, are among the agencies which limit their increase.

A. H. MUNDT, Fairbury, Ills.

August	No.	Mailed	Nov. 17,	1884.
Sept.	"	"	Dec. 1,	"
Oct.	"	"	" 17,	"
Nov.	"	"	" 24,	"