The writings of Nothnagel, Osler and Hawkins, and this recent paper by Smithies indicate that hemiplegia is a very rare complication in typhoid fever. But I am inclined to believe, that, while this is true, the complication is less rare than we had formerly supposed; for I have myself seen three well-marked cases, two of which I have reported. One of these (*Medical Record*, Dec. 19, 1903) is included in Dr. Smithies' table; the other (*THE JOURNAL A. M. A.*, May 27, 1905) is not. The third case I have never reported.

The complication deserves more study; and it is to be hoped that physicians will place on record such cases as may come under their observation, so that a more adequate idea may be obtained as to the genesis, character and frequency of this complication.

THEODORE DILLER.

Taxing Private Hospitals.

CLEVELAND, OHIO, Aug. 15, 1907.

To the Editor:—In *THE JOURNAL*, July 27, 1907, the matter of taxing private charitable institutions is discussed. The view there taken is a mistaken one. Public support given private institutions, no matter for what purpose organized, whether religious, charitable or otherwise, is unsafe, as it tends to foster the unjust taxation systems under which we labor. Exactly as I am glad to see half rates withdrawn from ministers of the gospel, so I would be happy to find all land or monopoly values (they amount in last essence to the same thing) taxed on their actual social value and that regardless of what use put to (except in the case of state institutions, where the reason for not taxing is self-evident, clear cut and incapable of misapplication). Human nature being such that men will support the system which is easy on them, i. e., which gives them an advantage, real or imaginary, over their fellows, irrespective of the justice of that system, I would rather not see the situation further muddled by more institutions on the free list. Let there be economic justice, and charity (such as we have now) will become a nightmare of the past.

J. E. TUCKERMAN.

Probilin.

PHILADELPHIA, Aug. 24, 1907.

To the Editor:—The report on Probilin in *THE JOURNAL*, August 24, explains my experience with the formula suggested by Dr. R. W. Wilcox in his article on "The Medical Treatment of Gallstone Disease" (*THE JOURNAL A. M. A.*, Aug. 4, 1906, page 347). Being desirous of carrying out the treatment suggested by Dr. Wilcox, I prescribed the formula of phenolphthalein with acid sodium oleate and salicylic acid. The pharmacist to whom the prescription was submitted informed me that it could not be compounded as written, but that he could dispense the materials massed separately in a capsule. Subsequently the same pharmacist suggested the use of sodium salicylate in place of the salicylic acid, the following modified formula being prescribed:

Sodium oleate	0.10
Sodium salicylate	0.10
Phenolphthalein	0.05
Menthol	0.015
Mix and dispense in 1 capsule.	

The sodium oleate was prepared by mixing the proper proportion of sodium hydrate and oleic acid.

While my experience with the above formula has been limited to four cases, the results obtained seemed to be due to free catharsis resulting in a lessening of the jaundice and the presence of bile in the stools. In no case were stones recovered. Two capsules daily were found to produce active catharsis and dosage was limited to that number.

HENRY F. PAGE.

Organization.—The spirit of the present organization of the medical profession is unique. It is not the labor union spirit of merely personal and material benefit; it is spirit of personal uplifting, of personal education, of personal improvement in all those things which go to make a man a better physician and a better guardian of the sick in his charge.—Editorial in *California State Journal of Medicine*.

Miscellany

Serum Treatment of Plague.—Experience with the serum treatment of plague illustrates clearly the difficulties under which the experimenter labors in producing a curative serum for a bacterial infection of the blood like the plague. The natural process of cure for plague appears to be the destruction of the bacilli which have entered the blood, while the toxins which they produce are destroyed or antidoted by the natural powers of the system. No great amount of antitoxin appears to be formed in the organism which is able by its natural powers to cope with the toxemia only to a limited extent. This explains in a manner the extreme fatality of the plague. The destruction of the plague bacilli in the blood sets free a large amount of toxin which overwhelms the system so that it can not by its unaided powers antidote or eliminate it and a condition known as plague marasmus results. This condition is graphically described by Choksy in his "Report on the Treatment of Plague with Yersin-Roux Serum at the Maratha Hospital During 1905." The plague serums so far produced are bacteriolytic serums with little antitoxic power. They hasten the destruction of the plague bacilli, but leave the system to cope with the toxemia unaided. If administered before the bacilli have multiplied extensively in the blood, they cut short the disease, but if administered at a later period they may intensify the toxemia by increasing the bacteriolysis and really aggravate the condition. Hence the administration of the serum in hospital practice, where a considerable number of cases are admitted somewhat late in the disease, shows but little advantage over the cases treated without the serum; indeed, a larger proportion of cases exhibiting acute marasmus with marked toxemia are observed under the serum treatment than without it. But if the hospital cases are analyzed according to the day of the disease when the treatment was applied a considerable improvement under serum treatment may be demonstrated. Choksy finds that by summing up three series totaling 500 cases there was a general mortality of 53.6 per cent., while the percentage of mortality among those inoculated on the first day, numbering 89 cases, was only 38.2. The mortality increases in a regular ratio (with the exception of a slight drop on the fourth day) to 75 per cent. of those inoculated on the seventh day. The results from serum treatment are better in private practice largely because of the fact that such patients apply for treatment earlier and have better care. The serum treatment not only secures somewhat better results in general, but it secures an earlier convalescence, marked by fewer complications than occur in cases that have not had serum treatment. Choksy instituted a comparison of the two methods of treatment under circumstances well calculated to give a fair representation of the value of each. Of 148 cases, 46 were rejected as unfit for serum treatment, and of the remainder each alternate patient received serum treatment, the others serving as controls. Of the 51 in the control group, 42 died and 9 recovered, while of those treated by serum 37 died and 14 recovered, the mortality being 82.3 per cent. and 72.5 per cent. respectively. On the whole, therefore, there can be no doubt that there was distinct gain under the serum treatment, and that more lives were saved than under the ordinary treatment. The results leave much to be desired, however. This result, it must be remembered, is obtained in a disease that carries away about 84 people, on an average, out of every 100 attacked. Choksy is inclined to believe that, for the present at least, the limit of success is reached. He suggests that the results might be better if the serum were administered intravenously. If a serum could be discovered combining antitoxic and bactericidal properties, more lives could doubtless be saved. Under the present circumstances, a grave responsibility would seem to rest on those in charge of plague hospitals, if they withhold the serum even for a few hours.

Efficient Treatment of Ivy Poisoning.—The extremely poisonous character of poison ivy or sumach, the especial susceptibility of some individuals, and the fact that cases of poisoning occur in which there is no evidence of positive contact with the plant, leading many to think that the poisonous constitu-