of the general history of syphilis there is one pathognomonic symptom I would again like to emphasize. If you scratch with your finger nail a very recent lesion which appears red, you will, even on a perspiring skin, immediately elicit fine scales. This is a symptom you will not find in any other erythematous disease and never in the erythematous syphilid. We have tried it dozens of times, and it helped us in the diagnosis always.

The bichlorid solution does a great deal of good; it relieves congestion and causes exfoliation of the skin. From this circumstance I would not say that the eruption is necessarily parasitic. As it stands now I think the general opinion is against its parasitic nature. I think I was able to show in these micro-photographs that the disease commences in contradistinction to the parasitic affections in the lower strata of the skin.

INFLUENZA IN CHILDREN.*

WM. CARVER WILLIAMS, A.B., M.D.
CHICAGO.

It is the writer's conviction that influenza is a more malignant malady in children than is generally acknowledged (a conviction fortified by his own experience and further strengthened by material gathered among his friends and from the literature), which has led him to venture this paper.

Without being able to furnish much, if anything, that is new on the subject, it yet seems justifiable here to emphasize some of the manifestations and possible calamities which may be traced to this disease.

Influenza vera, when the victim is a child, differs from the same disease in adults in proportion to the difference in constitution, vital resistance and recuperative power. It should not be forgotten in this connection that the disease is of a markedly toxic nature. The admirable article in Ashby and Wright's work states, however, that children are less likely to have some of the ordinary symptoms, such as backache, rigors and pain in the limbs, and more likely to earache, glandular swellings and a cough resembling whooping cough. The latter is also mentioned by Forchheimer. Blackader likewise states that the invasion in children is more often insidious and slow than in adults. These statements seem to be borne out in our experience. The very multitude of its manifestations, with the ensuing difficulty in accurate diagnosis, has so confused our statistics on influenza as to render many of them misleading; yet we are probably in greater danger of under- than of over-estimating its gravity.

Washburn and Ayre report 12 cases of death from pneumonia in children, of which the autopsies showed the Canon-Pfeiffer bacillus present as the main factor in the disease in 8 of them, though previously unrecognized.

Dr. F. E. Wynekoop, of the Chicago Department of Health, in a personal interview, stated that of the 2,460 examinations made just previous to the interview of cultures from throats in cases of suspected diphtheria, 677 showed the Canon-Pfeiffer bacillus as the only pathogenic organism, while of these 677 cases, in 173 the diagnosis of diphtheria was definitely made from the clinical findings—yet the Klebs-Loeffler bacillus was not present.

Age appears to have no influence on the susceptibility of the individual, and, while very young infants are somewhat less frequently attacked, there is some question as to whether it may not be due to better protection

rather than to natural immunity. In fact, there is nothing either in the literature nor in my own experience which warrants a belief in the existence of a complete natural immunity. The question of susceptibility would, therefore, appear to be one of the comparative degree to which different constitutions succumb to the disease—one child may have a so-called mild attack followed by recovery in a few days, while a brother or sister may be sick one or two months or longer. Much depends on the state of vital resistance of the victim, as in other diseases, but even more, in the writer's estimation, on the previous health of the mucous membranes. In a series of some fifty cases showing some of the more serious manifestations, he has rarely failed to obtain a definite history of indigestion, constipation, a common cold or some other previous disturbance of the mucous membranes.

Owing to the almost universal dissemination of the pathogenic organism during an epidemic, there is comparatively little possibility of accomplishing much in the way of prophylaxis at such times. And yet, because we know that the disease may be conveyed by contact, we should insist on such precaution as can be enforced to prevent this. Children should not be exposed to the contagion in the persons of other members of the family, nor sleep in illy ventilated rooms, nor breathe the vitiated air of public rooms and conveyances. Neither should they be allowed to risk contracting ordinary catarrhal "colds" from exposure. Mothers should be warned not to neglect attention to the children's bowels, and to allow them all the fresh air possible. Thus they may be enabled to resist and defeat an attack which might otherwise prove serious from this all too powerful enemy.

The division of the disease into various types for separate consideration is more or less arbitrary. The truth of the matter is, regardless of how the division is made, that every case one sees is of a mixed type. In other words, it is a case of influenza, and not of catarrh or neurasthenia—a specific infection, giving rise to various symptoms of which sometimes one, sometimes more than one, may be prominent, but of which a number are invariably present.

Those cases in which catarrh figures as the most conspicuous indication are perhaps more frequently met with. Starting with a rhinitis, or rhino-pharyngitis, the inflammation may spread either to the respiratory or alimentary tracts, or both. Wherever there is mucous membrane we may have this inflammatory reaction excited by the activity of the Canon-Pfeiffer bacillus. Thus we may find our little patients suffering from a rhinitis, a conjunctivitis, an otitis media, an inflammation of the accessory sinuses or of the pharynx, larynx, bronchi, lungs, stomach or bowels. The seat of choice for the severer forms of the infection is doubtless influenced greatly by the comparative resistance of the mucous membrane in various localities. The more nearly normal and healthy the membranes the fewer will be the grave manifestations of the disease. Hence the previously mentioned warnings in regard to the matter of prophylaxis.

The catarrhal manifestations of the disease appear, then, to be due mainly to the micro-organism itself. On the other hand, the nervous and circulatory and febrile types may be regarded as of toxic origin. The Canon-Pfeiffer bacillus, as in the case of that of Klebs-Loeffler, produces an exceedingly active and deadly toxin, capable not only of overpowering the vital organs, but of actually destroying their structure, to some extent, through degeneration. This being so, one can readily

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Diseases of Children, and approved for publication by the Executive Committee: Drs. H. M. McClanahan, A. C. Cotton and H. E. Tuley.

see how multiform in appearance and how serious in extent may be the history of such cases. One need not read far into the literature of the subject to find them, reported by careful observers, of meningitis, multiple neuritis, hemiplegia, neurasthenia and various mental disturbances—of endocarditis, pericarditis and myocarditis, arteritis, hemorrhagic nephritis and various other anomalies of the circulation with their consequences. Professor Monti of Vienna adds to these a glandular class, which includes those cases in which the lymphatic and other glands are chiefly involved, with a minimum of other clinical findings. I have included this class because I have known of several cases which would fall therein.

A form of the disease, until recently too little considered, is the chronic form, first described by Filatow. Cases have since been reported by Lord, Finkler, Wassermann, Ortner and others, where as high as 60 per cent. of cases having chronic cough, for instance, were caused by influenza bacillus, as proved by bacteriologic examination, in many instances it being found in pure culture.

Relapses and sequelæ during apparent convalescence are common, particularly the former. One can never be certain even when the child is to all appearances entirely restored to health that no sequels will follow. The writer has in two or three instances seen grave troubles develop days after recovery seemed complete.

The case reports which follow have been chosen without any attempt at covering the field. Aside from their individual interest, however, they indicate how widely divergent may be the chains of symptoms in disease traceable to the Canon-Pfeiffer bacillus, as well as the severity to which they not infrequently attain. It will be noticed that they are all more or less difficult to classify according to any of the rules laid down in different text-books.

CASE 1.—Baby W., aged 5 months, was taken ill in February, 1900, with what seemed a mild attack of influenza. The attack was, however, followed in four or five days by a bronchopneumonia, which ran an apparently typical course. As convalescence from this occurred, it was noticed that the child refused all food and even water. Examination of the throat revealed a highly inflamed catarrhal pharynx and fauces. Bacteriologic examination showed the characteristic influenza bacillus in large numbers. The angina and attendant dysphagia continued, in spite of all local and constitutional treatment, for so long a time that the child was in apparent danger of immediate starvation. There was also a middle ear complication. The little patient gradually developed decided symptoms of rachitis, which persisted until she was taken into the country in June of the same year. She did not regain the normal strength of children of her age until after she was a year old.

CASE 2.—M. C., aged 4 years, had a severe primary attack of influenza, verified by culture, in January, 1902, involving the middle ear and mastoid cells. The latter, however, never became suppurative and gradually responded to treatment—leeches, icebag, etc. Though the tympanic membrane was punctured three or four times, no pus was ever obtained. When the local trouble had subsided, the temperature still remained above normal, ranging mostly from 101 to 102.5, sometimes after exertion rising to even 104. A careful physical examination revealed a heart murmur, heard most plainly near the base and transferred most strongly toward the right. No other findings. This led to a diagnosis of post-influenzal endocarditis, and the child was kept in bed for three and a half months. During this time there was no appreciable improvement, the temperature curve and the heart murmur remaining about the same at the end as they were at the beginning of the confinement. About May 1 the little patient developed some indications of chorea, which yielded readily to treatment. She was allowed to get out of bed, was sent into the country

and began to improve from that time on. The temperature gradually resumed its normal status and the murmur has been slowly diminishing ever since, till at the present time, something more than a year after its appearance, it is barely perceptible. She is in every other respect apparently strong and well.

CASE 3.—Baby L., aged 8 months, taken ill during the month of January, 1900. Several other children in the family had been ill with undoubted influenza for several weeks. The little fellow became sick gradually. For several days a slight catarrhal condition of the mouth, eyes and nose had been present. Very soon he refused to eat and developed continuous fever, which was characterized by great severity. For days the temperature remained over 104, occasionally reaching 105.5 and even 106. His pulse was rapid, extremities were cold and blue, respirations were rapid and the patient showed some cyanosis. The most careful physical examination of the chest revealed only a few râles—indeed, the changes in the chest were insignificant as compared with the severe temperature range. The examination of the heart was negative. The abdomen at times was tympanitic. Food was taken in small quantities and only partly digested. The stools were thin, though diarrhea was not a marked symptom of the disorder. The ear was repeatedly examined by a competent otologist with negative findings. At another time the accessory sinuses, including the antrum of Highmore, were most carefully examined and no positive signs of disease found in these regions. After ten days of such alarming symptoms, the patient brightened, the temperature fell and he was undoubtedly on the mend. After several days of apparent improvement the patient again relapsed into the condition previously described. Two such relapses occurred. On each occasion alarming symptoms returned and it appeared that the resistance and the strength of so young an infant was hardly sufficient to withstand so severe an infection. After a course of several weeks, the patient recovered and very soon regained his strength.

The whole course of this infection was remarkable for the almost entire absence of symptoms save the hyperpyrexia and the great prostration. The catarrhal symptoms of the mouth, nose and pharynx were insignificant in comparison with the febrile condition of the child.

CASE 4.—J. R., 8 months old, taken ill about Christmas, 1899, with an otitis. During January had an attack of influenza, with typical catarrhal symptoms, followed by a pneumonitis, during which there developed symptoms of meningitis, such as retraction of head, strabismus and twitching. At this time there was nothing locally to indicate a connection between the meningitis and the ear trouble, which appeared rather better than worse. During the pneumonia the temperature ranged from 102 to 105. A typical crisis occurred, and the temperature resumed normal, or nearly so. After a few weeks the contraction of the neck abated and there followed a paresis so that the child was unable to hold up its head. This condition lasted nearly six months. There was a similar condition in all of the voluntary muscles. Reflexes were diminished. The child was three and a half years old before being able to walk.

At present, 5 years old, though he seems bright and interested in his surroundings, taking part in the games of his playmates, he is unable to talk more than two or three words. He makes his wants known cleverly enough by signs.

I have purposely omitted all mention of treatment of influenza in children, for the simple reason that my study and experience has led me to the belief that there is no specific treatment for any of its forms. The treatment is purely rational and symptomatic, and the indications can only be met as they arise. As these are so numerous and varied an attempt to cover the subject would be obviously useless as well as practically without value.

CONCLUSIONS.

I feel justified, both from experience and from perusal of the literature, in drawing the following general conclusions:

1. Children of any age, even early infancy, are susceptible to the influenza infection.

2. Repeated relapses and grave sequelæ are frequent in cases where the initial seizure is mild as well as where it is severe. Therefore, no attack should ever be considered as "mild" nor as recovered from until a sufficient time has elapsed.

3. Clinical diagnosis should always be supported by the evidence of the microscope, whenever possible, as the practical certainty thus obtained may be of great value in explaining future developments.

I desire to acknowledge no small indebtedness to Drs. I. A. Abt, J. C. Cook, M. P. Hatfield and J. W. White for their kindness in furnishing clinical histories and other data toward the writing of this paper.

58 East Forty-third Street.

DISCUSSION.

DR. R. B. GILBERT, Louisville, Ky.—I regret that the essayist has not given us the treatment. I have been looking for some article on the management of the disease and its sequelæ. I hope some one will elaborate this part of the subject. The ravages of the disease in the young are, of course, much more rapid and severe than in the adult, and often influenza is not recognized until bronchopneumonia has developed and a fatal termination has occurred. I recall a case during the height of an epidemic of influenza in our city in which the mother gave birth to a child while still in the febrile stage. The child presented the same symptoms and died in a few days.

DR. CHARLES G. KERLEY, New York—I agree that the treatment is wholly symptomatic and I do not understand how there can be a distinct method of treatment because in different seasons and epidemics it attacks by preference different parts of the body.

Two years ago there was an intestinal type in New York City, characterized by a diarrhea bordering on the dysenteric type and associated with intense prostration. This year the complication which seemed to be particularly frequent has been involvement of the ear. I have had ten cases this year in private practice in which it was necessary to perform paracentesis, and this does not represent all of the cases in which there was ear involvement. It is evident that the treatment must vary greatly with the prevailing type. One symptom which is usually prominent is prostration. I have seen two children die from the grip simply from the intensity of the intoxication, for the autopsy failed to show any lesion to account for the death. Another prominent symptom is fever, a point which has been well brought out in the paper. I have seen cases in which the high temperature could not be accounted for even by the most careful clinical examination by anything else than the grip. I am inclined to think that some of the cases reported as typhoid fever in children were really instances of grip infection. Many of these cases of grip run along for two or three weeks with hardly any symptoms but fever and intestinal disturbance. All that we can do is to carefully feed these children, control any tendency to hyperpyrexia and employ general supporting measures.

DR. P. J. BARCUS, Crawfordsville, Ind.—It has been pointed out in recent years that when there is an epidemic of scarlet fever at the same time as an epidemic of the grip, because of some possible antagonism between the two diseases, the type of epidemic of scarlet fever is somewhat milder than usual. I have observed such conditions present during the last six months in our vicinity, but I am unable to state whether this explanation is the correct one. It would be interesting to know the real facts. The recent epidemic of grip with us has been milder than usual. I do not think it is possible to formulate any general plan of treatment; it must be symptomatic because of the great variations in the type of the disease. Coal-tar remedies and indeed anything which tends to create depression should be avoided.

DR. H. M. McCLANAHAN, Omaha—We have seen a great deal of the grip in our city during the past winter. Two symptoms,

hyperpyrexia and extreme prostration, have impressed me as very important, and I frequently make the diagnosis on these symptoms. I have not had the opportunity of making the bacteriologic tests, and have, therefore, been compelled to depend on clinical examinations. I have noted that in these cases hydrotherapy has failed to control the fever as in other conditions producing high temperature. Another fact which has impressed me is the frequency of middle-ear trouble and mastoid abscess following the grip. The record of my cases shows that there are more cases of middle-ear disease from grip than from scarlet fever, and the same is true of mastoid abscess. I have adopted a treatment in a number of cases which is not at all new. It is simply the use of quinin bisulphate by suppository in doses of from six to twelve grains daily. It does not cause depression as do the coal-tar products. A few months ago I saw a case in consultation in which the attending physician was pushing phenacetin, although the child's finger nails were already blue. I remonstrated with him, insisting that such a line of treatment was very dangerous. I have seen recently a form of grip showing well-marked nervous symptoms.

DR. THEODORE J. CATLIN, Delano, Minn.—I recall a case, seen last winter, occurring in a boy who suffered a relapse. The fever kept up a long time, and the diarrheal symptoms were marked. The family called in another physician in my absence, and he made a diagnosis of typhoid fever, stating that he made this diagnosis not only on the symptoms mentioned, but on the presence at that time of a distinct eruption. I have met with other cases having skin eruptions similar to scarlatina and some of a marbled appearance. I would be pleased to hear if others have met similar skin symptoms in influenza.

DR. C. F. WAHRER, Fort Madison, Iowa—I should like to ask regarding a classification of the different forms of the grip, and also where is the anatomic seat of this disease. Some years ago, when it first appeared, it was thought that its anatomic seat was in the cerebrospinal system. We then met with the nervous and the catarrhal forms. The treatment is not satisfactory, but is symptomatic and purely empirical. If we knew the anatomic seat of this disease we might be able to suggest a more definite line of treatment.

DR. JOHN C. COOK, Chicago—In a hospital where I do the pediatric work I saw two babies die suddenly within forty-eight hours after birth, their mothers being ill with the grip. I have seen four cases of cerebral troubles this winter, all in consultation, in which the onset was as clearly that of grip as possible. In three of the cases at least one parent had the grip. A week from the time that one of the children had been seen by the attending physician I saw the case. The child was quiet and the only indication of meningeal trouble at the time was that one leg was a little stiff. Two days later the child had Cheyne-Stokes respiration, was rigid and presented the circulatory disturbances of the skin pointing to meningeal complication. Death occurred four days later. In another case, that of a baby nursing from a mother who was thought to have had the grip, the child had appeared stupid for four days before I saw it. Examination showed the pupils unequally dilated, and not responsive to light, the child totally blind. I was assured that the child had not been blind before this illness. It is still blind.

With regard to eruptions, influenza appeared in my neighborhood early in the autumn with an eruption, and the first two or three cases I saw presented an eruption closely resembling that of scarlet fever. As a rule, the throat symptoms were absent, as were also the rapid pulse and high temperature. In addition, all of the symptoms were those of grip. In the five or six cases which I allowed to mingle with the others in the family there was no spreading of the eruption to the other members, showing that the disease was not scarlet fever.

DR. J. H. HANNA, Covington, Tenn.—I saw a case presenting an eruption which I thought to be that of scarlet fever. A few months later measles appeared in the family, and the child developed the same set of symptoms minus the eruption. There was every symptom of measles at the time except the eruption, and although I looked confidently for the measles eruption it did not come. After this had subsided another

member of the family had measles. Three weeks ago the same symptoms developed and continued for several days, with all the symptoms of measles save the eruption. After the fourth day the measles eruption appeared and was well marked and typical.

DR. W. C. WILLIAMS, Chicago—I labored under a misapprehension when I prepared my paper, and consequently wrote so long a paper that it was necessary to omit many portions in the reading. This accounts for certain omissions to which reference has been made. There can be no question that the infection gains entrance through the mucous membranes just as in diphtheria. Some of the German investigators claim to have discovered the germ in the blood itself, and on the valves of the heart where there had been an endocarditis, but the evidence presented is not conclusive to my mind. I had an interesting series of four cases of eruption to report, in three of which the eruption closely resembled the rash of scarlatina. However, other children did not contract the disease. Along with this rash there were not the usual symptoms of scarlatina. All of these cases happened to be followed by an acute general adenitis, indicating that it was a sequence to whatever infection attacked the skin. With regard to treatment, I can only repeat what I said before, i. e., a bronchitis or an enteritis or other manifestation should be treated just as if there were no grip element. I would suggest the use of antiseptic douches to the mucous membranes, although admitting that the trial of this measure has not given encouraging results.

THE TREATMENT OF SCARLET FEVER, ITS COMPLICATIONS AND SEQUELÆ.*

H. M. McCLANAHAN, A.M., M.D.

Professor of Diseases of Children, College of Medicine, University of Nebraska; Physician to Child's Saving Institution and Methodist Hospital.

OMAHA.

It is difficult to discuss intelligibly the treatment of a disease having such wide variations in its clinical manifestations. In speaking of the cases, I shall use the designations: malignant, mild, severe and septic. Malignant cases are those overwhelmed by the poison and terminating fatally in one or two days. I shall say nothing of the treatment of this class. Mild cases are those presenting the characteristic symptoms without complications, the fever subsiding in from five to seven days. Severe cases are those having high temperature, profuse rash, often nervous symptoms, erythematous sore throat, but without membranous exudate. Septic cases are those presenting the symptoms of scarlet fever and, in addition, membranous exudate on the tonsils and other portions of the throat. These cases may or may not be severe at the inception. This is the class of cases in which we have the streptococci infection. These clinical divisions are simply made to indicate the kind of treatment more clearly. The subjects of quarantine, the isolation of the patient, the preparation of the room and the care necessary on the part of the attendants, while important, will not be considered in this paper. Certain general principles of treatment apply equally to all cases, and to these I shall first call your attention.

DIET.

To my mind the diet is exceedingly important and not rightly appreciated. For some years I have adopted the rule of keeping all patients on a liquid diet for not less than four weeks, the object being the prevention of renal complications. According to my observations, this sequela usually follows the mild cases. During the

last year I have seen five cases of nephritis following the mild forms of scarlet fever. These cases were not under my care during the acute attack. On the other hand, I have not observed this in any case where the principles of treatment indicated were faithfully carried out. By a liquid diet I mean milk, properly modified to suit the age of the patient. Many cases can be continued on this simple diet. In others we may vary it by the use of thin broths and fruit juices. In young children it is possible to modify the milk to suit almost every case. I have seen but two exceptions; for these whey was substituted with satisfactory results. In all cases the patients should have an abundance of water. In order to encourage the taking of food I recommend the use of carbonated waters, weak lemonade, orangeade, etc. I wish to emphasize the point that this diet is to be continued the full four weeks and not the first or second week, as it usually recommended. The continuance of the liquid diet during the period of convalescence is, to my judgment, the key to the situation.

REST IN BED.

The second general principle of treatment is to require that the patient be kept in bed during the entire course of the disease. I am aware that this is exceedingly difficult in mild cases, but, in my judgment, these are just the cases requiring this extreme care. The patient out of bed by the end of the first week and having the freedom of the house is more likely to be exposed to changes of temperature, to draughts of air, etc.; the influence of this exposure on the cutaneous capillary circulation adds to the work required of the kidney, and increases the danger from this source. In the severe cases it is not difficult to keep the child in bed, and I have a conviction that this is one reason why fewer cases of nephritis follow this severe form of scarlet fever.

LOCAL AND OTHER MEASURES.

In the mild cases, diet and rest in bed constitute the most important part of treatment. A spray or gargle of some mild antiseptic should always be used. Dobell's solution, properly diluted, or Seiler's tablets, dissolved in water, usually meet the indication.

I have very little confidence in any local treatment preventing otitis media. In my observation this complication occurs as frequently in mild as in severe cases. The bowels should be moved freely at the beginning of the attack and as soon as vomiting has ceased some mild aperient should be given. The bowels should move regularly during the entire course of the disease. The initial vomiting seldom requires special treatment. In cases where this is an early symptom, all food should be withdrawn for twelve to twenty-four hours. A simple fever mixture of liq. ammon. acet., properly diluted, may be given for the first few days.

ANTIPYRESIS.

Severe cases require more active treatment. When the fever is 104 it should be controlled. I have no doubt that the nervous manifestations are due to high temperature, as well as to the infection. Among antipyretic remedies, hydrotherapy ranks first. It is not only generally most efficient but, more important, it is the safest. However, in private practice the prejudice of the laity frequently prevents its proper application. Sponging with cool water is sometimes sufficient. More efficient, however, is the application of a towel dipped out of warm water about the body of the child, then, over this, is applied several thicknesses of a towel, wet with cold water. Should these means prove inefficient, I next em-

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Diseases of Children, and approved for publication by the Executive Committee: Drs. H. M. McClanahan, A. C. Cotton and H. E. Tuley.