THE ORIGIN OF GRAPHITE.

Sir: I am indebted to Mr. T. H. Clark for his comments in the September–October number of your journal on the origin of the graphite in the Sleaford Bay area of South Australia. A word in reply may perhaps not be out of place. Mr. Clark is of the opinion that “certain associations” described by me “seem to show that the graphite need not be wholly organic in origin.”

The associations leading Mr. Clark to this opinion have, however, been incorrectly interpreted by him. Their interpretation, in fact, is based on an erroneous view of the original nature of the sediments now carrying graphite. The first case is that of the garnet showing inclusions of graphite zonally arranged. Mr. Clark is here in error in supposing that the garnet is a product of the silication of the dolomites, and from a little closer inspection of the original paper he would have seen that the garnet is an almandine type, and not a grossular or andradite. The original sediment I consider to have been a siliceous shale, and the garnet to have been derived from biotite, as in the associated paragneisses, or in certain cases more probably derived from iron oxide in the original sediment (cf. pp. 196–7). Carbon dioxide can not be considered as a by-product in the synthesis of these zoned garnets.

It is to be noted that the majority of the graphite at Sleaford Bay is concentrated in rocks which were originally carbonate-free types and formed distinct stratigraphic units in an original sedimentary series. The assumption of an inorganic origin for the graphite necessitate the migration of CO_2 from the carbonate sediments to the siliceous and shaly types. While this is perhaps not to be regarded as a strong objection, it can scarcely be permitted to be used in support of an inorganic origin.

The facts brought forward in the original paper, and here in part restated, the absence of the vein type of graphite from this particular locality, are, in my opinion, more in harmony with an organic derivation than with one in which the reduction of CO_2 of an original carbonate molecule is involved. Alluding to the

DISCUSSION AND INFORMAL COMMUNICATIONS. 219

reactions which I have discussed, it may be emphasized again that in one of these only is experimental evidence available hitherto for the deoxidation of the oxides of carbon, namely, in the reversible system $2\text{CO} \rightleftharpoons \text{CO}_2 + \text{C}$. Even in the work of Dubrunfaut, which I have quoted, the carbon can be interpreted as derived from decomposition of carbon monoxide without intervention of hydrogen.

As has been noted in the original paper, when excavation work has proceeded further in the South Australian area, examination may show that more than one method of derivation of graphite is involved. Be that as it may, it is not improbable that the problem of graphite genesis, like not a few other vexed questions in petrology, will await its ultimate solution in high pressure experimental work in the laboratory.

CECIL E. TILLEY.

CAMBRIDGE, ENGLAND.

A FIELD METHOD OF REDUCING MAPS TO SCALE.

Two years ago, while engaged on some field work, the writer was called upon to reduce a geological map of a special area to a smaller scale in order to incorporate it in the general district map, then in the course of preparation. The usual method of locating the various points of the large scale map on the one of smaller scale by means of squares was found too slow and tedious and what was thought to be a new method, described in detail below, was devised which accomplished the work more quickly without loss in accuracy. It is based on the theorem of similar triangles and its very simplicity would suggest that it is already known and in use; but a diligent search in handbooks of Field Geology and verbal enquiry amongst a number of geologists would indicate that the application of this theorem to the reduction of maps to scale, while not new to some, at least has never been described. A brief note on its use may therefore be justified.

Let the upper polygon of Fig. 27 represent a closed plane table traverse which it is desired to transfer to a correlation sheet