medical men in this matter is beyond comprehension. With a 95 per cent. mortality rate staring them in the face they very often neglect to call in the surgeon until it is too late. Our homeopathic brethren are the worst offenders in this grave matter, for they usually give the patient a dose of medicine every fifteen minutes until he dies. The excuse usually given by the medical man for his neglect in these cases is that he was not sure of his diagnosis, and that he does not approve of the knife as a diagnostic agent. Exploration has enabled us to reduce the medical mortality of 95 per cent. in these cases to 45 per cent., and when we secure the co-operation of all practitioners it will fall much lower.

At this point I can not refrain from calling attention again to the restrictions that should be thrown around exploratory abdominal operations, for it is worse than useless for an inexperienced operator to open the abdomen in suspected perforation, or in cases of obstruction of the bowels, except when the obstruction is caused by strangulated hernia.

Chronic peritonitis is a condition very frequently demanding exploration. When the condition is tubercular the operation *per se* will often prove curative. We have learned with the development of abdominal surgery that the old idea that all cases of chronic peritonitis are tubercular was a mistaken one. The majority of cases of chronic peritonitis are the sequels of acute appendicitis, salpingitis or cholecystitis, but a great many cases develop independent of an acute inflammation. There are many chronic invalids in every community suffering from peritoneal adhesions caused by chronic non-tubercular peritonitis who could be restored to health and comfort through an exploratory opening. It is very important to remember that a patient may be a great sufferer from peritoneal adhesions who has never had an acute abdominal inflammation. When making an exploratory operation in a case of suspected peritoneal adhesions it is very important to so locate the incision that all parts of the abdomen may be reached. My personal preference is for an incision through the right rectus muscle, large enough to admit a hand, thus enabling the surgeon to explore the pelvis, the appendix and the gall bladder.

In conclusion, allow me to express the hope that this brief paper will be understood as denouncing the abuse of exploratory incisions as emphatically as it advocates their use. An earnest endeavor has been made to make such clear statements that no honest man can offer them as an excuse for the unwarranted substitution of the knife for surgical knowledge or diagnostic ability, and to convince the doubting ones that their refusal to use the knife as a means of diagnosis where other means have failed is to sacrifice human life on the altar of prejudice.

Fatal Emotions.—Keraval is director of an insane asylum in France, and he reports three deaths occurring there recently which demonstrate anew that a vivid emotion is capable of affecting the cerebral circulation to such an extent as to entail lesions immediately or rapidly fatal. One patient was a man of 44, with atheromatous degeneration, but apparently robust. He succumbed to a cerebral congestion induced by intense delight at the news that he was to be released from the asylum. The other cases were those of a man of 65, with ideas of persecution, and a man of 57, a general paralytic. Both died from cerebral congestion following a comparatively slight annoyance. The autopsy revealed in the last case the fresh hemorrhage among the lesions of old meningo-periencephalitis.—“Nord Méd.,” vi, 30.

Original Articles.

ILLUSTRATIONS OF MIXED INFECTIONS IN PULMONARY PHTHISIS,

WITH NOTES ON COMPARATIVE TREATMENTS.

PAUL PAQUIN, M.D.

Formerly Director of the Laboratory of Comparative Pathology, and Professor of Bacteriology and Comparative Medicine, University of Missouri.

ASHEVILLE, N. C.

The important significance of mixed germs and certain débris of the respiratory organs in phthisis is evidenced by the confusing variety of grave symptoms and lesions that accompany them and their too frequent fatal sequence. Yet the problems of mixed infections in this ruthless malady are still considered very largely with the same old indifference and skepticism that bred empirical treatments one hundred years ago. Impractical theories continue to sway a vast army of medical and laymen. The practitioners who seriously undertake to differentiate and individualize among their tubercular patients, with a view to ascertain the true conditions existing, and select appropriate remedies and proper nutrition are still in the minority; and the patients who look upon consumption with discriminating common sense are few. For many of the profession and laity consumption is consumption and usually fatal anyway; for a large number of others, in all walks of life, they will procrastinate till broken down, and then expect relief or cure in a few weeks or months—a thing impossible without a miracle. This is very unfortunate, because many consumptives are curable if they have the courage to begin to arrest the development early enough, the patience to persist, and if they are not too dissipated. No case is cured quickly. On the other hand, too many rash promises are made by too many doctors in desperate cases.

In order to present the significant differences in the microbiology and pathology of the various classes of consumptives, and suggest the errors of stereotyped treatments and the reasonableness of discriminate therapeutics, I beg leave to present a résumé of clinical and laboratory observations dating back a few years. For the purpose I have reviewed carefully a number of microscopic slides and macroscopic specimens and sections prepared for diagnosis and medical advancement in the course of practice. In fact, I have ransacked clinical histories related thereto and many works bearing on the subject. I shall here attempt to place briefly the most important data and findings before the profession. Each of the microscopic pictures represents the important things found in a given specimen of sputum. In some of them, however, I have grouped the findings of two or more differently stained slides of one expectoration in order to render more appreciable and clearer the various germs and débris present in the case. The macroscopic cut needs no attention at this point.

Fig. 1 shows the usual appearance of sputum in pure, so-called subacute pulmonary phthisis. Pus cells, bronchial epithelial cells, bacilli of tuberculosis and some mucus are present. There were none of the complication germs in this case of the kind and quality that do harm. The patient presented an area of dulness about three inches in diameter below the right apex, some fine moist murmurs and considerable expectoration, which was occasionally grayish and occasionally yellowish, or a mixture of both colors. Few bacilli of tuberculosis were present. The patient had come from

a country home away from the influence of smoke. In my opinion the grayish color was due to pathologic pigmentation and not to carbon inhaled. (This is often the case in most of gray sputa, many observers to the contrary notwithstanding.) He had been coughing six months and had lost twenty pounds and had had night sweats several weeks. His temperature had seldom risen above 101 2/5 F. in the afternoons; mornings it had often been subnormal. Being a case of clear tuberculosis, he was placed under the serum therapy as the specific basis of anti-tubercular treatment, and the various special phenomena of his malady were met as they appeared by such measures as were deemed best at the time. Nourishment and out-door air were of course insisted on. Neither creosote, guaiacol or inhalations were used. All these had been employed without good results for six months previously. Creosote and guaiacol seemed to have produced a severe form of dyspepsia, which yielded only to weekly lavage of the stomach. He steadily improved and got well in the course of eighteen months. He has shown no sign of illness for over four years. Infiltration seems to have disappeared absolutely. It would be difficult now, if not impossible, to find any lesion by any form of examination. Indeed, an insurance examiner of experience failed to find anything wrong in his chest.

The most significant features of this case are that it presented the evidence of pure pulmonary tuberculosis; that there never existed in the sputum any of the pathogenic germs that so frequently complicate tuberculosis; that the temperature seldom was higher than 101 2/5 F., and that the case recovered under the natural law measures indicated above, whereas he had been going down grade steadily and rapidly under the influence of the haphazard treatments which had been enforced before. This is one instance of several in my experience tending to show that pure tuberculosis is curable, often with comparative ease, by serum alone, provided such serum has been tested and is really effective. Such a product is not common on the market.

Contrast this case with the next two illustrated, viz.: pictures designated as Case 38 and Case C. The one marked No. 38 presents pus cells, bacilli of tuberculosis very few, and two forms of pus germs—the streptococci and staphylococci, each of them in small number. The sputum which contained these germs was gathered during the existence of an extensive area of more or less dense consolidation and involvement of lymphatic glands. All three of these classes of microbes are poison producers. Five weeks previous the patient had presented unmixed tuberculosis. He in the meantime contracted typhoid fever, and after his recovery from it the pus germs were present. Soon after small vomicae were found. His temperature arose a few times as high as 103 F. at night, and there appeared occasionally night sweats and chills, but generally the highest point was 101 F. This was when the pus germs were few. The disease gained ground steadily in spite of all possible care, and the remedial measures applied included many remedies approved by authoritative teachers. Classic inhalations, which were begun at the first sign of softening, served seemingly to increase the irritation. Creosote injured the stomach and kidneys. Combined antiseptic and antitubercle sera were used without changing the situation. The strychnin tonics, iodine and sera, combined, and climate had a better effect, causing reinforcements which prolonged life three years. Here we had not higher fever than in pure tuberculosis, except when pus

bacteria were numerous, and it was produced doubtless largely by pus toxins, or the mixture of these toxins and tubercular poison, and also by the influence of the lesions and their debris. Did not such complications argue earnestly for discriminate treatment?

Now consider picture of Case C and note all the differences carefully. The picture of its sputum shows pus cells, bacilli of tuberculosis, tetragenii and staphylococci (the latter numerous), each kind of which is capable of producing toxins that may separately, in a measure at least, cause cell, nerve and blood poisoning and fever. What special damage the combined toxins may cause I will not undertake to say. It is sufficient for our purpose to know that this mixed case, with only a very few small tubercular spots scattered in the upper third of one lung, exhibited temperature to 102 F. in the afternoon, coughed and expectorated a good deal of muco-pus with yellowish tinge, and did not do well under antitubercular serum alone. The disease yielded somewhat to a combined treatment, including antitubercle and antipyogenic sera hypodermically with iodine and various forms of judicious inhalations of respiratory sedatives and disinfectants. The patient is patched up, perhaps, only for a limited time. Yet to-day (five years after manifestation of disease) he is living in comparative comfort, without fever, and is working. But there yet remains in him evidence of lesions in quiescent state. These are liable to break down in the future, although with proper care he may retain his improvement.

I now draw your attention to Fig. 4, designated as of "Clinical Records," 1902. It shows sputum containing epithelial cells, few pus cells, tubercle bacilli, pneumococci, and some pyogenic bacteria. This patient exhibited infiltration in the upper half of the front of the right lung, with considerable aching opposite that locality. Fever arose as high as 104 F., but usually wavered between 100 and 103. A confused kind of pneumonia, limited in area, existed, and persisted for three months, after which resolution began and progressed slowly for weeks, leaving the lung impaired in elasticity, but not crepitant. Bronchial râles of a mucous kind were of course often present. Finally the pneumococci disappeared and the pus germs increased. As long as the latter were present in large quantities the temperature ranged from 99 F. in the morning to 102 F. in the afternoon. Later the pus germs decreased considerably, the bacilli of tuberculosis remaining scattered; then the fever wavered between normal and 99 or 100 F. At the time of this writing the temperature was between 97 and 99 F.

As soon as placed under my care this case received treatment applicable in limited pneumonia lesions of this type. It would, I am sure, have been useless to have put such a case under the influence of any therapeutic measure strictly within the line of the routine or specific treatments of consumption. What, for instance, could forced or voluntary inhalation have accomplished? Nothing but harm. The case lives in semi-comfort.

The microscopic display in Fig. 5 is from a case with a cavity four years old. La grippe and hemorrhages had been the first features of the development of the malady. The patient had done poorly most of the time for two years—la grippe recurring three times at least—but for the last year the case has presented comparatively few bacilli of tuberculosis, and this rarely. Cicatricial tissues and membranes seem to be slowly forming in the cavities and the healing process seems to be generally gaining ground. A mixed treatment,

A CASE OF UNMIXED INFECTION OF TUBERCULOSIS.
 ① BACILLI OF TUBERCULOSIS. ② PUS CELLS.
 ③ EPITHELIAL CELLS.
 TEMPERATURE 96 TO 101½ °F.

FIG. 1.

CASE 38-RECORDS 1899-1900. ① PUS CELLS.
 ② TUBERCLE BACILLI. ③ STREPTOCOCCUS
 ④ STAPHYLOCOCCUS.
 TEMPERATURE 99 TO 103 °F.

FIG. 2

CASE C-THREE YEARS AFTER EVIDENT MANIFESTATION OF
 CONSUMPTION. ① PUS CELLS- ② BACILLI OF TUBERCULOSIS.
 ③ MICROCOCCUS TETRAGENUS. ④ STAPHYLOCOCCUS ALBUS.
 TEMPERATURE 97½ TO 102 °F.

FIG. 3.

CLINICAL RECORDS 1902. ① EPITHELIAL CELLS. ② PUS CELLS. ③ BACILLI OF TUBERCULOSIS. ④ PNEUMOCOCCI. ⑤ MICROCOCCUS PYOGENES ALBUS. TEMPERATURE 99° TO 104°F.

FIG. 4.

FROM CAVITY THREE YEARS OLD. BACILLI OF TUBERCULOSIS ABSENT SEVERAL MONTHS. PATIENT NOW ABOVE NORMAL WEIGHT. ① PUS CELL. ② STAPHYLOCOCCI. ③ ZOOGLEA OF FINE COCCI UNDEFINED. ④ FILAMENTOUS BACILLI UNDEFINED. TEMPERATURE 95° TO 98½°F.

FIG. 5.

CASE 44. CLINICAL RECORDS 1902. ① PUS CELL. ② PNEUMOCOCCI WITH & WITHOUT VISIBLE ENVELOPE. TUBERCLE BACILLI HAD BEEN PRESENT UNTIL ONE WEEK PRIOR TO CONGESTION IN WHICH THESE GERMS APPEARED. TEMPERATURE 99° TO 103°F.

FIG. 6.

CASE 16 OF CLINICAL RECORDS OF 1900. ① PUS CELL. ② EPITHELIAL CELL ③ PNEUMOBACILLUS (OF FRIEDLANDER) ④ BACILLI OF TUBERCULOSIS. TEMPERATURE 99° TO 103°F.

FIG. 7.

CASE IN SERIES OF 1901-02. ① PUS CELL. ② BACILLI OF TUBERCULOSIS ③ PNEUMOCOCCI ④ CHARCOT-LEYDEN CRYSTALS OF ASTHMA. TEMPERATURE 99½° TO 103°F.

FIG. 8.

CASE OF DR. B. ANALYSIS 1901. ① BACILLI OF TUBERCULOSIS. ② GROUPS OF UNDEFINED COCCI. ③ SMALL BACILLI-PROBABLY PFEIFFER'S GRIPPE GERM. ④ FILIMENTS KNOWN AS LEPTOTHRIX BUCALIS. ⑤ EPITHELIAL CELLS-UPPER AIR PASSAGES.

CASE AA. ① PUS CELLS. ② BACILLI OF TUBERCULOSIS. ③ NUMEROUS PYOCYANUS BACILLI (GREEN PUS) TEMPERATURE 97° TO 101°F.

FIG. 9.

CASE II OF RECORDS OF 1899. FOETID EXPECTORATION ① PUS CELL. ② BACILLI OF TUBERCULOSIS. ③ LARGE BACILLI, PROBABLY THE LUMNITZER BACILLI OF FOETID BRONCHITIS. TEMPERATURE. 97 1/2° TO 101 1/2°F.

FIG. 10.

CASE No. 13 OF RECORDS 1898-99. LA GRIPPE & TUBERCULOSIS PRESENT. ① PUS CELLS. ② BACILLUS OF TUBERCULOSIS. ③ DIPLOCOCCUS IN CAPSULE. (PERHAPS AN INFLUENZA GERM) TEMPERATURE 98 1/2° TO 102 1/2°F.

FIG. 12.

FIG. 11.

No. 12 OF RECORDS 1901-02 (CASE BB) ① PUS CELL. ② BACILLUS OF TUBERCULOSIS ③ LOOKS LIKE THE ORGANISM KNOWN AS BACILLUS CRASSUS SPUTIGENUS. (PATHOGENIC GERM) TEMPERATURE 96° TO 100 1/2°F.

FIG. 13.

*MATERIAL IN SPUTUM OF BRONCHIECTASEY - PEYER -
AUTHOR FOUND SUCH SPUTUM MATERIAL IN
COMPLICATED CASE OF TUBERCULOSIS AND *
BRONCHIECTASEY - 1894.*

FIG. 14.

*ELASTIC FIBRES IN TUBERCULOSIS.
FOUND IN CASES OF SOFTENING, ETC.,
IMPORTANT DIAGNOSTIC & PROGNOSTIC
FEATURE, - (FROM PEYER, PAQUIN AND OTHERS.*

FIG. 15.

*FIBRINOUS CAST OF SMALL BRONCHI (PEYER & OTHERS
RARELY MET IN TUBERCULOSIS,
FOUND BY THE AUTHOR IN FEW INSTANCES
UNDER SPECIAL CONDITIONS.*

FIG. 16.

*TUBERCULAR LUNG. ADAPTED FROM DA COSTA AND
RECENT CASE OF PAQUIN'S. A CAVITY WITH PUS,
B RAMIFICATIONS OF LARGE BRONCHI. C DENSE
CONSOLIDATION IN APEX, D SCATTERED TUBERCULAR
LESIONS. E INFILTRATION.*

FIG. 17.

including serum as the chief remedy, the cacodylates and climatology, are constantly in use.

This expectoration shows the existence of staphylococci in small numbers, some zooglea of more minute cocci, which I am not prepared to name, and some bacilli, also of an undesigned character. At one time the bacilli of tuberculosis, different pus and air germs, were numerous, and the temperature then ran as high as 103 F. When this specimen was taken the tubercle bacilli happened to be absent, as they had been for some months, and the temperature did not reach higher than 99 F. In fact, the last year it rarely ran over this point, except during occasional periods of congestion. At present, however, in the morning it sometimes goes down as low as 96 and 95 F. without appreciable discomfort. (Such a depreciation of temperature in the morning is not an uncommon thing in many kinds of tuberculosis.) The patient is now better nourished than at any period of her life, being several pounds above her best weight of her healthiest days. Temperature still wavers between 96 and 98 F.

The sixth picture of the series, Case 44, shows pneumococci and pus cells. The patient had tubercular laryngitis mildly and a large consolidation with pain in the upper third of the left lung. The temperature ranged between 99 and 103 F. Three months before the bacilli of tuberculosis alone had been found. The patient escaped my observation during this period, after which he again presented himself for examination. At this time a congestion of the diseased lung was manifesting itself, the pneumococci appearing in small numbers, as in this cut. In two days, however, they increased enormously. Evidence of congestion and some inflammatory manifestations remained for several weeks. Then a few bacilli of tuberculosis were found. After the pneumonia symptoms had disappeared these bacilli appeared in greater numbers than ever before, but the temperature went down to between 98 and 100 F.

Here is a case of localized pneumonia, complicating consumption, something like in the preceding exhibit, and naturally it suggests also forms of treatment other than those for simple or different forms of mixed tuberculosis. Such measures were instituted, specific and general, to fit the conditions at the proper moment, including special diet, milk, cream, plasmon during sore throat particularly. The patient is now at work, walking miles daily, as an insurance agent, but he is hoarse, though not suffering.

The next picture, known in my records as No. 16, 1900, presents a peculiar mixed infection of tuberculosis. In it are exhibited Friedlander's pneumobacilli, with bacilli of tuberculosis, amid a considerable amount of pus and a few epithelial cells. This case, which died a couple of years ago, suffered at first in the right lung, front and back, from top to bottom, almost complete consolidation. No treatment succeeded in reducing it in the least. Portions of it softened after a while and were expectorated in pieces mixed with pus. About that time the left lung became seriously involved, and finally death resulted. These pneumobacilli remained almost to the last—long after various pus and putrefaction germs had appeared. The temperature never was normal and often went as high as 105 F. It usually wavered between 99 and 103 F.

The eighth figure or case in series 1901 and 1902 is intended to show the finding of the microscope in the sputum of a patient who was suffering from asthma and tuberculosis, with a very limited area of consolidation. The bacilli of tuberculosis and pneumococci were

present, and here and there we see Charcot-Leyden crystals and pus cells. As long as the pneumococci remained in large numbers and the infiltration kept prominent the temperature went as high as 103 $\frac{2}{5}$ F. By degrees resolution took place to a considerable extent, and these germs finally disappeared. The lungs remained impaired, however, and the bacilli of tuberculosis are still present at this writing. But fever with these germs alone does not rise higher than 100 F.

No argument is necessary to show in this case also the need of some circumspect, judicious differentiation in diagnosis and treatment.

Picture 9 (from a patient of Dr. B.) presents a rare combination of debris and bacteria. It includes nasal, bronchial and pharyngeal secretions and excretions with germs. In it we see bacilli of tuberculosis, some fine cocci that do not resemble much, and yet perhaps are, Pfeiffer's influenza (or grippe) bacteria, some long *Leptothrix buccalis* and some epithelial cells. I do not know the history of this case further than was indicated by this presentation, and that la grippe was said to exist in the patient when the sputum was gathered.

I now draw your attention to the tenth picture, designated as AA. In it you will find pus cells, bacilli of tuberculosis and small bacilli of green pus. In this case, analyzed in 1900, there existed scattered areas of consolidation, having respectively, doubtless, a tubercle or more as their nucleus. Patient suffered greatly from bronchorrhea, the matter from which was usually green in the morning. Bacilli of tuberculosis were not always present. Temperature seldom went higher than 99 F. Occasionally it reached 101. He has remained under my observation for nearly three years and is still living, although practically incapacitated for regular manual labor. Remedial measures, including hygiene, nourishment, climate, the cacodylates with iodine used at proper times, and occasionally antitubercle serum, seemed to have served the purpose of holding him in comparatively good condition and free from suffering. There is a decided improvement since 1900.

Fig. 11, designated as Case II, 1899, records bacilli of tuberculosis and other chief characteristics of sputum of a peculiarly offensive kind, which came from a case that fell under my observation in 1899. This patient also suffered from profuse bronchorrhea, but the color of the sputum was brownish and had a very fetid odor. He died with evidences of gangrene a few weeks after my first examination. Every form of supposedly useful treatment, including antiseptic inhalations, had been used without avail before I was called in consultation, and further efforts were not more successful. The patient had used creosote at enormous doses daily for a year without benefit.

The twelfth picture shows germs that I found in the sputum of a case which I examined for a St. Louis physician in 1898. You will notice that the bacilli of tuberculosis are present, some small pus cells and small diplococci and single cocci, with and without surrounding envelopes. (The difference in appearance of these supposedly grippe germs and those seen in Fig. 9 may be due to different staining processes and microscopic amplification.) Temperature 98 $\frac{4}{5}$ to 102 $\frac{4}{5}$ F.

The thirteenth picture, No. 12 of records, marked BB, shows the germs found in a patient who died in a sanitarium in 1901. He was not under my care and the history of the case is not at hand. This picture is shown chiefly to present a large germ which I had never found but once before in connection with tuberculosis.

While these thirteen pictures are practically all of different kinds, it will be noted that some pneumonia germs were present in four instances. In my experience of many years the pneumococcus has been present and a common complicator of pulmonary tuberculosis. It is perhaps as frequently found as either of the pus germs that reach the lungs, and sometimes assists in the pathologic processes.

If so many kinds of germs and complications in phthisis indicate anything, they surely indicate the necessity to pay the strictest attention to exact methods of differential diagnosis, whether the cases appear before us early or late, and that we can never be sure that we are applying proper measures without possessing proper knowledge of the pathology of each individual.

I shall now present to you a few pictures of another class, leaving out the bacteria, but showing certain debris and lesions which also have their indications in therapeutics. Two or three of these illustrations are familiar to most of my readers; they have been borrowed in part or in whole or adapted from Peyer and Da Costa.

In the first of these are to be seen the debris found in bronchiectasy. I have seen but one case in which tuberculosis and this bronchial affection clearly co-existed. But the combination is probably more common than this instance would indicate. The material in this expectoration includes pus cells, fatty globules (probably from broken-up leucocytes) and crystals of triple phosphates (irregularly formed) produced by the liberation of ammonia from the nitrogenous compounds in presence of magnesium. Bacilli of tuberculosis in such a mixture are not easily stained, and some other germs, too, are sometimes hard to discover. But the voluminous amount and mixed nature of expectoration in similar conditions should not mystify us or lead us to hasty conclusions eliminating tubercular complications. Prolonged observations, oft-repeated analyses in the course of weeks and months, may be required for proper diagnosis.

Next we come to the elastic fibers, Fig. 15, the significance of which every physician knows. Unfortunately, in sputum analyses they are too often overlooked or neglected. When it is known that they constitute evidence of the destruction of lung tissue, even in early stages of softening, their presence or absence should in all cases, if possible, be established as a guide in prognosis and treatment.

In the third of these, or Fig. 16, is represented a fibrinous cast of bronchi usually existing but rarely expectorated in cases of chronic croupous bronchitis, and very rarely found in connection with tuberculosis. In this particular case, however, the bacilli of tuberculosis had been present in small numbers, but there had been a prior history of chronic bronchial catarrh. At the time the patient came under my observation both sides of the bronchial tree suffered from this condition and dyspnea was very great. Partial infiltration of small areas existed here and there in both and increased slowly. All treatments failed to produce any good results. One day this patient, after using home inhalations of creosote, guaiacol and other antiseptics in glymol, which he had kept up under the advice of another physician for some months, undertook to climb a hill three hundred feet high. On his return he was seized with very serious congestion of both lungs and severe coughing, and died, asphyxiated, in two hours.

Next in order is the consideration of certain important gross lesions in pulmonary tuberculosis. For this purpose I here exhibit an illustration, Fig. 17, showing

the essential common features in certain stages of phthisis. With it before us, and bearing in mind the microbiology of tuberculosis, as discussed in this paper, let us endeavor to realize the merits and demerits of various treatments and their effects on the anatomy, the physiology and mechanism of the respiratory organs, as well as, in a measure, on the system generally.

This section of the right lung, trachea and large bronchi shows the following structures and lesions: first, the lower half of the lung appears sound, excepting a cavity (*a*) (although in this half of the lung the tissues may contain concealed small tubercles); second (*b*), the distribution of the large ramifications of the right bronchial tree (down to about the point where the bronchi are about one-sixth of an inch in diameter), penetrating infiltrated and consolidated spots in their course and communicating with the cavity mentioned above; third, a dense and old consolidation (*c*) at the apex, with slight surrounding infiltration; fourth (*d*), tubercle centers scattered in upper third with sound intervening tissue; fifth (*e*), a diffuse, comparatively new infiltration at the side in the middle lobe.

Now, it may be well at this point to refresh our memory about the normal anatomy and physiology of the lungs in order to realize the nature and significance of these lesions and the kind of therapeutics indicated.

Briefly, the lungs consist of a mass of aerial tissue made up of lobules which are "joined together by interlobular cellular membrane, each lobule constituting a distinct lung in miniature, having its own separate vein and artery; that these lobules do not communicate directly with each other;" that their ultimate terminals consist of a collection of microscopic air cells, "alveoli of exceedingly delicate nature, ranging in size from 1/200 of an inch to 1/100 of an inch in diameter," which are in communication with the air by means of filiform bronchioles; that these bronchioles are formed of delicate circular and longitudinal elastic fibers with flattened epithelium up to the point in their outward course, where they reach 1/24 of an inch in diameter; that further out they possess cartilage and a mucous coat, and gradually become larger and denser and stronger as they simplify and progress toward the trachea; that the pulmonary artery accompanies the bronchi branch for branch to the minutest divisions (making possible direct medication by means of the blood); "that the circulatory system forms a network of capillary blood vessels of the alveoli where the exchange of the gases (the oxygen of the air and carbonic acid of the system) takes place;" that amid this capillary network of the alveoli, beneath the endothelial layers, there exist lymph spaces which communicate with lymphatic vessels, that get eventually in communication with the air on one hand and the pleura on the other, thus opening innumerable channels for lymphatic and pleuritic infection by microbes inhaled into the lungs.

The important facts I wish chiefly to draw attention to are that the remote divisions of the bronchioles are of microscopic size, therefore invisible to the naked eye; that their infundibula are made up of minute cells of about 1/200 of an inch in diameter, and that the mass of the lungs consists of inflated, spongy tissue, which these cells and connected structures form, and that the blood vessels permeate them everywhere, assisting in the exchange of carbon for oxygen, and finally, that a system of lymph vessels, glands and interstices involves all the air cells and puts them in communication with the outside air and pleural cavity.

We may add that the bronchi, large and small, and the air cells secrete mucus and that, under irritation, they may form sufficient of it and of certain pathologic material to fill them more or less and obstruct patches of lung enough to prevent sufficient communication with the air; and, furthermore, that by inflammatory processes the plasma of the blood may in part filter through the vessels and cause infiltration of the lung tissue, making it more or less impermeable to the air inspired.

The facts related in the two preceding paragraphs suggest and explain some questions of interest in tubercular pathology and therapeutics. To begin with, the anatomy and physiology of the lymphatic system in the lungs show how nebular intrapulmonary medication is wholly impotent in perhaps all cases involving the necessity of reaching lymphatics, which is so common. The lymphatics do not breathe, therefore the vaporized drugs can not get into them. This anatomy and physiology show also how and why we have so many cases of tubercular pleurisy and so many kinds of secondary tuberculosis in various organs of the system. Diffusion of the disease germs takes place through the lymphatic channels.

In the circulatory system, affecting the spongy tissues so thoroughly, we find, on the other hand, the explanation of the success often obtained in the treatment of tuberculosis by internal or hypodermic medication where direct lung measures fail. More of this farther.

To bring the important question of therapeutics to a focus, I will specify that there are two chief divisions of treatment. First, the aerial division, which includes in part climatology, pure or medicated air, and the artificial efforts to drive remedies directly down into the lungs; second, the circulatory division, which means all the efforts to reach the lungs and influence the system generally by means of the blood and lymph circulation. This includes internal medicine, hypodermic injections and dietetics, either of which can be of service only after absorption of the remedy or food in the system. Both of these divisions have their advocates, both have their usefulness and limitations, and both are abused by those who prescribe mindless of pathology (among whom, by the way, are very widely known men, who seem not to understand analytical work or have it done crudely by inexperienced persons). In the first or aerial division belong numerous, often reckless, usually haphazard forms of inhalations, tracheal injections, sprays and nebula, which in their place and at the proper time have a measure of usefulness. How far can inhalations reach, for instance? Even if the normal anatomy and physiology of the lungs permitted it, it is evident that the consolidations exhibited in the picture of the section of the lung here presented would allow very little or no air to reach the alveoli or to reach properly even the periphery of the lesions, far less the structure of the central portions. By what law, I ask those who commend and apply intrapulmonary medication generally or as a routine practice, can any form of pressure cause medicated air, or any other remedy, to reach the ultimate structure of such diseased tissues and heal them? Is there any way to force nebula into such patches? Decidedly no, and when it is remembered that all tubercles, large or small, are to some extent accompanied by more or less consolidation, which precludes even proper natural respiration, who can not realize the futility of all efforts to force inhalations into them? Yet it is claimed that not only such consolidations *en masse* can be reached by fine nebula, but that the air cells that constitute them can also be medicated by this means under a pressure of thirty, forty or eighty pounds! This claim

is so remarkably contrary to known physiologic and pathologic facts that one who reads it or hears of it wonders at the daring of those who advocate it before the intelligent minds of the medical profession. It is wholly incompatible with the laws of physics and physiology. Indeed, while nebulized medicines are really very fine divisions, these are individually visible to the naked eye, and to drive one of them down in an air cell would be practically like forcing a globe the size of a baseball into a receptacle the size of a marble.

It must be remembered, besides, that the air cells are always partly filled with residual air, which can not be forced out, and that these cells can not be unduly dilated without tearing them open. Furthermore, the lungs within and without sustain an average (at our altitude, at least, Asheville, N. C.) of 10 to 12 pounds of pressure. They are maintained in their natural condition of partial expansion by this in and out equilibrium of air pressure. Suppose the pressure within the lungs is increased without compensation on the outside, what law would prevent dilation or rupture? We see, then, that certain kinds of intrapulmonary medication preached to the profession and public are far from always being right and safe. In truth, only three benefits can be derived from the best forms of direct pulmonary medication; that is to say, from tracheal injections and inhalations, to-wit: a limited degree of disinfection of such parts as such treatments may reach; second, a limited palliative influence over such irritated points as they traverse or touch (more or less poorly), and, finally, a limited degree of influence in subjugating cough. In no case is the *indiscriminate* use of such measures and their medicines right, and some conditions sometimes forbid them; for instance, seriously inflamed bronchi and acute congestions. To obtain the results possible by nebular medication, all that is necessary is to have an apparatus capable of forming fine nebula or sprays, such as the Globe Nebulizer or the Robertson and De Vilbiss instruments, and then have the patient inhale proper remedies at their leisure; or a room may be filled with nebula or medicated vapors. High pressure simply makes finer nebula; these, as made, are sent with the full force back of them into the throat, from which the patients breathe in what they can. It does not appear to be good practice to apply 40 to 80 pounds of pressure into the throat of tubercular patients. They have more than enough trouble in keeping their throats from hyperemia and secondary infections without increasing the tendency to these dangerous conditions by unnecessary rough treatment of sensitive and delicate membranes. Evidence showing the correctness of this estimate is not wanting.

I need not dwell on the value of the second division or circulatory medication. It is established beyond controversy that certain drugs in certain forms can be sent through the lungs by means of the blood just as easily as they can be sent to all tissues that receive nourishment. Whoever knows the physiology of the circulation must accept this as a fact. It is true that those lesions which are more or less solid fail to receive blood freely, but even they absorb the essentials of the blood from neighboring sources. Besides, this form of medication, administered hypodermically or otherwise, may be used so as to produce most beneficial effects by direct and indirect influences on the blood, the nerves, the digestive organs, etc., etc.

Having in mind the numerous different kinds of pulmonary tuberculosis, we can not escape the conviction

that many forms of treatment now in vogue, besides the mechanical means discussed here, are also useless and some of them dangerous, if not fatal, when applied without respect to the pathologic conditions present at the time, as, for instance, the use of creosote in a reckless manner and the abuse of alcoholics. Nor is full knowledge and realization of tubercular pathology and microbiology sufficient for the best work. It is, moreover, necessary to realize the condition of the digestive, eliminative, circulatory and nervous organs and to apply properly the means necessary to correct whatever faults they may exhibit. Observe, for instance, the sudden rise of temperature from normal to 103, 104 and 105 by indigestion alone, or in connection with absorption of pus, etc.! Such conditions do arise, complicating consumption by addition of auto-intoxication to it. I have seen such occurrences (auto-intoxication from the alimentary canal) arrested by one stomach and one intestinal lavage, which, by the way, is a particularly common necessity for the control of digestion, because indigestion is so common in almost every case of pulmonary phthisis in which ordinary drug measures are not always applicable.

To be specific as far as possible on a general question, what, then, constitute the most reasonable and most promising methods of treatment of tuberculosis pulmonalis, particularly in mixed conditions?

The details are not appropriate here. In fact, I have gone already beyond my first intention on the subject of treatment. But there is reason to indicate the basic elements that need be understood in mixed and simple tuberculosis if one is to pursue a right course and try to cure. In all cases we need first, nourishment, by which I mean the assimilation of proper food supplies, including under this head, for convenience, the breathing of proper air. Without these in sufficient quantity and good quality, no remedy will succeed. With them as foundation, diverse treatments, applicable in different conditions, may be properly selected after understanding each case. Thus the reliable, tested antitubercle sera (not the ordinary, untested commercial stuff), the cacodylates, the assimilable and non-irritable iodins, tonics, appetizers, stomach lavage, raw beef juice are useful in probably all cases, provided the proper moments for each or certain combinations of them are not neglected.

No matter what their nature, special complications demand special measures of relief. For instance, pulmonary congestion may be relieved by a hygroscopic plaster; cough by camphor, hyoscyamus, heroin, codein, hydrocyanic acid, etc.; dyspepsia by stomach and intestinal lavage, pepsin, takadiastase, and other dietetics; intestinal auto-intoxication by high colon lavage and intestinal antiseptics; anemia by soluble iron, milk albumin, protonuclein, etc.

I repeat that the most important problem in the treatment of consumption of any kind is that of nourishment. It must be sufficient in every instance or failure is unavoidable. Speaking in a general way, milk, cream, meats, a dozen of whites of eggs in water every day, whole eggs in sherry wine, fish, oysters and appropriate meat juices are among the very best foods. Potatoes and bread are among the most useful of vegetable origin. Unfortunately, in the majority of cases that come to the physicians, there arises the necessity of feeding them, at times, with more highly nutritious and assimilable substances than is afforded by natural products. This is especially common in enfeebled stomachs and sore throats. Thus, in throat tuberculosis, for instance, both local pains and indigestion are usually present. The

pain produced by swallowing is so excruciating at times that even water produces torture. In such unfortunates, raw eggs, milk and cream are usually the most easily swallowed natural foods. But they may not always suffice, and often they upset digestion very much. This is often true of sweet milk. I have tried all sorts of artificial nourishments to assist these, or alone in many conditions. I have made numerous trials of many manufactured and home-made products in cases accompanied by poor digestion, assimilation and painful deglutition. Several are pleasant, but few are nourishing enough.

There is one kind of food used by many now which is possessed of special virtue as nourishment and medicine at the same time. It is the fresh juice made daily from raw beef.

Professors Charles Richet and J. Héricourt, both medical experimentalists and teachers of international reputation, presented to the Congress of Medicine in 1900, and afterward to the Academy of Medicine of Paris, many reports on their clinical experiments with such juice in consumption. Since then numerous physicians in every country have verified the claims of these scientists that raw beef juice is not only nutritious to the highest degree, but that it possesses specific curative virtues to arrest the malady. But it must be raw, not heated in the least. Its serum must be unimpaired, for this is internal sero-therapy, so to speak. To get good results large quantities must be given daily—all of the juice of four to six pounds of steak between meals. Strange to say, this does not interfere with the regular meals or the application of a special diet. It will be a pleasant surprise to those who have horror of raw meat to know that this fluid is very palatable and may be made delightful. My experience with raw beef juice in large quantities has been most gratifying, particularly in connection with plasmon, the cacodylates, serum, iodine, etc.

The unfortunate part of this most useful measure is that few presses can extract more than 25 per cent. of the juice from raw meat, thus reducing the chances of its use and making it expensive. I have tried everything in the shape of press I could find on the market, excepting high-priced hydraulic presses, and only the Hayes Meat Crusher have I found to extract more or less easily 75 to 85 per cent. and stand the effects of hard pressing without breaking, but it is in the present size a little expensive. As it is essential, as I have said, to use large quantities of beef juice to get any good results, and, as all that exists in six pounds of good beef is usually necessary every day, it will be seen at once how costly it would be to use presses extracting only 25 or 30 per cent.

In conclusion, I urge the study of the results obtained in France and elsewhere with the cacodylates, reliable sera, iodine and beef juice in consumption.

THE NEW UNITED STATES PHARMACOPEIA.

JOSEPH P. REMINGTON, PH.M., F.C.S.

CHAIRMAN COMMITTEE OF REVISION OF THE U. S. PHARMACOPEIA.
PHILADELPHIA.

The numerous inquiries from all parts of the United States with regard to the appearance of the new Pharmacopeia would seem to require at this time a communication as to the progress of the revision, and the writer takes this opportunity of stating that the work is progressing favorably. On account of the death of the distinguished chairman of the Committee of Revision, Dr. Charles Rice, May 13, 1901, revision was necessarily retarded; nevertheless, the subcommittees