the tumor, which is seen to be confined entirely to the group of adductors (*e*).

The hæmorrhage was almost *nil*, but the shock was so great that during the latter part of the operation his pulse became very feeble and rose to 180. In order to try its effect in a case of simple shock, Dr. S. B. Woodward, who kindly assisted me at the operation, injected, at my request, twenty ounces of normal salt solution into the median cephalic vein of the right side. The pulse fell at once to 150, of increased strength, and although it rose during the afternoon to 170, on the following morning it had come down to 140. Nourishment and stimulants were freely administered, and he began slowly to regain his strength. The temperature, which had risen as high as 102°, remained normal after the fifth day; and on the seventh primary union of the wound, except along the track of the drainage-tube, was complete. He continued to

FIG. 2.
a, a, a, Inner hamstring muscles. b, Sciatic nerve. c, Biceps femoris. d, Head of femur. e, Tumor.

improve steadily, was up on crutches; and on January 12th, six weeks after the operation, was reported to be gaining both strength and flesh. On the following day, however, the inner part of the stump became red, hard, oedematous and painful. It was manifestly a recurrence, but seemed to be limited to a small movable nodule, and its removal was therefore attempted by my successor at the hospital, Dr. S. B. Woodward. The whole stump proved to be infiltrated with soft broken-down material, similar to that found at the first operation, and the disease had extended round on to the buttock and involved the gluteal muscles. Everything possible was removed, but recurrence was noted again two weeks later. He gained temporarily in strength, left the hospital in February, and died at home the last week in March, eight months after the receipt of the original injury.

The examination of the tumor by Dr. W. S. Miller, the hospital pathologist, showed that it consisted of a

large soft mass confined to the adductor muscles. The centre of this mass was composed of broken-down coagula, around which was a layer of soft, round-celled sarcomatous tissue containing numerous anastomosing blood-vessels. Diagnosis: angio-sarcoma.

Clinical Department.

A CASE OF HEMIANOPSIA.

BY CHARLES M. HOLMES, M.D., NORTHAMPTON.

Mr. H. R., was wounded through the head by a rifle-ball when fourteen years of age, forty-five years ago. In passing an open window through which some men were firing at a target, his head intercepted a bullet which struck him in the right temporal region, at a point three-fourths of an inch in front of a vertical line from the right external auditory canal and two inches above the level of the same point. The bullet passed through the head in a backward and slightly upward direction, and came out in the median line three inches above the external occipital protuberance of the occipital bone, the distance between the entrance and exit measured on the convexity of the skull being seven inches.

At the point of entrance the wound was apparently but little larger than the bullet which caused it, and this point is now marked by a slight but distinct bony prominence. The damage to the skull was greater at the point of exit, where two fragments of bone as large as a "finger-nail" were removed by the attending physician, Dr. Rice, of Leverett, Mass., deceased. The patient says that these pieces of bone were preserved for many years, and he has often seen them. This wound is marked by a large pit in the scalp. This spot is very sensitive to pressure at times, and the patient says that a very slight blow there will cause a convulsion. According to the statement of the patient, Dr. Rice told him afterward that he saw on the grass "several spoonfuls of white brain substance lying there in the clotted blood."

Loss of consciousness was immediate, and no one expected that he would live. After a hard struggle of six months he was again able to get about; but he was hemiplegic, and this condition has remained, though he now has some use of his left leg, and can walk with the aid of a cane if he uses great care and keeps on a level and smooth surface.

The whole left side shows the results of the injury. He had so nearly attained his growth in height at the time of the accident that there is little if any difference in the comparative lengths of the long bones of the body; but there is a difference of from half an inch to two inches in the girths of the two sides, on account of failure in development and also on account of atrophy from disuse of the left side necessitated by the paresis of that side.

His tongue is not markedly affected, though he says he cannot manage his food so well if it is chewed in the left side of his mouth. Sensation is dulled on the whole left side of the body, though there is no complete anaesthesia. The left side feels numb and cold, especially if he is tired. The senses of taste and smell are impaired on the left side. Hearing is also a little im-

paired, but on both sides, and is perhaps due to other causes than the injury to the brain. The knee-jerk is exaggerated in both legs, and equal. No ankle clonus.

Soon after he was injured, perhaps immediately, epileptic convulsions came upon him, and have continued since. They begin on the parietic side, but become general, and are very severe. At times, also, he has attacks of epileptic mania, in which he is very violent and unreasonable. At these times he has false hearing and delusions of persecution. He recovers from these attacks with very incomplete memory of what has occurred. He is at all times very irritable.

The patient complains of poor vision in the left eye; and on examination it is discovered that he presents the symptom of left bilateral hemianopsia, with the line vertical and passing through the point of fixation. His pupils react readily to light and distance, and Wernicke's symptom is absent.

Medical Progress.

RECENT PROGRESS IN LEGAL MEDICINE.

BY F. W. DRAPER, M.D., BOSTON.

DEATH BY DROWNING.

In a discussion of the subject of the resuscitation of those apparently dead by submersion, Professor Brouardel insisted that death by drowning presented itself under several distinct forms, according to the mechanism of asphyxia which predominated.¹ Indeed, the submersion may not cause asphyxia at all, but death may result the moment the individual falls into the water, the suddenness of the fatal effect depending on the influence of shock and of cold upon peripheral branches of the trigeminal and superior laryngeal nerves. In such an event, respiration and circulation are inhibited. Water does not enter the air-passages, and the person if rescued seasonably may be resuscitated.

A second mechanism of death by submersion has the following phenomena. The person sinks to the bottom of the water and remains there a brief time. During the first minute, the air-passages resist the intrusion of the water successfully, the muscles of respiration preventing this effect by their tonic contraction. This muscular spasm then ceases, and the water penetrates into the bronchi. But even under these circumstances, resuscitation is practicable if the rescue is prompt.

The third and most common variety of drowning is this: The individual struggles, tries to swim, sinks to the bottom, then rises to the surface; and in these irregular movements he causes contractions of the diaphragm which promote the regurgitation of the contents of the stomach and these pass into the windpipe when efforts at respiration are made. Water also enters the air-passages and the blood takes up a quantity equal to a quarter or a third of its own total amount. This effect is seen when comparison is made between the relative number of the blood globules in the normal condition and after drowning. In cases of drowning of this variety, the air-cells and bronchioles are filled with mucus and water in the form of fine froth and the cells of pavement epithelium in the mucous mem-

brane undergo an appreciable degeneration in consequence of the passage of water through them. These combined conditions are such as to prevent absolutely a restoration to life by any known method of resuscitation.

DEATH FROM SUBMERSION IN FLOUR.

Dr. Ragnar Falck, of Lund, publishes in the *Nordiskt Medicinisches Arkiv*² the case of a man who had been buried in flour, who had fallen down into the room used for blending the various kinds of flour, and which was capable of containing 220 sackfuls. Rigor mortis was present in the body; fine flour was found in the hair, the external meatus, within the eyelids, in the nares, within the lips and all about the throat. The mouth and œsophagus were filled with "paste," the stomach was empty, the cerebral membranes and vessels were hyperæmic to a marked degree, the heart was small and contained a little dark blood in the left auricle, and the lungs were incompressible and full of blood. It is particularly interesting to note that microscopical examination resulted in finding flour in the smallest bronchial twigs. This circumstance makes the case resemble death from drowning. French medico-legal authorities, however, state that in cases of submersion in dust-like substances, as when children are buried in ashes, nothing enters the respiratory organs, and Berenguier and Jaudien, after experimental studies, came to the conclusion that spasm of the glottis makes the entry of dust-like material impossible. Inferences, however, drawn from children's accidents, and from experiments with small animals, are hardly to be depended upon in cases of adults, and in Germany Casper, Hoffmann, and Maschka have published several cases of submersion in sand and similar substances, in which the material could certainly be found in the larger bronchi.

A NEW METHOD OF RESUSCITATION IN CASES OF DROWNING.

Laborde, of Paris, has demonstrated the practical utility of a new method of procedure for the restoration of those apparently dead by submersion. Having observed in his physiological experiments on animals that, in cases of suspended animation during anaesthesia, it was possible to stimulate the function of respiration by rhythmical traction on the tongue, he applied this knowledge to some cases of asphyxia by submersion and found the application to be both easy and satisfactory. The theory of the manipulation is that irritation of the base of the tongue, especially such as is effected by traction, stimulates the respiratory reflex. The details of the procedure are as follows: The tongue is seized either with forceps or with the fingers protected by a handkerchief or other similar fabric to prevent slipping and is then subjected to regular, rhythmical traction and backward movements, corresponding in frequency to the normal respiration. The teeth meanwhile are held apart by a spoon or other expedient adapted to the need, the object being to give room for forcible and vigorous to-and-fro motions of the tongue. The author urges persistency in the measures taken for the restoration of those apparently drowned and promptly taken from the water; and he insists that prolonged efforts will sometimes be rewarded even in cases that seem hopeless from the beginning.

¹ La Semaine Médicale, August 3, 1892.

² Lancet, December 26, 1891.