Annals and Magazine of Natural History

Series 1

ISSN: 0374-5481 (Print) (Online) Journal homepage: <http://www.tandfonline.com/loi/tnah07>

XXXIX.—Descriptions of Chalcidites discovered in Valdivia by C. Darwin., Esq.

Francis Walker F.L.S.

To cite this article: Francis Walker F.L.S. (1842) XXXIX.—Descriptions of Chalcidites discovered in Valdivia by C. Darwin., Esq., *Annals and Magazine of Natural History*, 10:65, 271-274, DOI: [10.1080/03745484209445234](https://doi.org/10.1080/03745484209445234)

To link to this article: <http://dx.doi.org/10.1080/03745484209445234>

Published online: 04 Dec 2009.

Submit your article to this journal [↗](#)

View related articles [↗](#)

XXXIX.—Descriptions of Chalcidites discovered in Valdivia
by C. Darwin, Esq. By FRANCIS WALKER, F.L.S.

Pachylarthrus Sariaster, Mas. *Viridis*, antennæ luteæ, pedes flavi, alæ limpidæ.—Fem. *Cupreus*, antennæ nigre, pedes fulvi, femora viridia.

Mas. Corpus crassum, convexum, nitens, viride cupreo-varium, scite squameum, parce pubescens: caput magnum, transversum, breve, viride, thorace latius; vertex latus, cupreo-varius; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: os flavum; palpi maxillares clavati; antennæ luteæ, graciles, subfiliformes, 13-articulatæ, thorace paullo longiores; articulus 1^{us} longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes usque ad 10^{um} mediocres, subæquales; clava longiconica, acuminata, articulo 10° fere duplo longior: thorax ovatus, robustus: prothorax transversus, brevis, antice angustus: mesothoracis scutum longitudine latius; parapsidium suturæ bene determinatæ, postice approximata; paraptera et epimera magna; scutellum subconicum, sat magnum: metathorax mediocris, obconicus, declivis: petiolus sat longus: abdomen rhombiforme, contractum, læve, glabrum, latitudine non longius, thoracis dimidii longitudine; segmentum 1^{um} maximum, dorsum obtegens: pedes flavi, simplices, subæquales; coxæ virides; ungues et pulvilli fusci: alæ limpidæ; squamulæ luteæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis ulnari vix brevior, cubitali duplo longior; stigma minutum.

Fem. *Cupreus*, parum nitens: caput thorace vix latius: palpi maxillares simplices: antennæ nigre, subfiliformes, thorace non longiores; articulus 2^{us} cyathiformis; 5^{us} et sequentes usque ad 10^{um} subæquales, approximati; clava conica, articulo 10° longior: petiolus brevis: abdomen ovatum, supra fere planum, subtus carinatum, apice acuminatum et attenuatum, thorace vix longius; segmentum 1^{um} magnum, 2^{um} et sequentia breviora: pedes fulvi; coxæ virides; femora viridia, apice fulva. (Corp. long. lin. 1—1½; alar. lin. 1¼—2¼.)

Dicyclus Lynastes, Fem. *Viridi-æneus*, antennæ nigre, pedes fulvi, femora ænea, alæ limpidæ.

Corpus breve, convexum, viridi-æneum, nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ nigre, graciles, subclavatæ, thorace vix longiores; articulus 1^{us} longus; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes usque ad 10^{um} subæquales, approximati; clava conica, compressa, acuminata, articulo 10° multo longior: thorax ovatus: prothorax transversus, brevis, antice angustus: mesothoracis scutum longitudine latius, parapsidium suturæ remotæ, sat bene determinatæ, postice approximata; paraptera et epimera magna; scutellum subconicum, mediocre: metathorax brevi-obconicus, declivis: petiolus brevis: abdomen subrotundum, supra fere planum, subtus carinatum, apice acuminatum, thorace brevius; segmentum 1^{um} magnum: pedes fulvi, simplices, subæquales; coxæ æneæ; femora ænea, apice fulva; tarsi apice fusci: alæ limpidæ; squamulæ picæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis ulnari vix brevior, cubitali longior; stigma minutum. (Corp. long. lin. ¾; alar. lin. 1¼.)

Lamprotatus Natta, Mas. *Viridi-cyaneus*, antennæ nigre, pedes flavi, alæ limpidæ.

Corpus angustum, sublineare, convexum, viridi-cyaneum, nitens, scitissime squameum, parce hirtum: caput transversum, breve, viride, thorace latius; vertex latus, æneo-varius; frons impressa, abrupte declivis: oculi rufi,

mediocres, non extantes: antennæ nigræ, graciles, subfiliformes, thorace vix breviores; articulus 1^{us} longus, sublinearis; 2^{us} longicyathiformis, 3^{us} et 4^{us} minimi; 5^{us} et sequentes usque ad 10^{um} subæquales, approximati; clava longiovata, acuminata, articulo 10^o multo longior: thorax longi-subovatus: prothorax transversus, medietate, antice paullo angustior: mesothoracis scutum longitudine latius; parapsidum suturæ bene determinatæ, postice approximata; paraptera et epimera magna; scutellum subconicum, mediocre: metathorax magnus, declivis, obconicus: petiolus sat longus: abdomen breviovatum, læve, glabrum, thorace multo brevius: pedes flavi, simplices, subæquales; coxæ virides; tarsi apice fusci: alæ limpidae; squamulæ fuscæ; nervi fulvi. (Corp. long. lin. 1; alar. lin. 1½.)

Lamprotatus Bisaltes, Mas. Aeneo-viridis, antennæ nigræ, pedes flavi, femora viridia, alæ limpidae.

Corpus convexum, aeneo-viride, nitens, scitissime squameum, parce hirtum: caput transversum, breve, thoracis latitudine; vertex latus; frons impressa, abrupte declivis: oculi rufi, medioeres, non extantes: antennæ nigræ?: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ bene determinatæ, postice approximata; paraptera et epimera magna; scutellum conicum, mediocre: metathorax brevi-obconicus, declivis: petiolus brevissimus: abdomen sublineare, læve, glabrum, fere planum, thorace angustius et multo brevius; segmentum 1^{um} maximum: pedes flavi, simplices, subæquales; coxæ virides; femora viridia, apice flava; tarsi apice fusci: alæ limpidae; squamulæ picæ; nervi fusci; nervus humeralis ulnari fere duplo longior, radialis ulnari paullo brevior, cubitali paullo longior; stigma parvum. (Corp. long. lin. 1; alar. lin. 1¼.)

Lamprotatus Orobia, Fem. Viridis, cupreo-rufus, antennæ nigræ, pedes rufi, femora basi viridia, alæ limpidae.

Viridis, convexus, nitens, scitissime squameus, parce pubescens: caput transversum, mediocre, thoracis latitudine; vertex sat latus; frons abrupte declivis, ad antennarum receptionem excavata: oculi picci, medioeres, non extantes: antennæ subfiliformes, nigræ, graciles, pubescentes, thorace non longiores; articulus 1^{us} longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} vix conspicui; 5^{us} longicyathiformis; 6^{us} et sequentes breviores: thorax longiovatus: prothorax medietate, transversus, antice angulum utrinque fangens non angustior: mesothoracis scutum longitudine vix latius; parapsidum suturæ bene determinatæ, postice approximata; paraptera et epimera magna; scutellum mediocre, conicum: metathorax medietate, declivis, obconicus: petiolus brevis: abdomen ovatum, nitens, læve, fere glabrum, apice acuminatum, thorace paullo brevius: segmenta antica magna, postica breviora: oviductus non exertus: pedes graciles, recti, subæquales, pallide rufi, pubescentes: coxæ virides; trochanteres picæ; femora basi viridia; tarsi apice fusci: alæ limpidae, medioeres; squamulæ viridi-picæ; nervi fusci; nervus humeralis ulnari fere duplo longior, cubitalis radiali brevior; stigma minimum. (Corp. long. lin. 1¼; alar. lin. 2¼.)

Var. β.—Capitis vertex viridi-aeneus: thoracis discus cupreo-aeneus: abdomen cupreo-varium.

Pteromalus Megareus, Fem. Viridi-aeneus, antennæ nigræ, pedes flavi, femora viridia, tibiæ fusco-cinctæ, alæ limpidae.

Corpus viridi-aeneum, convexum, parum nitens, scitissime squameum, parce pubescens: caput transversum, breve, thorace paullo latius; vertex sat latus; frons abrupte declivis, excavata: oculi picci, medioeres, non extantes: antennæ nigræ, subelavate, pubescentes, sat graciles, thorace non longiores;

articulus 1^{us} viridis: thorax breviovatus: prothorax brevissimus, supra vix conspicuus: mesothoracis scutum latum; parapsidium suturæ non bene determinatæ; scutellum subrotundum, mediocre: metathorax brevis, declivis, obconicus: petiolus brevissimus: abdomen longiovatum, nitens, læve, fere glabrum, supra planum, subtus carinatum, apice acuminatum, thorace longius et angustius: pedes flavi; coxæ virides; femora viridia, apice flava; tibiæ fusco-cinctæ; tarsi apice fusci: alæ limpidæ; squamulæ piceæ; nervi fulvi; nervus humeralis ulnari multo longior, cubitalis radiali multo brevior; stigma minimum. (Corp. long. lin. 1; alar. lin. 1½.)

Closterocerus Xenodice, Mas et Fem. *Viridis cyaneo-varius, abdomen cupreum, antennæ nigræ, pedes flavi, femora viridia, tibiæ fusco-cinctæ, alæ fusco-maculatæ.*

Mas. Corpus sublineare, angustum; depressum, læte viride, nitens, læve, parce hirtum: caput transversum, brevissimum, inter oculos impressum; vertex sat latus; frons abrupte declivis: oculi mediocres: antennæ subsectæ, nigræ, hirtæ, thorace non longiores: thorax ovatus: prothorax brevissimus: mesothoracis scutum longitudine vix latius; parapsidium suturæ vix conspicuæ; scutellum subovatum: metathorax obconicus, declivis, mediocre: petiolus brevissimus: abdomen sublineare, cupreum, basi cyaneo-viride, thorace angustius non longius: pedes flavi; coxæ virides; femora viridia; tibiæ fusco-cinctæ; tarsi apice fusci: alæ mediocres, ciliatæ, fulvotinctæ, in discum obsolete fusco-maculatæ; squamulæ piceæ; nervi fulvi; nervus ulnaris humerali multo longior, radialis vix ullus, cubitalis brevissimus in alæ discum abrupte declivis; stigma minimum.

Var. β.—Tibiæ omnino fuscæ: alis maculæ vix conspicuæ.

Var. γ.—Alæ omnino limpidæ.

Fem. Cyaneo-viridis: antennæ setacæ, thorace breviores: abdomen ovatum, thorace paullo brevius vix angustius. (Corp. long. lin. ½—¾; alar. lin. ½—1.)

Var. β.—Tibiæ nigro-fuscæ, apice flavæ.

Platygaster Paches, Mas et Fem. *Atra, antennæ nigræ, pedes nigri, tarsi picei, alæ subfuscæ.*

Mas. Corpus convexum, atrum, nitens, læve, fere glabrum: caput transversum, breve, thorace vix latius; vertex sat latus; frons abrupte declivis: oculi picci, parvi, non extantes: antennæ nigræ: thorax ovatus: prothorax brevissimus, supra vix conspicuus: mesothoracis scutum longitudine latius; parapsidium suturæ non bene determinatæ; scutellum subrotundum, non productum: metathorax parvus, obconicus, declivis: petiolus brevis: abdomen longiovatum, thorace paullo angustius non longius; segmentum 1^{um} magnum, 2^{um} et sequentia breviora: pedes nigri, subæquales; femora clavata; genua picea; tibiæ clavatæ; tarsi picei: alæ subfuscæ; squamulæ piceæ. (Corp. long. lin. ½; alar. lin. ¾.)

Fem. Caput thoracis latitudine: antennæ subclavatæ: abdomen longiovatum, apice acuminatum, thorace angustius et paullo longius.

Inostemma Quinda, Fem. *Atra, antennæ flavæ, apice fuscæ, pedes flavi, alæ limpidæ.*

Corpus atrum, angustum, sublineare, convexum, læve, nitens, pubescens: caput transversum, subrotundum, thorace vix latius; vertex latus; frons abrupte declivis: oculi parvi, non extantes: antennæ subclavatæ, flavæ, apice fuscæ: thorax ovatus: prothorax brevissimus, supra vix conspicuus: mesothoracis scutum transversum, sulcis 2 indistinctis postice approximatis; parapsidium suturæ non bene determinatæ; scutellum obconicum, non productum: metathorax brevi-obconicus, declivis: petiolus crassus, brevissimus: abdomen longiovatum, glabrum, apice acuminatum, thorace paullo longius

non angustius; segmentum 1^{um} maximum: pedes flavi, simplices, subreinales; alæ limpidæ, mediocres; squamulæ picæ; nervi fulvi; nervus subcostalis alæ basi emissus nervulum in discum rejiciens et spatio brevi ante costæ medium in stigma subfurcatum decedens: nervus quoque spurius alæ basi emissus in discum excurrens et nervo subcostali nervuloque triangulum fingens. (Corp. long. lin. $\frac{1}{2}$; alar. lin. 1.)

Romilius (n. g. *) Zotalæ, Fem. *Ater, antennæ nigræ basi fulvæ, pedes fulvi, alæ limpidæ.*

Corpus longum, angustum, sublineare, atrum, vix convexum, obscurum, pilis canis pubescens, subtilissime et confertissime punctatum: caput transversum, breve, subrotundum, thorace vix latius; vertex latus; frons convexa, non impressa: oculi parvi, non extantes: antennæ nigræ, graciles, subclavatæ, prope os insertæ, thorace non breviores; articulus 1^{us} longus, subfusiformis, fulvus; 2^{us} longicyathiformis; 3^{us} et 4^{us} longi, lineares; 5^{us} et sequentes ad 10^{um} breves, approximati, clavam fingentes longifusiformem: thorax longiovatus: prothorax brevissimus, postice excavatus: mesothoracis scutum magnum, trisulcatum, longitudine vix latius; sulci laterales postice approximati; scutellum parvum, semicirculum fingens: metathorax mediocris, obconicus: petiolus brevissimus: abdomen longifusiforme, subtus convexum, apice acuminatum, thorace multo longius; segmentum 1^{um} breve; 2^{um} et 3^{um} magna; 4^{um} et 5^{um} paullo breviora; 6^{um} adhuc brevius: pedes fulvi, longi, graciles, simplices, subreinales; tarsi articulus 1^{us} longus, 2^{us} multo brevior, 3^{us} adhuc brevior, 4^{us} 3^o brevior, 5^{us} 4^o paullo longior: alæ mediocres, limpidæ, abdominis apicem dum quietem agunt non attingentes; squamulæ fulvæ; nervi fulvi; nervus subcostalis alæ basi emissus ad costæ medium eam attingens et fere ad alæ apicem percurrans; nervus 2^{us} quoque alæ basi emissus in discum excurrens ibique furcatus, furca antica nervo subcostali triangulum fingens, furca postica ad alæ marginem posticum decedens; nervus cubitalis rectus, sat longus, nervo subcostali ubi costam attingit emissus, stigmate terminatus minuto. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. $2\frac{1}{4}$.)

[To be continued.]

XL.—*The Crustacea of Ireland. Order Decapoda.* By Wm. THOMPSON, Esq., Vice-Pres. Nat. Hist. Society of Belfast.

CRUSTACEA. Order DECAPODA.

DECAPODA BRACHYURI.

Macropodia phalangium, Leach, Mal. pl. 23. f. 6; Desmar. p. 155. pl. 23. f. 3.

Stenorhynchus phalangium, Edw. Crust. tome i. p. 279.

Cancer phalangium, Penn. Brit. Zool. vol. iv. p. 11. pl. 9. f. 3, edition 1812.

This species has already been recorded by Templeton as "not uncommon on the Irish coast," and by Mr. J. V. Thompson as "very abundant in the deep water of the harbour of Cove." Ent. Mag. vol. iii. p. 371.

It has occurred very commonly to my scientific friends and myself when dredging in the loughs of Strangford and Belfast; and to Mr.

- *Scelio Duris* (Monogr. Chalcid., ii. 61.) also belongs to this genus.