of the planet is very beautiful; but my experience is that on these rare nights one will see fewer strange phenomena about the Ring and the shadows than when the images are blurred and indistinct⁴.

Even if I could have been so greatly deceived as to represent for realities the deformations undergone by images in rapid vibrations, I am pretty certain that I could not have seen the delicate dark angular forms which I have represented, but rather rounded, ill-defined forms totally wanting in the blackness and sharpness of those which I saw. Contrary to Professor Hall's suggestion, it is precisely when the definition was the most perfect that the „strange phenomena“ could be seen with more distinctness, and at the moment the image became tremulous in the least, it disappeared confounded with the dark division of the rings.

The fact that Professor Hall has not been able to see the „Pencil line“, even during one of these beautiful nights he speaks of, sufficiently indicates his failure to see the jagged appearance of the principal division; and indeed, he could not have expected to see it, as these forms are almost as difficult to make out as the grayish line of the outer ring.

I have no positive evidence that these markings continued visible after the end of September, 1875; as, after that time, I discontinued somewhat my observations on Saturn, looking at it only occasionally, until the present year, during which I have observed it on every possible occasion. But, of course, the obliquity of the Ring is too great now to allow the observation of such delicate forms, although I still continue to see the principal division on the ansae. It is not impossible that the obliquity of the Ring was the cause of the failure of Professor

Hall to verify my observations, or the phenomenon may be a temporary one, and it may have been absent when he made his observations.

The phenomenon of the jagged border of the principal division, as I have represented it, was seen so often and with such distinctness in 1872, when the Ring was wide open, that it was impossible for me to doubt its reality; and, besides, it was verified at least on two occasions by Professor Winlock, the late director of the Harvard Observatory, who once was accompanied by Mr. Miliken, manager of the Western Union Telegraph Company, who also saw the dark angular forms on the following ansa.

Professor Hall seems to be in doubt as to the reality of the anomalous curvature of the shadow of the planet on the Ring, and appears inclined to attribute this appearance to some illusion caused by the varied conditions of our atmosphere. In reply, I will remark that, if such was the case, how could we explain its long duration as concave, and its no shorter duration as convex, which has been alternatively observed since the time of Cassini?

Several years ago, a very distinguished and industrious observer, F. Angelo Secchi of the Roman Observatory, pointed out that the deformation of the shadow of the Ball on the Ring was the natural consequence of the unevenness of the surface receiving it. If this is the true explanation, as I think it is, the natural consequence, as derived from the observations, is that the form of the surface is not permanent, since the shadow has evidently shown different outlines; appearing at different times either as a straight, a convex, a concave or a notched line.

Cambridge, Oct. 5, 1877.

Correction of the Elements of Antigone (129).

By *E. P. Austin*.

From the observations of 1873 — 1876 the following normals have been deduced:

Nr.	Date	α	δ	λ	β
I	1873 Febr. 20	9 ^h 4 ^m 24 ^s 33	+ 17° 23' 5" 1	133° 26' 50" 7	+ 0° 36' 49" 4
II	May 24	8. 49. 27. 12	20. 14. 6. 7	129. 15. 19. 5	2. 22. 9. 3
III	April 21	8. 57. 10. 17	20. 50. 58. 8	130. 49. 38. 0	3. 27. 16. 0
IV	May 23	9. 25. 53. 68	+ 19. 41. 1. 9	137. 37. 20. 6	4. 19. 59. 5
V	1874 June 29	18. 19. 32. 44	— 9. 4. 0. 3	274. 58. 43. 4	+ 14. 18. 15. 5
VI	1875 Oct. 7	1. 24. 1. 85	— 7. 15. 23. 2	16. 35. 26. 0	— 14. 55. 29. 8
VII	1876 Dec. 14	5. 6. 3. 47	+ 8. 5. 52. 1	76. 11. 21. 2	— 14. 42. 46. 0

The above are for Berlin midnight and give the true rightascensions and declinations and the corresponding longitudes and latitudes referred to the mean equinox of the beginning of the respective years.

The place for 1874 June 29 depends on a single meridian observation at Berlin for that date, all the others are derived from a considerable number of observations.

With the following elements I have computed the perturbations by Jupiter and Saturn which are given in the annexed table.

Elements I

Osculating for 1873 Febr. 14.

$$\begin{aligned} \pi - \Omega & 102^{\circ} 50' 55'' 96 \\ \Omega & 138. 0.21.02 \\ i & 12.10.54.62 \end{aligned} \left. \vphantom{\begin{aligned} \pi - \Omega \\ \Omega \\ i \end{aligned}} \right\} \text{Mean Equ. 1873.0}$$

$$\begin{aligned} \mu & 727'' 88557 \\ \varphi & 11^{\circ} 58' 2'' 98 \\ M_0 & 281.38.51.37 \text{ Febr. 14.0 Ber. M. t.} \end{aligned}$$

Perturbations.

	Δi	$\Delta \Omega$	$\Delta \varphi$	$\Delta \pi$	$\Delta \mu$	ΔM
I	— 0.18 0.00	— 0.04 0.00	— 4.03 — 0.10	— 7.46 + 0.24	— 0.00315 + 0.00030	— 0 55 — 0.30
II	— 1.20 + 0.02	— 0.50 + 0.01	— 25.49 — 0.19	— 38.60 + 1.55	— 0.00563 + 0.00200	— 12.22 — 1.90
III	— 2.38 + 0.04	— 1.63 + 0.03	— 43.00 — 0.32	— 43.70 + 2.71	— 0.02600 + 0.00358	— 40 26 — 3.23
IV	— 3.80 + 0.06	— 3.85 + 0.06	— 62.68 — 0.61	— 29.10 + 4.12	— 0.09440 + 0.00556	— 88.49 — 4.61
V	— 9.21 + 0.13	— 32.14 + 0.07	— 183.00 + 0.28	+ 294.48 + 14.91	+ 0.87260 + 0.00455	— 212.99 — 9.72
VI	— 16.69 + 0.13	— 48.84 + 0.07	— 162.00 + 7.10	+ 648.86 — 8.44	+ 0.45064 + 0.01580	— 289.01 + 5.61
VII	— 16.64 + 0.31	— 117.70 + 0.12	— 168.61 + 7.00	+ 918.04 — 26.09	— 0.33890 + 0.02310	— 514.98 + 39.13

In the above table the upper number gives the perturbation by Jupiter, the lower that by Saturn from 1873 Febr. 14, to the dates of the respective normals.

With the above elements and perturbations the latitudes and longitudes have been calculated for dates of the several normals and the residuals are given in the following table marked (I).

The corrections of the elements have been derived by given equal weight to the several normals.

$$\begin{aligned} \Delta \mu & = + 0'' 020803 & \Delta \varphi & = - 6'' 35 & \Delta \Omega & = + 3'' 18 \\ \Delta M_0 & = + 14.60 & \Delta i & = + 0.87 & \Delta \pi & = - 26.94 \end{aligned}$$

giving the corrected Elements II.

$$\begin{aligned} \pi - \Omega & 102.50.25.84 \\ \Omega & 138. 0.24.20 \\ i & 12.10.55.49 \\ \mu & 727.906373 \\ \varphi & 11.57.56.63 \\ M_0 & 281.39. 5.97 \end{aligned}$$

Referred to same Equinox osculating for same dates as Elements (I).

With these elements the places have been recomputed giving residuals II agreeing closely with the results of substituting in the equation of condition.

I Residuals II.

C—0

C—0

	$\Delta \lambda$	ΔB	$\Delta \lambda$	ΔB
I	1'' 9	+ 3'' 4	+ 1.2	+ 2.6
II	— 0.7	+ 3.0	— 1.1	+ 2.9
III	+ 0.5	+ 0.8	— 1.8	+ 0.1
IV	+ 0.6	+ 1.0	+ 2.7	+ 2.2
V	— 2.3	+ 6.4	+ 0.5	+ 6.7
VI	+ 12.2	+ 4.1	+ 0.8	+ 4.4
VII	— 8.8	+ 5.0	— 1.3	+ 3.2

The preceding residuals indicate an error in the latitude of Normal V which as already observed depends on a single observation but as the correction of the elements must now be quite small. I have not thought

it necessary to compute new corrections by excluding that normal.

A Computation with the corrected elements gives the following corrections to my ephemeris for the opposition 1878.

1878 Jan. 27	Corr. of α	Corr. of δ
Febr. 12	+ 2 ^m 3 ^s 8	— 4' 57''.
Febr. 28	2.11.4	— 4.36
	+ 2.12.4	— 3.39
Washington. 1878 Jan. 16.		

Beobachtungen auf der Sternwarte zu Athen.

Von Dr. J. F. J. Schmidt.

Veränderliche Sterne. 1877.

7. β Lyrae. 1877.

Hauptminima.			Nebenminima.			Hauptminima.			Nebenminima.		
Mai	10.12	p = 4	Mai	4.10	p = 3	Sept.	16.35	p = 3	Sept.	10.00	p = 4
"	23.22	4	"	17.00	4	Octbr.	11.55	1	"	22.70	4
Juni	4.75	4	"	29.05	4	"	24.80	3	Octobr.	5.87	3
"	17.50	4	Juni	10.87	4	Nov.	6.75	1	"	18.25	3
"	30.90	4	"	24.37	4	Dec.	2.55	1	"	31.75	1
Juli	13.62	4	Juli	6.75	4	"	28.40	2	Nov.	13.40	3
"	26.65	4	"	20.50	4				"	26.45	3
Aug.	8.37	4	Aug.	1.90	4				Dec.	10.10	3
"	21.45	4	"	15.10	4				"	22.60	1
Sept.	3.33	4	"	28.00	4						

8. δ Cephei. 1877.

			Maxima.					
Jan.	5.65	p = 4	Mai	30.55	p = 2	Sept.	14.55	p = 4
"	10.75	3	Juni	4.77	3	"	19.67	4
"	16.00	3	"	9.85	3	"	25.50	3
"	21.60	3	"	14.77	3	"	30.70	2
"	26.70	4	"	20.87	3	Octbr.	5.80	1
Febr.	11.77	4	"	25.95	3	"	11.67	2
"	17.33	3	Juli	1.82	4	"	16.75	4
"	22.35	1	"	6.95	3	"	22.70	3
"	28.37	2	"	12.30	3	Nov.	1.60	1
März	5.70	1	"	17.75	4	"	7.50	2
"	10.85	1	"	23.00	3	"	12.75	4
"	16.75	1	"	29.00	3	"	18.10	0.5
"	26.90	1	Aug.	2.70	4	"	23.50	4
April	12.00	1	"	7.90	4	"	28.62	4
"	17.20	2	"	13.70	4	Dec.	3.87	4
Mai	3.90	1	"	18.75	4	"	14.87	1
"	10.20	2	"	24.00	4	"	20.30	0.5
"	14.00	2	"	29.80	4	"	25.67	4
"	19.80	1	Sept.	3.75	4	"	30.70	4
"	25.10	1	"	9.00	4			

Für den Athener Horizont liegt δ Cephei in den Abendstunden des März, April und Mai so tief, dass genaue Beobachtungen in dieser Zeit nicht zu erwarten sind.