REPORT OF A CASE OF ABDOMINAL ANEURISM TREATED BY WIRING.*

MURAT WILLIS, M.D.

(Adjunct Professor of Abdominal Surgery in the Medical College of Virginia; Attending Surgeon to the Memorial Hospital Richmond, Va.)

The cure of aneurism affecting the larger blood vessels of the body is a question that has given much concern to medical men for many years. In attempting to formulate a rational treatment for this serious condition, surgeons have been particularly active, and by some it is classified as a distinctly surgical disease. A study of the literature, however, is not very encouraging, and, in the light of our present knowledge, it must be confessed that the rational treatment of this malady, either surgical or medical, has by no means been attained.

It is the purpose of this paper to place on record one case of aneurism of the abdominal aorta, treated by the insertion of a wire through which a galvanic current was passed after the wire had been introduced. At the same time attention is called to the previous similar operations, which, as collected from the literature, are twenty in number. A critical analysis of these twenty-one cases brings out several facts from which important conclusions may be deduced.

The case which I wish to report was a negro, 32 years of age, whose occupation was that of a Pullman car porter. He was referred to the surgical service of Dr. Geo. B. Johnston from the practice of Dr. E. A. Truman, Richmond, Va. The patient had enjoyed perfect health until June, 1907, and denied all history of lues. At that time he began to suffer with occasional attacks of diarrhœa, which have persisted up to the present time in spite of treatment. He now has from five to ten stools a day, which are watery, offensive, con-

tain a slight amount of mucus and blood, but there is no tenesmus. Three months ago he began to suffer with severe pain in the region of the umbilicus, with frequent colicky exacerbations. This lasted for nearly a month but has not since returned. Following this he noted considerable throbbing in the abdomen. Physical examination showed a distinctly emaciated patient who was moderately anemic. The heart and lungs were clear throughout on percussion and auscultation. In the abdomen there was a visible prominence in the right hypochondrium. On palpation over this area there was a long rough systolic thrill, which on auscultation gave rise to a bruit that was so loud as to almost hurt the ears. The tumor appeared to be the size of a small cocoanut and was situated just above and to the right of the navel. There was very marked expansile pulsation. A distinct diastolic shock could be made out in addition to the systolic bruit. The abdomen was elsewhere negative. Both femoral pulses were distinctly felt and were synchronous in time. An examination of a specimen of the fresh stool showed that it was liquid with small yellow masses like milk curds and two small particles of dark red coagulated blood with a slight amount of mucus. Microscopical examination was negative for amebæ and other parasites.

Patient was operated upon March 28, 1908, under ether anesthesia by the drop method. A median incision was made and exposed the tumor, which appeared to arise from the aorta just below the celiac axis. The tumor was

*Read before the Southern Surgical and Gynecological Association, St. Louis, Mo., December, 1908.

distinctly sacculated and came off on the right side of the aorta. A needle, covered with lacquer, was inserted into the sac with its point away from the aorta and through this eleven feet of silver wire was fed to form a coil. Following this the electric current was passed as follows:

- 10 milliamperes for 20 minutes.
- 20 milliamperes for 10 minutes.
- 30 milliamperes for 10 minutes.
- 40 milliamperes for about 5 minutes.

When this had been raised to 40 milliamperes the patient's condition suddenly changed, the pulse went up to 140 and it was necessary to give artificial respiration for several minutes. He gradually reacted, however, and the needle was removed without hemorrhage and the patient was put back to bed. Recovery was without incident.

The notes of a physical examination three weeks after operation showed, as we thought, a tumor of distinctly smaller size with less marked pulsation, and patient stated that the pain was very much less. One month after operation he left the hospital and two weeks after this he walked sixteen squares to present himself to the clinic. At this time he complained that the diarrhoea from which he suffered previous to operation was again giving him trouble. On account of this he was unable to go back to his work as Pullman car porter, but took up a lighter occupation. Three months after operation the diarrhoea increased in severity and he had repeated hemorrhage from the bowels and died forty-eight hours after the bleeding commenced. It was impossible to obtain a complete autopsy or to get the relatives to consent to our removing the tumor. A partial autopsy, made through an incision at the site of the original wound, showed the abdominal cavity entirely clean and free from any evidence of either hemorrhage or infection, and we believe that the sac ruptured into the bowel. On palpating the sac the coils

of wire could be very distinctly felt through the thin walls and there was absolutely no evidence of any clot formation. Although before death we were satisfied that the tumor was reduced in size and that there was less pulsation, still at the autopsy there was no evidence to support such a view. The wall of the sac could be easily compressed between the fingers and would evidently have collapsed were it not for the framework of wire.

The first attempt to cure an aneurism by the insertion of a wire was made in 1864 by Dr. Moore of the Middlesex Hospital. He inserted seventy-eight feet of wire into a thoracic aneurism on the theory that the first indispensable condition for the cure of aneurism is to provide a means of eliciting fibrin from the blood, and a second like condition is to extend the surface within the sac on which the fibrin may be deposited. This method was not applied in the case of an abdominal aneurism until twenty years later, when Loreta, in 1884, introduced six feet of fine wire through a trochar. In this case there appears to have been an almost instant relief of symptoms and a progressive improvement for ten weeks succeeding operation. Thirteen weeks after the sac was wired, however, the patient suddenly died from rupture of the aorta into the abdominal cavity.

A distinct improvement over the early methods was the suggestion by Corradi of passing a galvanic current through the wire after it had been introduced. This was first applied by Corradi himself in the case of a large aneurism of the ascending aorta, April, 1879. It was not until 1892, however, that the Moore-Corradi method was employed for the cure of abdominal aneurism. This operation was performed by Stewart of Philadelphia.

SUMMARY.

For the reason that aneurism of the abdominal aorta has proved absolutely incurable by internal medication, it may be regarded as one

of those conditions in which a wide range of experimentation and trial methods is justified. Much has been hoped for in the Moore-Corradi method of treatment by wiring and electrolysis. Notwithstanding the fact that a technic of this operation has been perfected through most elaborate and painstaking studies and experiments, it would seem that those cases of aneurism of the abdominal aorta in which it has been applied have terminated in disappointment to the operator.

A critical study of the twenty-one cases thus far reported inevitably leads to the conclusion that surgery so far offers no certain relief for this malady. Of these twenty-one cases, the subsequent history of three is not recorded; in the remaining eighteen, death ensued in every case. The duration of life after operation in this series varied from one hour to nine months with an average duration of less than fifty days. Who knows that the length of life may not have been shortened instead of prolonged by this method of treatment? The operative technic that has developed for this condition would seem to be absolutely perfect and based upon sound surgical principles, but the fact remains that this method of treatment is disappointing and the last word has not yet been said concerning the surgical treatment of abdominal aneurism.

BIBLIOGRAPHY.

1. Loreta. 1885. "Surgical Treatment of Abdominal Aneurism." (Editorial abstract of report in Memorial Royal Academy de Bologna, 4 S., Vol. 6). *British Medical Journal*, London. Vol. 1, April 11, pages 745-746; May 9, page 965.
2. Lange. 1886. "Aneurism of the Abdominal Aorta, Treated by the Insertion of Wire Into the Sac." *Medical News*, Philadelphia; Vol. 49, pages 582-583. Also *British Medical Journal*, November 20; Vol. 2.
3. Morse. 1887. "Cure of Abdominal Aneurism by Loreta's Method." *Pacific Medi-*

cal and Surgical Journal, San Francisco; Vol. 30; February, pages 65-67.

4. Pringle, J. J. 1887. "A Case of Aneurism of the Abdominal Aorta Treated by Laparotomy and the Introduction of Steel Wire Into the Sac." *Med.-Chir., Tr.*, London, Vol. 70, pages 261-276.

5. Stevenson. 1895. "A Case of Abdominal Aneurism Treated by Laparotomy and the Introduction of Wire Into the Sac; Death." *Lancet*, London, Vol. 1, pages 22-23.

6. Langton. 1899. "A Case of Aneurism of the Abdominal Aorta Successfully Treated by the Introduction of Silver Wire into the Sac." *Tr. Clinic Society*, London, Vol. 32, pages 138-141.

7. Stewart. 1892. "Treatment of Sacculated Aortic Aneurism by Electrolysis Through Introduced Wire." *American Journal Medical Science*, Philadelphia. N. S., Vol. 104; pages 422-438.

8. Noble. 1898. "Aneurism of the Abdominal Aorta Cured by the Introduction of Gold Wire and Galvanism." *Philadelphia Medical Journal*; Vol. 1; pages 1203-1205.

9. Reeve. 1899. "On the Proper Wire for Introduction into an Aneurismal Sac." *Ann. Surg.*, Philadelphia; Vol. 30, pages 704-706.

10. Stewart. 1898. "Further Account of the Treatment of Aneurism by the Conjoint Use of Galvanism Through Introduced Coiled Wire." *Report of cases. Philadelphia Medical Journal*; Vol. 2; pages 1011-1019.

11. Parham. 1901. "A Case of Abdominal Aneurism Treated by Abdominal Section and Introduction of Wire." *American Journal Surg. and Gynec.*, St. Louis; Vol. 14; pages 193-194.

12. Maunsell. 1904. "The Surgical Treatment of Abdominal Aneurism." *Tr. Royal Academy of Medicine, Ireland, Dublin*, Vol. 22, pages 109-126.

13. Holt. 1904. "Aneurism of Abdominal Aorta and Coelic Axis; Attempted Cure by Power and Colt's Wire Apparatus." *Death*

from Double Pneumonia. J. Roy. Army Med. Corp., London; Vol. 3; pages 175-178.

14. Matas. 1901. "The Treatment of Abdominal Aortic Aneurism by a Preliminary Exploratory Celiotomy and Peritoneal Exclusion of the Sac. Followed at a Later Sitting by Wiring and Electrolysis, with the Report of Two Hitherto Unpublished Cases." Am. Med., Phila., Vol. 1; pages 546-589. Also Tr. Southern Surg and Gynec. Association; Vol. 13; pages 272-330.

15. Hunner. 1900. "Aneurism of the Aorta Treated by the Insertion of a Permanent Wire and Galvanism (Moore-Corradi method.) Johns Hopkins Hospital Bulletin, Baltimore; Vol. 11; pages 265-279.

16. Power and Colt, 1903. "A Case of Aneurism of the Abdominal Aorta Treated by

the Introduction of Silver Wire, with a Description of Instruments Invented and Constructed by Mr. G. H. Colt to Facilitate the Introduction of Wire Into Aneurisms." Med. Press and Circ., London, N. S.; Vol. 76; pages 161-164. Also Lancet, London, Vol. 2; pages 808-813. 1903.

17. Eastman. 1906. "Aortic Aneurism Treated by Introduction of Fifteen Feet of Silver Wire." Death after four months. New York Medical Journal; Vol. 84; pages 1170-1172.

18. Griffith. 1905. "A Case of Sacculated Aneurism of the Abdominal Aorta, Treated by the Introduction of Silver Wire and the Passage of a Constant Current." Lancet, London; Vol. 2; August 12; pages 442-444.