PASTEURIZATION.

BY M. J. ROSENAU, M.D.

Director Hygiene Laboratory, U. S. P. H. and M.-H. S.

Advantages.—The advantage of pasteurization is that it is a cheap and effective means of preventing the transmission of infectious diseases such as tuberculosis, typhoid fever, diphtheria, scarlet fever, etc., commonly spread by milk.

Disadvantages.—1. Pasteurization promotes carelessness on the farm and dairy, etc. (This may be controlled by proper regulations, inspections and laboratory examinations.)

2. Pasteurization renders milk less digestible. (While it is generally conceded that boiled milk commonly induces constipation, the majority of the evidence plainly indicates that pasteurization has little, if any, effect on the digestibility of the milk.)

3. Pasteurized milk favors the production of rickets and scurvy. (There is no proof to this effect and authorities agree that the danger is slight; and, further, that it may readily be obviated.)

4. By destroying the non-spore-bearing bacteria, pasteurization sometimes allows toxic organisms to grow and produce serious poisons in the milk. (On the other hand, these same poisons are more frequently produced in milk that has not been pasteurized, and this danger may be obviated in pasteurized milk by cooling it quickly, keeping it cold and shortening the time for distribution.)

5. Pasteurization is inefficient as a preservative; the milk keeps only 12 to 24 hours longer than otherwise. (This is really no disadvantage, for the quicker bad milk sours, the better.)

6. Pasteurization injures the taste of the milk. (This is not so, if properly done.)

7. Pasteurization increases the cost of the milk. (True, but it is the cheapest safeguard, and the expense of pasteurization is offset by the keeping quality of the milk.)

Methods.—The milk must be heated to a temperature of 65 C. (149 F.) for 20 minutes, or 70 C. (158 F.) for 10 minutes, as soon as practicable after the milking, in enclosed vessels, preferably the final containers. After pasteurization, the milk must be promptly cooled, kept cold and distributed with promptness.

The advantages so far outweigh the disadvantages that I unhesitatingly recommend compulsory pasteurization of all milk not certified under Class 1 and Class 2 of Dr. Melvin's classification.

DISCUSSION

ON PAPERS BY DRS. GOLER, DARLINGTON, JORDAN AND MAGRUDER.

DR. G. LLOYD MAGRUDER, Washington, D. C., said that Washington holds a unique position in regard to government and is somewhat backward in some points in which it should be forward. It is governed by a board of commissioners, but the laws are enacted by Congress and it is necessary to go to Congress for almost everything. The commissioners have power over certain police regulations, but they can not control without legislation from Congress. The District of Columbia is very close to Maryland and Virginia, and the congressional committees on District of Columbia affairs frequently have members from these states who are influenced by their constituents. For some time, Dr. Magruder said, a senator, who had a dairy farm in Virginia, was on one of these committees. Almost all the legislation which it was attempted to secure for stricter regulations of the milk supply of the District was opposed by this man, because he had trouble in carrying out the requirements on his farm while he lived in Washington. Health officers and commissioners, year after year since 1895, have begged for additional legislation; in 1895 legislation was secured which improved conditions immensely, but it was not sufficiently strong to enable the health officers to do what they should. They are now working for additional legislation.

DR. C. H. JONES, Baltimore, said that the papers on this subject have given him an idea as to what can be accomplished

under bad conditions. He stated that at the Scientific Exhibit Hall he had presented some pictures that were taken for the milk exhibit gotten up by Dr. Fulton, secretary of the State Board of Health of Maryland, in the spring of 1906. There are about 108 photographs presenting the conditions within the city, in the suburban district and in the rural district. So far as he has been able to learn, the greatest difficulty is found in the suburban district. Within the city there is a fairly strict supervision—so far as one man can carry it out. The improvement in the milk has been very great; practically no disease except that introduced in the milk in houses, could be traced to these particular dairies. The rural district is far better than what one would expect, as there is no supervision; but Dr. Jones thinks that is accounted for in the following way: The distance that the milk has to come from the stables to the point of distribution within the city, and the fact that the railroads do not provide any means of conveying the milk at the proper temperature compel the farmers to exercise a certain amount of care, as otherwise they know that they can not get the milk to the city and have it remain sweet. On the other hand, the suburban section is a district close to the city where the milk is brought twice daily to the consumers in wagons, not by the railroads. Here is found the greatest amount of filth. The short haul permits the milk to be consumed within three or four hours after milking. Dr. Jones thinks that he can demonstrate a typhoid epidemic, due entirely to milk infection, in the suburban district—not the urban district nor the rural.

DR. W. F. SNOW, Leland Stanford University, Cal., said that California is carrying out three experiments which seem pertinent to the subject. All the speakers have emphasized four things: First, the education of the public through the press; second, the necessity for the establishment of demonstration dairies and experimental farms within public parks, or in similar ways; third, the emphasizing before the public of pure production of milk rather than pasteurization; and fourth, the necessity for field inspection work. In Alameda County, California, the supervisors have established a model dairy in conjunction with their county infirmary. They are spending considerable money in making it what a dairy should be, with the object of bringing the people from the surrounding cities (those near the Bay of San Francisco) to see the place. They are experimenting with a milking machine for the benefit of the public, and are using the milk also in an experimental way (so far as good milk may be an experiment) with the patients in this infirmary, to see whether good milk will show good results in the dietetics of this hospital. Further, in regard to the question of education itself, Dr. Snow has experimented somewhat with some of the high schools in putting before the teachers of biology such blanks and instructions as have been shown in the Section, asking these teachers to go with him, and later, with their students to the dairies supplying their communities with milk. The high school students are then asked to write, in their course on biology, papers on their impressions with regard to how fully these rules and regulations are complied with. The effect has been rather amusing, and very encouraging from the point of view of throwing a lot of amateur inspectors into the field; amusing in its influence on the dairymen, encouraging in the public sentiment which it has created in the homes of these school children. Within the last year and a half four high schools (those of Los Angeles, Stockton and some other communities), representing probably 5,000 homes, have decided to put education in public health into the regular school curriculum. Just what they are going to accomplish, or how persistently they will carry it out, no one knows, but at any rate the scheme of instruction in these four high schools includes some sort of simple field-inspection training in examining the sanitary conditions of dairies and other food-production centers. The third experiment of which Dr. Snow spoke is one which seems valuable to those who are in states which have had very little money to draw on for this public work as yet. The United States Government utilizes the best students in the zoologic, biologic and geologic departments of the universities for all sorts of valuable experimental field work. The health departments of the

various states could use the students in biology and physiology fully as well for very careful volunteer inspection work in the field of public health. Merely as an experiment, the State Department of Health in California has been trying such a plan with the students from one of the universities, with a very great deal of value from two points: First, that of supplying the overburdened officers of the health department with accurate, trustworthy facts; second, that of educating many of the young men who will soon be physicians, and, more important still, Dr. Snow said, our voters and trustees or lawyers in incorporated communities, both large and small.

DR. SENECA EGBERT, Philadelphia, said that one important point would be the requirement that milk cans and containers in transit on railroads should be sealed. Under the auspices of the State Commissioner of Health in Pennsylvania, it was found that often the train employes made free with milk in transit; and to get it they used anything convenient, such as a tin cup or glass or the tumbler from the cooler in the caboose, or sometimes the lid of the milk can itself. Another thing that the cities can do, he said, is to insist on far better sanitary conditions in the milk shops and at the distribution points. He also thinks that it would be wise for each city to have one or more model sanitary milk shops so that the public might have a practical example of what such shops should be, and thus might learn to patronize only those of that character. Another idea which has occurred to him as feasible, is that the milk should be delivered to the consumer in an absolutely frozen condition. It could be frozen at the dairy and delivered to the consumer as is ice cream, either in bulk or in blocks, enclosed in paper cartons. Investigations have shown slight changes in the composition and digestibility of frozen milk, but it is doubtful, he said, if these changes are any greater than those which occur when milk is pasteurized. It seems to Dr. Egbert that at very slight additional expense such milk could be supplied and people could be educated to look for milk in the solid form. This would prevent the bacterial growth that undoubtedly occurs if milk is kept at any temperature above 32 F. Newman and others positively state, he said, that 32 F. is the maximum temperature at which the bacteria will not increase in milk, and the plan suggested would prevent any increase from the time the milk leaves the refrigerating plant until it reaches the consumer. Dr. Egbert is inclined to feel that in the ordinary dairy pasteurization it is not a safe thing for the consumer. Pasteurization does not kill all the bacteria; after that process and after the milk reaches a temperature of say 60 F., the bacteria may grow and be just as numerous as ever before and the harm done may be just as great as if the milk had not been pasteurized. The ordinary layman, if he gets pasteurized milk, thinks it is free from germs, which is not the case. Milk refrigerated (as it should be) immediately after the milking and kept cooled down to at least 30 F., will be a much safer article to supply to the ordinary consumer than the pasteurized milk.

DR. H. M. BRACKEN, Minneapolis, said that in small communities it is discouraging to try to get people interested in such matters. In Minnesota there are no very large cities where the problem of securing a safe milk supply is a difficult one. He declared that a city with a population of 300,000 or less, and a good farming community around it, should have no difficulty in securing a good milk supply, and that cities with populations of 50,000 or less should have no difficulty in keeping the milk standard near that of certified milk. He referred to the papers and illustrations showing the variations in the quality of milk. The standard of Boston for commercial milk—500,000 bacteria per c.c.—seems liberal enough, yet it appears to be difficult for Boston and New York City to keep up to that standard. The inspected milk of New York (100,000) is rather a high grade milk. In Minnesota an effort was made through the State Board of Health to encourage smaller cities to insist on a better milk supply; and the board, while it is not in the position to give mandatory directions on such matters, did recommend that all the smaller cities should fix a standard equivalent to that of Boston, viz., 500,000 bacteria per c.c. Some of the smaller cities are now taking the matter up and by ordinance are adopting this as

a standard. When this standard becomes somewhat general, it will then be possible to move for a still higher standard. He said that if the milk is properly handled it should not be necessary to freeze it to keep it. Certified milk can be kept for some time, and Dr. Bracken thinks that certified milk is certainly superior to milk reduced to a frozen state.

DR. G. LLOYD MAGRUDER, Washington, D. C., said that as a result of this agitation in Washington, one dealer has tried hard to comply with the requirements for inspected milk. He voluntarily offered the producers an additional 3 or 4 cents a gallon for the purpose of getting clean tuberculin tested milk. He has spent \$500 in advertising. Before Dr. Magruder left Washington, these advertisements had been going on for ten days, and the dealer said that he had had ten applications for this higher grade of milk—tuberculin-tested and low in bacteria—for which he wanted to charge 10 cents a quart. Dr. Magruder saw Mr. Gordon, the representative of the Walker-Gordon establishment in Washington, the day before he left Washington. He has a depot directly opposite Dr. Magruder's office, where he puts out 1,600 quarts a day. Mr. Gordon says that most of his customers are people coming to Washington on the recommendation of the plants elsewhere, and he can not get over 10 cents a quart for his milk in Washington; and that if he raises his price to 12 cents in September he is sure that he will lose more than one-third of his customers. People want milk at 6 and 8 cents a quart.

SOME ADVANCES IN RENAL AND URETERAL DIAGNOSIS.*

CHARLES LESTER LEONARD, A.M., M.D.

PHILADELPHIA.

The uncertainties that frequently surrounded the diagnosis of lesions in the deeper urinary tract have been materially lessened by recent advances in the physical methods of diagnosis. This is particularly true of the intrarenal and intraureteral disorders, where improvements in diagnosis have resulted in a clearer differentiation between the symptoms of conditions that had been grouped as identical; in a more exact determination of the character and location of the lesion; and hence in a more accurate analysis of the indications for treatment, resulting in valuable modifications, more precise operative intervention, and, finally, a marked betterment in results.

The facts that have led to a differentiation in the symptomatology and a more accurate localizing of intrarenal and intraureteral lesions have been derived from the study of the symptoms after an accurate localization of a calculus has been made by the Roentgen method of diagnosis. Thus the localizing symptoms observed as grouped about calculi at different points in the urinary tract have been utilized to locate other than calculous lesions.

These symptoms are not of themselves sufficient on which to base a diagnosis, nor is any method of diagnosis so accurate as to be self-sufficient. The complete study and diagnosis of any renal or ureteral condition can not be made by any one method, and all advance is the composite of advances made in all methods combined. The cystoscope, the ureteral catheter, the wax-tipped bougie, the divided and centrifugalized urines, the clinical history and symptomatology are all important factors in increasing the accuracy of diagnosis. Unfortunately the technic and clinical experience to obtain valuable results by any of these expert methods is as difficult to attain as that demanded of the Roent-

* Read in the Section on Obstetrics and Diseases of Women of the American Medical Association, at the Fifty-eighth Annual Session, June, 1907.