This article was downloaded by: [New York University]
On: 17 February 2015, At: 22:26
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK

Annals and Magazine of Natural History: Series 4

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah10>

The caudal styles of insects sense-organs, i. e. Abdominal antennæ

Dr. A.S. Packard Jun.

Published online: 16 Oct 2009.

To cite this article: Dr. A.S. Packard Jun. (1871) The caudal styles of insects sense-organs, i. e. Abdominal antennæ , Annals and Magazine of Natural History: Series 4, 7:38, 176-176, DOI: [10.1080/00222937108696345](https://doi.org/10.1080/00222937108696345)

To link to this article: <http://dx.doi.org/10.1080/00222937108696345>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly

or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

The Caudal Styles of Insects Sense-organs, i. e. Abdominal Antennæ.
By Dr. A. S. PACKARD, Jun.

Dr. Anton Dohrn has published a note, in the 'Journal of the Entomological Society of Stettin' (1869), to the effect that the abdominal appendages of the female of the mole-cricket (*Gryllo-talpa*) are true sensory organs (Tastorgane).

In the 'Proceedings of the Boston Society of Natural History,' May, 1866, the writer states that "while, as we have shown above, the genital armour of insects is not homologous with the limbs, there are, however, true jointed appendages attached to the ninth or tenth abdominal ring, or both, which are often antenniform, and serve as sensorio-genital organs in most [many] Neuroptera and Orthoptera" (p. 290).

In the same 'Proceedings' for February 26, 1868, he thus writes: "Regarding the insect as consisting of two (fore and hind) halves, the two ends being, with this view, repetitions of each other, these anal stylets may be considered as abdominal antennæ, so that the antennæ look one way, and their homologues, the many-jointed antenniform anal stylets, the opposite" (p. 398).

The subject is also referred to in the 'Guide to the Study of Insects,' p. 17; and the remarkable antenniform abdominal appendages of *Mantis tessellata* are figured in illustration.

I have been able to detect sense-organs (probably endowed with the sense of smell) in the short, stout, jointed anal stylets of the cockroach (*Periplaneta americana*), beautifully mounted by Mr. E. Bicknell, having recently, after reading Dr. Dohrn's note, observed the sense-organs and counted about ninety* minute sacs on each stylet, which are probably smelling or auditory organs, such as are described by Hicks (see 'Guide,' p. 26). They are much larger and much more numerous than similar sacs in the antennæ of the same insect, and are situated in single rows on the upperside of each joint of the stylets. During the breeding-season a peculiar odour is perhaps emitted by the female, as in vertebrate animals; and it is probable that these caudal appendages are endowed with the sense of smell rather than of hearing, that the male may smell its way to its partner. This is an argument that the broadly pectinated antennæ of many moths are endowed rather with the sense of smelling than of hearing, to enable the males to smell out the females. I have observed the same organs in the lamellæ of the antennæ of the carrion-beetles, which undoubtedly depend more on the sense of smell than that of touch or hearing to find stinking carcasses in which to place their eggs.—*American Naturalist*, vol. iv. Dec. 1870.

* Mr. Bicknell has counted more carefully than I did the exact number of these pits, and made out ninety-five on one stylet and one hundred and two on the other, adding, "there were none on the underside of their appendages that I could see."