ACUTE MENINGITIS TREATED BY DRAINAGE OF THE SPINAL CORD.

By STEPHEN PAGET, F.R.C.S. ENG. &c.,
SURGEON TO THE WEST LONDON AND METROPOLITAN HOSPITALS.

THE relief of tension in meningitis by incision of the dura mater of the cervical cord has been already practised in several instances, and my reason for publishing this isolated case is the hope that similar cases will be recorded, and thus show whether the surgeon may expect, by earlier operation, better success than I obtained. The improvement in the patient in the present case was sufficiently marked to justify the operation, which was done under the advice of Dr. Tooth and Dr. Risien Russell. A year ago I trephined the skull in a similar case, but the slight relief thus obtained was followed by hernia cerebri.

A boy eight years of age was admitted to the Metropolitan Hospital on Sept. 3rd under the care of Dr. Risien Russell. There was a history of phthisis in the family. A fortnight before admission he got wet through, and since then he had complained of pains in the body and lower limbs, with headache and constipation. During the previous week there had been frequent vomiting. On admission to hospital he was thin, pale and drowsy; the skin was hot and dry, the temperature was 102° F.; the pulse was 120; and there was frequent vomiting. On Sept. 4th he was drowsy, and resented being disturbed; he vomited several times during the night and also in the morning. At times he was restless, moaning and crying, but he did not speak. The knee-jerk was abolished, there were slight retraction of the head and occasional strabismus and the pupils were dilated. He vomited several times during the day. On the 5th he vomited in the morning. The cheeks were flushed, the head was much retracted, the left eye turned inwards, and the pupils were widely dilated and immobile. Both optic discs were swollen and blurred; he appeared to be blind, and he was speechless and unconscious, passing urine and faeces under him. On the 6th there was no improvement and he was much wasted. The next day I exposed the spine according to Horsley's method, and removed the arches of the fourth and fifth cervical vertebrae. The dura mater was very tense and did not pulsate; when it was punctured the fluid spurted up with great force in a high jet till four or five ounces had escaped. The dura mater was now freely opened, a horse-hair drain was put in, and the wound was closed round it. It was observed that the pulse gained in strength when the fluid was evacuated, but the pupils did not change. On the 8th there was marked improvement; the vomiting had stopped and the cheeks were no longer flushed. During the night the pulse had gradually fallen to 118 and the respiration from 40 to 26. He spoke a few words, saying that his neck hurt him, and he put out his tongue when he was told to do so and smiled when he was gently tickled. There was no change in the pupils and he still passed everything under him. On the 9th he had relapsed into his former condition, except that the vomiting had not returned. The cheeks were flushed and he was unconscious and restless, moaning and gesticulating. The wound had drained well and it had been unnecessary to change the dressings daily. He died on the afternoon of Sept. 11th.

A post-mortem examination was made on Sept. 12th. The head and neck alone were examined. The wound was healthy, and the drain still acted. The veins of the dura mater were distended. In the middle cerebral fossæ there was about an ounce of sero-purulent fluid, and in the posterior fossæ there were between two and three ounces of clear serous fluid. The pia mater was slightly infiltrated in a few places. The convolutions were so much flattened that they were almost obliterated: this extreme flattening was well marked over both hemispheres. The brain was firmer than natural to the touch. There was marked dilatation of the third and fourth ventricles; the lateral ventricles contained an excess of fluid, but they were not distended or markedly dilated. The pia mater at the base of the brain was infiltrated with sero-purulent lymph, and showed a few minute tubercles.

Remarks.—The operation gave relief, but only for a day; and the drainage of the cerebro-spinal fluid after the operation did not effect the removal of the collections of fluid in the cranial fossæ or reduce the inflammatory swelling of the brain. Dr. Tooth is of opinion that this general swelling

of the brain was the chief cause of death, and thinks that it might be advisable, in a similar case, to make a very free opening in the vault of the skull after incising the dura mater in the cervical spine.

Wimpole-street, W.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL AND THERAPEUTICAL.

THE EFFECTS OF A WASP'S STING.

By A. IVOR THOMAS, M.R.C.S. ENG., L.R.C.P. LOND.

THE following account of the effects of a wasp's sting is interesting, in so far as it mentions some peculiar features which hitherto appear to be unrecorded.

A man aged fifty-six was stung on the middle finger of the left hand by a wasp at 1 P.M. on Sept. 25th. Immediately on admission to hospital he became faint and experienced severe pains extending up his arm and ultimately all over his body. A severe rigor followed, ending in violent vomiting and diarrhoea, which lasted for about ten minutes. The patient noticed that his tongue and throat were becoming "larger" and he became unable to protrude the tongue or swallow; but there was no difficulty in breathing. I arrived at 1.45 P.M. and found the patient in a cold, collapsed condition, having a thin, wiry pulse, with the temperature 98° F. The finger, hand and forearm were swollen, but not painful; his tongue and tonsils had by this time decreased in size, but were still perceptibly swollen and reddened. The sting had been abstracted from his finger before my arrival by his wife, who stated that it measured about a quarter of an inch long. The patient was ordered to bed and to have a poultice to the finger and half-ounce doses of brandy every two hours. On the 26th his general condition had improved, the temperature being 99.8°, and the tongue and throat had become normal. The forearm was swollen, cedematous and painful. Red streaks indicating lymphangitis were present, the supra-olecranon lymphatic gland being enlarged and tender, but there was no axillary tenderness. The patient noticed, on being undressed for bed on the previous day, that his trunk and legs were covered by a vivid redness—probably erythema—a condition mentioned by certain writers. This redness had disappeared on the morning of the 26th. The urine had diminished in quantity, was acid and high-coloured, and contained large quantities of phosphates. I ordered further poulticing and a diuretic mixture. On the 27th the patient, after passing a good night, had wonderfully improved; the redness and cedema of the forearm had subsided, he was in every way in a comfortable condition, and he was allowed to get up, being convalescent.

EMPHYEMA OPENING INTO A BRONCHUS; RECOVERY.

By EDWARD MACKEY, M.D. LOND., L.R.C.P. EDIN.

THE letter in THE LANCET of Sept. 30th from "T. H. G.," inquiring as to cases of the above kind, induces me to record the following case:—

A boy aged six years was admitted to the Children's Hospital, Brighton, on July 18th last, who was said to have had "inflammation of the side" for five weeks. He had a rapid, feeble pulse, hurried breathing, and moderate cough, the temperature being 100.2° F.; there was dull percussion from the third rib in front, in the axilla, and at the right base, with feeble breath sound and some friction; the heart had an apex beat in the fifth space outside the nipple. On the 25th there was Skodaic resonance at the right apex, and the hypodermic needle on being inserted between the fourth and fifth ribs in the mid-axillary line brought laudable pus; he was referred for incision.—27th: Whilst being anaesthetised he had a violent cough and brought up from two to three ounces of pus, with some risk of asphyxia; the operation was deferred.—29th: He has continued to cough up non-offensive pus, but there are no signs of pneumothorax; the general condition is improved.—Aug. 2nd: The improvement continues, and after I had consulted with my surgical