ADENOIDS.*

W. FREUDENTHAL, M.D.

Consulting Laryngologist to Montefiore Home, Beth Israel Hospital,
and Bedford Sanatorium for Consumptives; Laryngol-
ogist to St. Mark's Hospital, Etc.

NEW YORK.

About seventeen years ago, when I came to this country, after having seen and operated on many cases of adenoids at Berlin and Freiburg, at one of the first meetings of the New York Academy of Medicine which I attended, a prominent colleague made the remark that while there might be found such things as adenoid vegetations among the children in Denmark or in Germany and Russia, the Anglo-Saxon race was free from that anomaly. In the years that have elapsed, I am confident that the gentleman has been convinced that they do exist in England as well as in the United States, and nowadays I have yet to see the practitioner who has not encountered cases of so-called adenoids in his own practice. They have been, so to speak, part of my daily bread for more than eighteen years in clinics and private practice. Still, when your Chairman did me the honor of asking me to read a paper before this Section, I was somewhat embarrassed as to what to say to a body of experienced practitioners that would be new to them on this topic. The libraries abound in articles and pamphlets on adenoids, which for some time have been viewed, not as a curiosity, but as an every-day occurrence. They have become so frequent in all parts of the globe that every practitioner must of necessity be familiar with them.

Adenoids grow in a place which I long ago considered the most important of the whole breathing apparatus, viz., the retropharynx. We can and do examine easily all other parts of the upper respiratory tract, but the retropharynx, which is the *locus minima resistentiæ* for many a disease, is too often overlooked. No wonder that adenoids, which have existed as long as civilization, were overlooked, and discovered only as late as 1868 by Wilhelm Meyer of Copenhagen. It is undoubtedly true, however, that Voltolini removed them already in 1865 with the galvano-cautery without recognizing their importance.

Now, what are adenoids, or, as they are also called, adenoid vegetations, adenoid growths, pharyngeal tonsil, third tonsil or Luschka's¹ tonsil? We had known since the time of Morgagni that the parts in question belonged to the tonsillar tissue, but only through the works of Henle, Kölliker and His have we acquired a thorough microscopic knowledge of these structures. It was my former teacher, Prof. Wilhelm His of Leipzig, who called it adenoid tissue, and since then the name adenoids has become so popular that we can no longer change it. Waldeyer showed that the adenoid tissue in the upper pharynx is only a part of a lymphatic pharyngeal ring ("lymphatischer Rachenring"), which now goes under the name of Waldeyer's ring.

Any part of this ring may undergo acute and chronic changes. Just as the faucial tonsil may become the seat of acute and chronic inflammation, the pharyngeal tonsil may be similarly affected.

ACUTE CATARRHAL INFLAMMATION OF THE PHARYNGEAL TONSIL.

This is called by the French adenoiditis, with which most likely the *fièvre ganglionnaire* is identical. It is

not as infrequent as is ordinarily believed, for, according to my experience, it often occurs in young children. As an examination is difficult or often impossible the diagnosis is very seldom made. A digital exploration should be omitted, if possible, on account of the reactionary inflammation and pain following this procedure. The main symptom of an acute adenoiditis is earache. It is sometimes very severe, and as there is pain in swallowing at the same time, the child refuses all food and runs down quickly. Frequently the child pulls on the ear, and this is taken either as a diagnostic sign of teething or of an acute otitis media. I have examined the ears of such children, but found nothing pathologic there, and the whole process subsided without affecting the ears at all. This, unfortunately, does not always occur, for pus forms not infrequently, as you all know. But this happens mostly after or in connection with an acute adenoiditis.

In other cases the sound of the voice is muffled and the breathing difficult, which again is regarded as "a cold in the head." Here the examination of the nose reveals nothing, except perhaps a slight hyperemia of the mucous membrane. In grown children the pain in swallowing is very characteristic, and resembles very much that seen in cases of tonsillar or peritonsillar abscess. In my cases fever has always been present, sometimes as high as 104 F.

If there already exists a hypertrophy of the adenoid tissue in the vault of the pharynx, then the tendency to acute inflammation is so much the greater, just as with the faucial tonsil.

THE TREATMENT OF ACUTE ADENOIDITIS.

The treatment which has given me the quickest and best results is the following: First evacuate the bowels and put the child in a hot bath with some mustard in it. Then give small doses of quinin, or aspirin, or salicylate of soda, and order inhalations of the normal salt solution or of some aromatic oils every two or three hours. Sometimes I direct a few drops of adrenalin (in the proportion of 1:6000, 1:8000, or even 1:12000) to be sprayed or dropped in through the nose, of course, warm or lukewarm. This can be done every few hours according to the age and condition of the child. A Priessnitz compress around the neck over night is sometimes of great service.

I will omit here the acute follicular and the acute phlegmonous inflammation of the pharyngeal tonsil, and come now to the well-known chronic form.

HYPERTROPHY OF THE PHARYNGEAL TONSIL.

As a matter of fact, this affection ought to have been recognized one hundred or more years ago, if some one only had had the idea, that apparently is not too far off of applying digital examination of the retropharynx in cases of obstinate nasal obstruction. By digital examination alone has it been made possible without any special skill to generally recognize the hypertrophy of the pharyngeal tonsil so very frequent among children. Surgical operations with cutting instruments in place of the inconvenient and difficult galvano-cautery operation has opened the therapy of this affection to the whole medical world (R. Kayser). It is important that this has been finally accomplished, since no single affection in the whole domain of medicine and surgery has so far-reaching an influence on the public health as these "innocent little overgrowths," as Dr. Francis² called them. I should rather call them "nocent," knowing what harm they can do. When we consider that Max

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Diseases of Children, and approved for publication by the Executive Committee: Drs. H. M. McClellan, Edwin Rosenthal and Samuel W. Kelley.

Schaeffer,³ twelve years ago, found among his 1,000 cases of adenoids that in 467, or in almost 50 per cent., the ear was involved, and that William C. Braislín⁴ concluded in 1900 that ear diseases almost invariably accompany adenoids, we have no right to call these growths innocent. They have done an immense amount of harm. "But by sheer weight of evidence, practitioners of medicine, parents and others who have to deal with the young, have at last become convinced of the practical importance of the discovery, with the result that many cases which forty years ago would have been allowed to lapse into hopeless deafness, or to fall victims to all the baleful consequences of neglected middle-ear suppuration, are quickly recognized and relieved."⁵ These last remarks were made by Yearsley only a few months ago. When we analyze statistics, however, we must ask ourselves: Are these little overgrowths on the increase among children or not? It seems to me that it is a difficult matter nowadays to find a city child without adenoids. There is, to my knowledge, no difference as regards climate, race or sex. No race is exempt from them, and while we formerly believed that they were more prevalent in damp or cold countries, we now know that this was a fallacy. In all countries, whether warm or cold, dry or moist, adenoids prevail, and they occur *ceteris paribus* in about the same proportions. The cause of their existence in such enormous numbers is due to our modern mode of living, to the poor and faulty hygiene that surrounds our children. But what special agent is at the bottom of it? Is it the housing of our young ones; is it the simply appalling neglect of all laws of hygiene in our public and private schools, not to mention the "kindergartens"; or is it something else? Many of our kindergartens, indeed, deserve much more the name of pest-holes than that of a "garden." A "kindergarten" in a low, overheated, ill-ventilated room, as most of them are, is a menace to humanity. Fröbel would turn around in his grave if he were to see what sort of hygienic methods we apply to the growing generation. And most of the other schools are no better. These, indeed, are the most favorable conditions for developing catarrhal inflammations of the upper air passages, as I have attempted to prove elsewhere. These inflammatory processes of the nose, and especially of the retropharynx, are the first consequences of the evils just spoken of, and on the basis of these "catarrhs" adenoids develop. They are nothing but the products of a reactionary inflammation, which sets in easily under the influence of various causes. If, for example, such a child is infected with scarlatina, measles, diphtheria or other diseases affecting the upper air passages, the already existing "catarrh" is very much exaggerated, and then, as happens so often, we have adenoids following almost immediately these infectious diseases. Such children are sometimes brought to me for the removal of their adenoids, at a time when I consider it too early to operate, being afraid that the infectious germs have not yet left the air passages. To give you an illustration: A child came to me two weeks after it had recovered from a pronounced attack of faucial diphtheria, and as the Klebs-Loeffler bacillus has been met with at a much later period in such cases, I thought it possible that a new infection might be stirred up by the operation, and therefore postponed it.

Even tuberculosis can make its primary entrance through the retropharynx, as I proved some six years ago. It is evident that the organism frequently reacts against the invasion of tubercle bacilli and hypertrophy

of the lymphoid tissue at the vault of the pharynx is one of these evidences. There is no doubt that tuberculous infection of the adenoid tissue is found only exceptionally, and the reason for this is that the mucous membranes of the head, as has been shown by Bollinger, manifest a marked tendency to allow the tuberculous virus to pass through their structures without themselves becoming diseased. And this quality is considerably enhanced by pathologic and especially inflammatory changes of the mucous membrane. Thus we can understand how the tubercle bacillus will penetrate even the intact mucous membrane. But what becomes of it? No doubt when it penetrates into the deeper tissues its passage is followed by a reaction, the result being adenoids in the first instance, and then, lower down, swellings and suppuration of the glands of the neck. That these lymphomata colli are in direct connection with the adenoids is demonstrated by a case like the following:

CASE 1.—N. McC., twelve years of age, was brought to me from Newark, N. J., with a lymphoma colli. I opened it, removed a quantity of pus and all diseased tissue and sent him to the mountains for four months, where he was under the best care possible. The boy had slight adenoids which I did not remove at once, as the parents would not give their consent. The wound did not close, and finally, after three months more had elapsed, I operated on the adenoids, with the result that within two weeks the wound of the neck had healed thoroughly. The boy is now in his seventeenth year and well and strong.

I am under the impression that since adenoids are removed so freely, glandular swellings of the neck have become more rare. I am not positive whether further statistics will verify this.

If, for some reason or other, the path of the tubercle bacillus downward is closed up, by adenoids or other causes, it makes its way to the cranial cavity, where a tuberculous meningitis, so frequent among children, is set up. This idea of mine is as yet only a theoretical deduction which I can not prove, but which seems very plausible to me. In an interesting paper read before the International Medical Congress at Paris, 1900, Dr. Paul Gallois spoke on infection through adenoids. He also believes that they are not only the seat of primary infection, as tuberculosis, etc., but that they are apt to spread various infections to the neighboring and remote parts of the organism. Thus, for example, many cases of infectious sore throat result through adenoids; rheumatism in different parts of the body also has its primary origin in the retropharynx, the seat of these little growths. This is a point which I mentioned some time ago. We have come to recognize the fact that articular rheumatism often gains access to the system through diseased tonsils. While this may be true to a certain degree, I have always, and still do, attribute the more important rôle in this respect to the retropharynx. We see the tonsils, but we do not see the retropharynx, that is to say, in over 50 per cent. of all cases it is not examined, and therefore overlooked. Adenoids only help to make it the *locus minima resistantia*.

Normally with the establishment of puberty an atrophy of the pharyngeal tonsil sets in, which B. Fränkel called the physiologic involution. This involution also takes place when the adenoid tissue is hypertrophied and has not been removed. One of the results of this is the familiar form of postnasal catarrh; another the cicatricial process known as Tornwaldt's disease. Schwabach has shown that what is called by Tornwaldt the bursa pharyngea, was perhaps nothing else than the median recess of the pharyngeal tonsil. On

the other hand, Lehmann has described a congenital cyst of the bursa pharyngea. There can be no doubt however, that the bursa itself is an anomalous structure, as has been demonstrated by Mégevand, Max Schäffer, Jonathan Wright and others. It must surely be of rare occurrence, since it has been observed by so few authors.

As the time allotted me has nearly expired I shall discuss only a few additional points. To speak before such a society as this is, of the symptoms, diagnosis and effects of adenoids on the ears, eyes, respiratory tract, abdomen (hernia⁶), and on the system at large, would be to carry coals to Newcastle. All this is so well known that I believe it to be the common property of every practitioner.

Let us rather discuss the questions: 1. When shall we operate? 2. How to operate? And 3. What are the after-effects and after-treatment?

WHEN SHALL WE OPERATE?

As these growths make their presence felt from the first weeks of infancy we sometimes have to operate very early. The youngest child that I have operated on was seven months old. In this case there was a stubborn mucopurulent discharge from the nose that did not yield to any treatment, and as the whole postnasal space was filled with adenoids the child took nourishment with great difficulty, did not gain in weight, but was losing ground constantly. All these symptoms improved after removing the adenoids. Still, I do not like to operate on children under 2 years of age, as almost all the reactions I have noted after these operations were in infants under 2 years. I could not always explain their occurrence satisfactorily. In older children I can not recollect having witnessed fever of any account. But I will admit that I have not examined my books with a view of elucidating this point. To go through all my private records and clinical notes for this purpose would mean an immense sacrifice of time, which I did not have at my disposal. Let me narrate some of these instances:

CASE 2.—M. S., 15 months old, was operated on at 8 a. m., May 16, 1887, for simple adenoids. In the afternoon the child was restless, refused all nourishment and felt "very hot." My assistant found an elevation of the temperature of 106 F. at 5 p. m. As I was out of town, I did not see the child until the next morning, when the temperature was perfectly normal. The child recovered quickly.

CASE 3.—Another case was that of H. K., 13 months old (adenoids—otitis media purul.—mastoiditis). A colleague in this city had advised removal of adenoids prior to the mastoid operation. The parents consulted me, and I coincided with his opinion. On Feb. 16, 1894, I removed the adenoids under chloroform. February 18, the child had a temperature of 104.2 F., profuse discharge from the nose and ear, was restless and took no nourishment. There was a good deal of edema of the uvula and the neighboring parts, which I directly ascribed to the operation. The manipulations with the curette were sufficient irritation to produce the edema in this delicate infant. There were no other signs of a reaction. The high temperature persisted for three days, when it became normal again. The discharge from the ear ceased completely two months later, and there was no longer any indication for operating on the mastoid.

I was under the impression that in this case the infection was carried from the ear to the retropharynx. As the result was so excellent I should not hesitate to take the risk in similar cases. On the contrary, I consider any danger arising from the ear, and the inability to take sufficient nourishment as positive indications for removing the adenoids, even in very young infants.

When, however, we have an infant with adenoids that breathes comfortably, sleeps restlessly only when it has a "cold," has no ear complications, and takes his food properly, I invariably prefer to defer the operation until the child is 2 years old or more.

When the child is older, shall we operate whenever we find a hypertrophy at the vault of the pharynx? By no means. Slight hypertrophies are so common that they can hardly be called deviations from the normal, and to operate in such cases is, to say the least, unnecessary. But, unfortunately, it is done, and like every other procedure that has proved of benefit, it is overdone nowadays. "Many a pharyngeal vault is scratched because it contains a perfectly normal amount of lymphoid tissue, and the child is really suffering from the vasomotor results of too much candy on the stomach, or of too much clothing on the back, or of too much coddling in steam-heated houses." (Jon. Wright.)

Fifteen or twenty years ago there was need of encouraging the performance of these operations, as the majority of practitioners were more or less afraid of handling adenoids. Naturally enough many cases were neglected at that period. Nowadays it is better to sound a word of warning in the other direction. I know that too much operating is being done already. I could narrate many interesting instances of this kind coming under my observation, were I not limited in time. Not every child that snores, or keeps his mouth open, has adenoids. There are other things under the sun which we can and must treat!

In spite of this, there remain thousands of cases where we absolutely have to remove the adenoids. When a child is brought to us with adenoids which even only partially close up the choanæ, or where an affection of the ear coexists, not to speak of all the other well-known lesions that occur as sequelæ of adenoids, then I advise the parents that an operation is absolutely necessary and that its omission is a very risky and perhaps dangerous matter for the child's health.

Let me remark, apropos, that I have not had the courage to operate in hemophylia, of which four cases came under my care.

THE OPERATION ITSELF.

When we do operate, we must not forget that we have here to deal with a field that is already infected. No amount of antiseptics will do any good in these cases, but they are liable to do harm. My views on this point have not changed for the last fifteen years, viz., that a strict antiseptics is impossible and unnecessary. And this is much more true of the retropharynx than of the nose.⁷ To wash out the mouth and throat with any antiseptic solution will be found bacteriologically entirely useless. The contact of the fluid with the mucous membranes is too short to kill even one bacterium. To wash the face of the child with an antiseptic or alkaline solution before the operation is playing with antiseptics, and syringing the nose before operation is dangerous.

For general anesthesia I invariably use chloroform, and am glad to be in accord with Sir Felix Semon and W. Williams.⁸ The few times I have given ether or ethyl chlorid I was not pleased with the narcosis. Still, I must admit that there are many with wide experience who prefer either of these two.

In what position shall the patient be placed during the operation? For several years, i. e., from 1887 to about 1893, I used the upright position, recommended again lately by T. R. French⁹ in a very instructive paper. I have given it up again: firstly, because in older chil-

dren it was not very convenient, although I must admit that at that time I did not have Dr. French's chair, but an arrangement of my own; secondly, because I encountered a few times unpleasant accidents which were due to the upright posture. I remember one case, a boy of 14, without any cardiac lesions, where breathing stopped for a while, the pulse became very feeble and I had to work for about two hours in order to thoroughly restore these functions. Similar, though not marked accidents have happened in several other instances, so that I came to the conclusion that the danger of anesthesia in the upright position was well founded. Nor do I favor the so-called Rose's position, i. e., with the head of the patient drawn over the edge of the table and bent well back. I am under the impression that the severe hemorrhages that have followed operations in this posture were indirectly caused by it. The surgeon is more apt to cut deeper into the normal tissue and wound a larger vessel, as the parts are more exposed.

Without any previous preparation, as douching or syringing, etc., my patients are put on a table, lying on their backs, the head not being raised and only a small pillow placed under it. The chloroform is given "slowly and cautiously and not pushed to the abolition of the cough reflex which protects the larynx against the entry of blood (*loc. cit.*, p. 720). The operation is performed thoroughly, which means all pathologic growths are removed and not a particle more. We "shave the surface clean" and no more. The attempt to remove all adenoid tissue will only be made by those who forget that we normally have and need such tissue. I have seen the operation done so "thoroughly" that I was convinced that the greater part of the mucous membrane was removed at the same time. On the other hand, there is often found a mass of thickened tissue at the lower edge of the growth, just below the velum, and I agree perfectly with French that this should always be well removed. A failure to do so is invariably followed by incomplete restoration of nasal breathing.

AFTER-TREATMENT.

As mentioned above, no preliminary antiseptics are used. If the patient bleeds during the operation and the blood flows through the nose and down the pharynx, this is the best antiseptic we could wish for. At the same time the flow of blood brings away masses of pus and purulent matter that have accumulated there for long periods. My after-treatment consists in keeping the child quiet for a day or two and giving it a carefully selected diet. The application of a nasal or postnasal douche I consider dangerous right after the operation.

Now, some children will not lose their nasal or postnasal catarrh after the operation, and the most important after-treatment has to commence then. I prefer to perform these operations in the spring, so that the children can stay, if at all possible, a few months in the country, preferably in the mountains. I have had excellent results in these cases, although naturally I have had to operate during all seasons, and with good results, too. In other instances, we have to treat the nasal catarrh systematically. If this be neglected, and it is neglected often, then a relapse of the adenoids will surely follow. In six months or a year we will have the same amount of adenoid tissue as before, in spite of the fact that they had been removed thoroughly. In very old and neglected cases we sometimes find hypertrophies of the turbinals which do not recede after the operation. In such cases, which are nothing but *stasis hyperemia*, the galvano-cautery was formerly used. I warn

against such a procedure, as atrophy of the nasal mucosa will often follow. The congestion that is present ought to be removed by treating the postnasal catarrh with all perseverance possible. Among the different factors in the management of adenoids none is of greater importance than the treatment of these nasal and postnasal catarrhs after operation, and by persistent efforts in this direction I am sure that we will obtain still better results than have been achieved heretofore.

I trust these brief remarks will bring out a full discussion which will prove of benefit to all of us, although I am fully aware that I have touched on only a few points of this extensive and important subject.¹⁰

BIBLIOGRAPHY.

1. Der Schlundkopf des Menschen, 1868.
2. "The Versatility of Adenoid Vegetations," Australasian Med. Gaz., Jan. 20, 1898.
3. "Bericht über 1,000 adenoid Vegetationen," Wien. Med. Woch., 1890.
4. "Effects of Adenoids on the Ear," Brooklyn Med. Jour., p. 520, 1901.
5. Macleod Yearsley: "Adenoids." London, 1901, p. 11.
6. W. Freudenthal: "Die chronischen Erkrankungen der oberen Luftwege und Unterleibsbrüche." Berlin, 1892.
7. W. Freudenthal: "Asepsis or Antisepsis in Nasal Surgery?" N. Y. Med. Jour., Feb. 11, 1893, p. 154.
8. Allbutt's System of Medicine, vol. iv, p. 720.
9. Lymphoid Growths in the Vault of the Pharynx. Reprinted from Brooklyn Med. Jour., 1893.
10. After I had finished this article I received a book by Sir Felix Semon, entitled "Some Thoughts on the Principles of Local Treatment in Diseases of the Upper Air-Passages." London, 1902. Dr. Semon deals here extensively on adenoids and I appreciate that we agree on most points. His remarks are so excellent and touch in a broad sense on so many questions that I believe every practitioner ought to read this booklet.

CHLOROSIS.*

C. F. WAHRER, M.D.
FORT MADISON, IOWA.

ETIOLOGY.

This affection is, with very few exceptions, confined to the female sex, and that principally during adolescence, usually heralding some disturbance of menstrual function, but chiefly characterized by deficiency of hemoglobin of the blood. It seldom affects early childhood, and if so, is usually connected with precocious development of sex. It is also seldom seen after the thirtieth year, but is by far most common at the beginning of puberty. It is also claimed to be at times hereditary, or apparently so. Sedentary occupation, indoor life, unsuitable food, excessive physical and mental activity, emotional or psychic disturbances are all causative in the majority of cases of chlorosis. It is remarkably frequent in girls who menstruate prematurely, indeed, one can usually predict a case of chlorosis in girls whose first menses appear very early. Often this affection is associated with anemia, coexisting or following soon in its wake. Careless observers may make a mistake in diagnosis, though this is one of the most easily diagnosticated of all diseases. These patients are often plump, even fat, or may be plethoric, making a striking contrast to the various cachexiae for which it is sometimes mistaken. It is never due to menstrual troubles; these may follow, not precede or cause chlorosis. Nor does it ever give rise to tuberculosis, as popular fallacy teaches, but the reverse is exactly true, although at times it follows long protracted sickness and wasting discharges.

The worst cases we find among factory girls in large cities, for obvious reasons; however, it is not rare in all

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Diseases of Children, and approved for publication by the Executive Committee: Drs. Edwin Rosenthal, H. M. McClanahan and Samuel W. Kelley.