LETTERS TO THE EDITOR.

(The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.)

The Size of the Brain in the Insectivore Centetes.

I DO not think that there is any recent mammal which has so small a brain in proportion to the size of the skull as has *Centetes caudatus*. In an individual of this species, the property of the Hon. Walter Rothschild, which was lately living in the Zoological Society's Gardens, I found the total length of the brain to be 28 mm., of which no less than 8 mm. were taken up by the enormous olfactory bulbs. The dried skull of that individual—measured along the base and not taking into account the projecting occipital region—was 96 mm. The greatest diameter of the brain is 16 mm.; the skull in that region is from 28 to 40 mm. broad. The small size of the brain relatively to the skull has been frequently commented upon and figured in certain of the early genera of Ungulate mammals; and it may be noted that the measurements which characterise *Centetes*, undoubtedly an early type of mammal, are by no means unlike those of such a genus as *Coryphodon*, judging, that is to say, by the published figures of the brain cast and skull of that animal. The resemblance is increased by the small size of the cerebral hemispheres, and by the complete exposure of the corpora quadrigemina in *Centetes*. I hope shortly to give a fuller and illustrated account of the brain of this Insectivore.

FRANK E. BEDDARD.

Thermochemical Relations.

IN a letter in NATURE of February 7 (p. 348), Signor Carlo Del Lungo calls attention to certain thermochemical equalities, and asks for an expression of opinion as to whether these are the result of a casual coincidence or of a definite law. The equalities in question, if they were not accidental, would have an extremely important bearing on thermochemistry generally, but unfortunately, I believe, they are probably mere accidents of coincidence.

The two instances given would seem to indicate that when two elements combine together, the heat evolved is equal to the sum of that which is liberated when these same two elements combine separately with some third. Thus we have (modifying Signor Del Lungo's equations so as to indicate more clearly that the heat of combination refers to that of the elements taken in their ordinary molecular condition):—

and

I have little doubt but that such coincidences might be shown to be accidental by the existence of similar sets of compound, the heats of formation of which show no such relationship. Unfortunately, I have with me no thermochemical works to supply the requisite data. An examination of the above instances, however, will, I think, suffice to prove that the coincidences cannot be the result of any definite law.

When we expand the first of the above equations so as to represent all the stages involved, and the combinations and separations of the atoms in the various molecules, we get:—

which simplifies into

This would mean that any pair of the four groups of atoms in the following diagram would combine together in any way, and yet liberate the same amount of heat—i.e. the thermal result would be the same whether the products were oxygen gas and hydrogen chloride, or water and hypochlorous anhydride.

NO. 1634, VOL. 63]

This, if established as a generality, would be of fundamental importance. But it is inherently improbable, and the existence of the second instance given by Signor Del Lungo seems to render it untenable. If it were true as regards the gases figuring in the above reaction, it would certainly not be true if one of the products concerned were in the solid condition; if, for instance, we dealt with solid instead of gaseous oxygen, there would be a divergence from equality equal to the heat of condensation and solidification of oxygen. Yet the second instance given by Signor del Lungo contains terms of this very character. When treated as above, assuming for the sake of simplicity that solid carbon molecules contain only two atoms, it gives

Thus the two instances given, inasmuch as they are necessarily dissimilar, must by their coincidence show that this coincidence is the result of accident, and not of any thermochemical law.

Woolacombe, February 12.

SPENCER PICKERING.

The Fraunhofer Lines in the Spectrum of the Corona.

IN some of the preliminary reports of the observations made during the last eclipse, undue importance appears to have been attached to the supposed absence of the Fraunhofer lines from the spectrum of the corona. Mr. Newall, for instance (*Roy. Soc. Proc.*, vol. lxvii. p. 365), says, "it is difficult to reconcile the marked polarisation (of the coronal light) evidenced in this investigation with the absence of Fraunhofer lines in the spectrum of the corona." Mr. Abbot, whose bolometric observations appear to him to indicate that the corona does not reflect much solar light, states that "additional evidence against the theory of reflecting particles is found in the Indian eclipse spectroscopic results of Campbell, who found a continuous spectrum from the inner corona with total absence of dark lines" (*Astrophys. Journ.*, vol. xii. p. 75).

Prof. R. W. Wood (*NATURE*, vol. lxiii. p. 230) has already suggested a partial reconciliation of the apparently conflicting observations, but there are other points which do not seem to have received sufficient attention.

It must be remembered that the Fraunhofer lines were observed in the corona of 1871 by Janssen, and were subsequently photographed by Schuster in 1882, and by Hills in 1893, besides having been seen by various observers on other occasions. Great weight seems to have been attached to the photographs taken in 1898 and 1900, in which no dark lines were recorded, but I venture to suggest that the apparent absence of the lines on these occasions was mainly due to the use of spectroscopes of too great a dispersion to exhibit them. While great dispersion is well adapted to render the bright line spectrum more effective, it is clearly fatal to the dark line spectrum, as in the observation of solar prominences. As a matter of fact, the recent photographs only show continuous spectrum extending to a few minutes from the sun's limb, and if one may take the evidence of the prismatic cameras, this arises mainly from the bright line giving inner corona, which probably does not shine chiefly by reflected light, and would, therefore, not necessarily show the Fraunhofer lines. This view accords well with Schuster's account of the photograph of 1882, in which it is stated that near G there was a sharp decrease in intensity at 0.29 of the solar radius from the limb, a further falling off in intensity at 0.60, with the spectrum fading out of view at 1.47; in the lower regions the spectrum was perfectly continuous, but in the upper regions the solar line G was reversed. Captain Hills also found, in 1893, the Fraunhofer lines only at a considerable distance from the limb. Stone, too, in 1874, observed the dark lines in the outer, but not in the inner corona. In these early observations and photographs we may take it that the dispersion of the spectroscopes employed was much less than in 1898 and 1900, and it would therefore seem desirable to search for the Fraunhofer lines in the next eclipse with instruments of smaller dispersion than those recently in use for the bright line spectrum and special observations.

It may be further remarked that as we do not see a section of the sun, the outer corona must be superposed upon the inner, and the fact that the Fraunhofer lines have not been observed or photographed in the inner corona, even with small dispersion, is sufficiently explained by the superposition of the dark line spectrum on the much stronger continuous spectrum of the inner corona. Such superposition would, of course, be more effective

in obliterating dark lines than bright ones. Prof. Wood's suggestion as to reducing the effective intensity of the truly continuous spectrum by passing the light through a suitably placed Nicol's prism, would accordingly be specially applicable to the inner corona.

Prof. Wood has obtained experimental evidence on a point to which attention was drawn many years ago by Ranyard (*Mem. R.A.S.*, vol. xli. p. 353), namely, that if the solar light be reflected by small particles in the corona, the reflected rays will be deficient in the rays of greater wave-length. In this way the bolometric observations indicating that the corona was cooler than the bolometer are partly explained. Is it possible that the observations are to be completely explained by further supposing that the bolometer strip was outside the image of the shallow inner corona, which is probably the chief part of the corona directly emitting light and heat? The image thrown on the strip appears to have been little more than one-third of an inch in diameter, and in the account of the observations Mr. Abbot simply states that the image was brought almost tangent to the strip, so that the strip may very well have been 4 or 5 minutes from the sun's limb.

At all events, there seems to be no sufficient ground, as yet, for rejecting the view that the luminosity of the outer corona is largely due to reflected solar light, while that of the inner corona is partly due to the incandescence of solid or liquid particles and partly to gaseous radiations. A. FOWLER.

Royal College of Science, South Kensington, February 9.

Malaria and Mosquitoes.

I THINK most of those who have had much experience of the Indian jungles would be prepared to corroborate the remarks made by Mr. D. E. Hutchins in last week's *NATURE* (p. 371), and would perhaps be inclined to think there may be something in the opinion alluded to by him, that "Dr. Ross's splendid discovery does not quite cover the whole ground." In 1873-4 I spent some months in the notoriously malarious region at the foot of the Darjiling Himalayas, which contained some tea gardens here and there at that period, while many new ones were being opened out. The planters suffered greatly from malarial fever, and I was told by several that it was far more prevalent, and of a worse type, on gardens in process of formation, by clearing the jungle and breaking up the ground, than on either the undisturbed jungle itself or on gardens that had existed for some time. In other parts of India my camp has suffered badly from malarial fever when mosquitoes were certainly not prevalent, and when, to the best of my belief, there were none, or perhaps I should rather say, none made themselves noticeable by stinging.

On the other hand, when in Upper Assam in 1874-5, I was informed by some of the planters there, and it seemed to be a commonly received opinion, that mosquito curtains were a valuable protection from malaria. Of course at that period no explanation could be given for the supposed fact, which seemed a very mysterious one, as for the invisible 'germs,' which were thought by some to float in the air, to be excluded by curtains of ordinary mesh would be something like a man being prevented from crossing a road through inability to squeeze himself between the milestones.

18 The Common, Ealing, February 12. F. R. MALLETT.

Audibility of the Sound of Firing on February 1.

THE following note was recorded here (at Littlemore, Oxford) immediately after I was called away from listening; and as it is more detailed than any I have seen, I daresay you will think it worth printing.

I held my watch in my hand and observed the sequence of the booms for some ten or twelve successive minutes. The second one showed there was regularity, and after the third the facts were clear. The sounds reached us thus:

0s. to 20s., continuous unbroken roll of guns of slightly different strength.

20s. to ± 24 s., silence for 4s. or 5s.

± 24 s. to ± 29 s., similar roll.

± 29 s. to ± 34 s., silence for 3s. or 4s.

± 34 s. to ± 38 s., a similar roll.

± 38 s. to 42s., silence to just the 42s.

42s. to 45s. exactly, a short roll culminating in three or four guns much louder than any of the preceding ones.

45s. to 60s. exactly, dead silence for a quarter minute.

NO. 1634, VOL. 63]

This recurred with such perfect regularity that there was no doubt whatever about the precision of the observation.

Then I had to ride away on a bicycle about a mile, where I met another cyclist, and we stopped and listened again. The long 20s. roll was now missing, but punctual to the second or two came all the others, the last one with big guns precise.

It seems likely that the long roll was the *simultaneous* salute of the long line of ships; but I have not yet noticed any authoritative statement as to how the saluting was done. It was interesting to think that at any moment while we were listening there were no fewer than six other such salutes travelling up towards us from the Solent.

From the strength of the sounds I should quite agree with the opinion expressed in Oxford that they might well be heard another twenty miles.

W. J. HERSHEY.

February 12.

The Origin of the "Tumbling" of Pigeons.

IN reading the account of tumbler pigeons in Darwin's "Animals and Plants under Domestication," the question arose to what the habit of "tumbling" might be due.

I have since seen, in a report of an address by Prof. J. A. Thomson to the North British Branch of the Pharmaceutical Society, a statement that the peculiarities of movement of the Japanese waltzing mouse are due to the fact that only one of its semi-circular canals is developed.

It occurred to me that a similar condition might explain the movements of the tumbler. I should be much obliged if you or any of your readers could tell me where information concerning either of these examples is to be found. E. P.

February 3.

[Prof. Thomson's statement is not quite accurate. The paper to which he referred is undoubtedly that by Rawitz, in *Archiv. f. Anat. und Physiol.* (Physiol. Abth.), 1899, pp. 236-244, where it is shown that of the three canals the anterior is alone normal, and that the other two—though reduced and abnormal in their inter-relationship—are present.

The physiology of the organ has been since experimentally investigated by Alexander and Kreidl, whose paper is in *Pflüger's Archiv. f. Ges. Physiologie* (Bd. 82, pp. 541-552); and as to the pigeons, they, too, have all three canals well developed. Concerning their membranous labyrinth, some experiments of interest were made in, I think, the early nineties by Spamer and others, and an account of these and of allied investigations bearing on the question raised (by McBride) will be found in the *Journ. Anat. and Physiol.* (vol. xvii. pp. 211-217). There does not appear to be any foundation for the view to which the writer of the letter inclines.—G. B. HOWES.]

Lummer's "Photographic Optics."

THE interesting review of Prof. Silvanus Thompson's translation of Dr. Otto Lummer's "Photographic Optics," which was published in *NATURE* on January 3, has come under my notice. I should be obliged if you would permit me to correct a mistake which occurs both in Prof. Thompson's book and the review. Concerning the two views of Munich published in Prof. Thompson's work, which are there attributed to Dr. Miethe, you remark: "Miethe's two views of Munich from a distance of about two miles—the one taken with an ordinary lens, the other with the teleobjective—show what a powerful weapon the latter is."

Both the views in question were taken by my firm, one with an antiprism and the other with this antiprism in combination with a magnifying system (negative lens) as a teleobjective of our own construction and manufacture. I send you a prospectus of my firm concerning the said teleobjective and containing also the two views of Munich.

I also observe another error in Prof. Thompson's work; the lenses recently introduced by Messrs. R. and J. Beck being described on p. 80 as Messrs. Voigtländer's Collinears, instead of our orthostigmats, for which I have given them a manufacturing license, as you will see in the notice on page 1 of the prospectus I also send by book post.

RUDOLF STEINHEIL.

München, February 5.

NOTHING could have been further from my intention than to do any injustice to Dr. Steinheil or his firm, and I hasten to acknowledge the justice of his criticisms upon the slips