ON NOCTILUCINE, THE PHOSPHORESCENT PRINCIPLE OF LUMINOUS ANIMALS.*

By DR. T. L. PHIPSON, F.C.S.

Member of the Chemical Society of Paris, and of the Royal Society of Medical and Natural Sciences of Brussels, etc.

In reviewing the various phenomena of phosphorescence, which are so numerous, both in the mineral world and in organic nature, it is not very difficult to convince one's self that in the great majority of well-observed cases belonging to the former, the production of light can be directly referred to electrical action, or to electrical action resulting from chemical action, and the latter mostly from oxidation.

As examples of purely electric action, may be cited the light produced in the formation and cleavage of crystals; the sparkling of boracic acid when cooling after fusion; the light emitted by benzoic acid during its sublimation; the flash of light that accompanies the cleavage of a plate of mica, when one fragment is found to be electro-positive, and the other electro-negative, immediately after the operation; the light emitted when quartz pebbles are violently rubbed together, which is accompanied by an odor of ozone; the flickering scintillations observed when a mixture of sulphate of soda and sulphate of potash crystallizes, when arsenious acid crystallizes, when crystals of sugar or nitrate of uranium are broken, and a very great number of similar cases which it would be far too long to mention here. In fact, the more the phenomenon is observed, the more general it appears to be throughout nature.

As examples of electrical action following on chemical action, may be cited the luminosity of phosphorus, of potassium and sodium, of arsenic, sulphide of antimony, and a considerable number of cases where oxidation, or other chemical action is intense. I have referred to all these in another place.†

In the organic world, with one exception, the luminosity of animals and certain plants can be with tolerable certainty referred to the production of the principle which I have termed noctilucine. The

*Read before the British Association for the Advancement of Science, Bristol, 1875.

† "Phosphorescence; or the Emission of Light by Minerals, Plants and Animals." London, 1862.

one exception is when the production of light can be traced also, with the greatest probability to electrical action. This is evidently the case with the sparkling radiations which are occasionally seen round the flowers of the marygold and several other plants at a certain period of their growth, when the atmospheric conditions are favorable. The phenomenon is coincident with the rupture of the anthers and the ejection of the grains of pollen. We have other examples of a similar production of light resulting from the sudden rupture of plant tissue in dry weather.

Noctilucine is a peculiar organic principle, which appears to be very widely spread through nature. It is the cause of the light of the glow-worm, the scolopendra, the fireflies, the pholas, and numerous other luminous animals, both during life and for a certain time after death. The first allusion to the existence of this substance occurs in my note, "*Sur la Matiere Phosphorescente de la Raie*," published in the *Comptes Rendus* of the Paris Academy in 1860, where I have spoken of it as a peculiar organic matter, devoid of phosphorus, but shining in the dark like that element itself. It was found, by the most careful tests, to contain neither phosphorus nor phosphoric acid. It is again alluded to in my work on "*Phosphorescence*," published in London in 1862, (p. 103), and finally in my paper, "*Sur la Noctilucine*," in the *Comptes Rendus* of the Paris Academy for August 26, 1872, (p. 547). The same substance is noticed again, exactly one year later, (August 25, 1873), in the same publication by Professors Ch. Robin and M. Laboulene.

Noctilucine is not only the cause of the phosphorescence of dead fish and animal flesh of every kind, but it is secreted by living animals, such as the glow-worm, the scolopendra, the fireflies (*Elater*), the common earth worm, and by a very considerable number of others, amounting probably to many hundred distinct species, that are seen to shine in the dark. It appears to be produced also by certain living plants—*Agaricus*, *Rhizomorpha*, and *Euphorbia* (?)—and occasionally in the decomposition of vegetable matter (putrid fermentation of potatoes, etc).

At the ordinary temperature of summer in this climate, noctilucine is a fluid nitrogenous substance, slightly viscous or oily in appearance that can be mixed with water, but does not dissolve, though the watery liquid can be filtered; its specific gravity is slightly less than that of water; its color is white, when impure, yellowish white, and when decomposed becomes brown.

from the glow-worm or the scolopendra, possesses the same chemical properties. It is secreted in a state of great purity by *Scolopendra electrica*; and in the month of September it is possible, by causing several of these myriapoda to run about on a flat glass dish with vertical slides, to collect enough of it to examine its principal characters. From the luminous organ of *Lampyris noctiluca*, and from the surface of phosphorescent fish, (stockfish, mackerel, herring, etc.), it can also be obtained, in a less pure form, by collecting on a damp filter the luminous matter, washed or scraped from the surface, or, better, by stirring the shining matter with water and allowing the liquid to stand in a narrow vessel till the next day, when the noctilucine will form a layer upon the surface that can be separated; but it is already partially decomposed.

The experiments I have made with this substance for several years past, whenever I have had an opportunity of observing it, lead me to the conclusion that it is of a basic or neutral, rather than an acid, nature; and, though I have not yet obtained it either pure enough or in sufficient quantity to investigate it very minutely, I believe that noctilucine belongs to the propionic series. It may, perhaps, hereafter be proved to be a cyanic derivative of propylic aldehyd, belonging to the same class of bodies as *leucine* (cyanhydric derivative of amylic aldehyd) or *creatine*. This would also tend to account for the final resolution of noctilucine into *urates* of soda and ammonia, which are almost always found in the luminous organs of various animals.

The secretion of noctilucine by the higher classes of luminous animals, such as the insects (*Elater*, *Lampyris*, etc.) and myriapoda, is, doubtless, to a certain extent, under the influence of the nervous system, which gives them the faculty of causing the light to cease. In this case the secretion is arrested for the time. (It has been long known that the eggs of the glow-worm shine for some time after they are laid, so that they also must contain a small quantity of noctilucine.)

In animals much lower down in the scale, such as the little *Noctiluca miliaris* of the English Channel, the polypes, medusæ, etc., it appears to me quite evident that there exists also a *special organ* for the production of noctilucine; and even here, where we find scarcely any indication whatever of a nervous system, the secretion of this luminous matter often appears to be subject to external influences.

London, October 28, 1875.