bark, and belonging to the same order, and a native of India, of the Malay Archipelago, and of tropical Australia.

For over 18 months I have employed a tincture of the bark as a general tonic and more particularly in influenza. I start with it as soon as the feverish stage is over and give from five to 10 minims in chloroform water four times a day, preferably some little time before meals. Acting as it appears to do on the skin and kidneys it can be administered with safety even before convalescence has set in, seeing that it helps in the elimination of the poison. I can confidently recommend its use as a valuable agent in the treatment of influenza. Like every other drug it does not act in every case, but generally speaking one finds it useful. The Addendum dose of the tincture is from a half drachm to one fluid drachm. This is too much for a start, and I prefer to start with 10 minims or even in many cases with five minims, and, as I have said, chloroform water is the best vehicle.

Leeds.

DIPHTHERIA ANTITOXIN USED SUCCESSFULLY BY THE MOUTH.

BY W. CAMPBELL M'DONNELL, M.R.C.S. ENG.,
L.R.C.P. LOND.

IT being thought that the antitoxin of diphtheria must for successful exhibition be hypodermically injected the following may be useful.

A girl, aged 14 years, having both tonsils covered with a soft white pellicle and with no symptoms but a sore-throat, very slight enlargement of the related lymph glands, and a temperature of 38.2°C ., at 11 A.M. on Nov. 12th, 1900, was given 1500 units of antitoxin by the mouth on the second day of the illness. The antitoxin had been issued by Burroughs and Wellcome in the dry state and was 20 months old. One patch yielded the diphtheria bacillus, as was certified by the medical officer of health. Within 20 hours of giving the antitoxin the right tonsil was quite clear. No application was made nor was any gargle used to the throat. On the third day there was no pellicle to be seen and at 11.30 A.M. the temperature was 35.8°C . On the fourth morning the temperature was 37°C . On the tonsil which was at first the most affected was seen a slight recurrence of pellicle. For this another 1500 units of the dissolved dry antitoxin were given by the mouth. She recovered uneventfully and as quickly as possible.

In an earlier case, where the struggling on attempting to hypodermically inject the antitoxin of diphtheria made that method impracticable, the antitoxin was rectally injected. The benefit was equal to that in the case described above. In this case, however, diphtheria was not bacteriologically proved, but clinically it was certain.

Stoke Newington, N.

A TUMOUR SIMULATING EPITHELIOMA WHICH APPEARED TO FOLLOW THE STING OF AN INSECT.

BY EDWARD J. WOOLLEY, M.B., B.C. CANTAB.

THE following case is of interest chiefly on account of the appearance of an uncommon causation, the tumour itself presenting no characteristics to differentiate it from epitheliomata from commoner causes and in more usual positions.

The patient was a labourer, 54 years of age. He was thin almost to emaciation, but said that he had always been so and that he had noticed no loss of flesh in the last few months. In July, 1900, he was stung, or bitten, on the external surface of the forearm by an insect which he described as "a green fly." A swelling of the size of a common nut appeared at the site of the sting and the patient rubbed it and attached no importance to it, thinking that it would go down. This, however, was not the case. Six weeks after the injury there was considerable itching in and about the swelling. The patient poulticed it and bathed it frequently with hot water. He noticed at this time that the swelling was getting larger. He continued his poultices and bathings until Dec. 14th, when he came to me. The tumour was then circular in shape and of about one inch in diameter, being raised above the surface of the surrounding healthy skin not

more than one-sixteenth of an inch. The centre of the growth was hollowed out slightly and had the appearance of a dry ulcer. There was some hardness, but not the hardness to be expected in a malignant growth; and there was no pain. The glands in the axilla and the supra-condylar gland were not enlarged. The skin in which the growth was situated was freely moveable on the subcutaneous tissues. Although the history and the appearance of the growth were not strictly those of epithelioma, they seemed sufficiently so to call for its excision and on Dec. 20th I removed it, under ether, by an elliptical incision. I found the growth to be confined exclusively to the skin, and the tissues under it to be quite healthy and not at all adherent to the tumour. A microscopical examination showed the tumour to be a squamous-celled epithelioma with well-developed cell-nests—a typical epithelioma of the skin. I should add that no members of the patient's family on the male or female side have suffered from cancer.

Kingston-on-Thames.

A Mirror OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv., Proœmium.

LONDON HOSPITAL.

A CASE OF HYDATID CYST OF THE LIVER RUPTURING INTO THE TRANSVERSE COLON.

(Under the care of Dr. F. J. SMITH.)

EVACUATION of the contents of a hydatid cyst of the liver into the bowel is not a common occurrence; more frequently the cyst makes an opening into the general peritoneal cavity, or through the diaphragm into the pleural cavity, or into the bronchial tubes of an adherent lung. When, however, the intestinal route is chosen by the hydatid, suppuration generally occurs in the cyst, and recovery is often complete and permanent. For the notes of the case we are indebted to Dr. H. N. Clarke.

A man, aged 32 years, was on April 2nd, 1900, admitted into the London Hospital with jaundice. He stated that he had been quite well until November, 1899, when he had an attack of "biliousness" and "constipation." Ten days later he had severe pain across the lower part of the chest; the pain was sufficient "to double him up" but did not make him vomit; the next day he became jaundiced. These signs all cleared up in about a week. Since then he had had five attacks, all the same in character—viz., pain first and then jaundice. He described his stools as being "like white putty" during the attacks. His family history was good; there was no history of syphilis or gonorrhœa. He had always been a temperate man.

The patient was admitted to hospital during his sixth attack of jaundice; he was deeply jaundiced and was much wasted, but he did not know how much weight he had lost. On examination the heart was found to be normal; there was a good deal of general bronchitis, and there was some fulness of the upper part of the abdomen. The liver was enlarged and the edge could be felt about two and a half inches below the costal margin; the organ was tender and a rounded smooth lump three inches across could be felt in the epigastrium, rather more to the left than to the right of the middle line; there was no œdema of the abdominal wall. On palpation the lump gave the sensation of deep fluctuation. The stools were white and clay-like. The urine contained a trace of albumin; the specific gravity was 1040 and bile was present.

Three days after admission—viz., April 5th—the patient had a rigor; the next day he was seen by Mr. Mansell Moullin who advised an exploratory laparotomy. It was decided that this should be done as soon as the bronchitis had cleared up sufficiently to allow the administration of a general anæsthetic. On the morning of April 7th the lump in the epigastrium was observed to be much smaller, and

later in the day some large pieces of white translucent membrane were passed in one of the stools, which had been carefully sifted since the patient's admission. No hooklets or scolices were found. The temperature ranged from 102° to 98° F. From this point the patient made an uninterrupted recovery; the swelling in the epigastrium entirely disappeared, though the edge of the liver could still be felt when the patient left the hospital. The jaundice gradually faded away, the appetite improved, and the patient rapidly put on weight again, being discharged as cured on May 8th.

Remarks by Dr. CLARKE.—Though no hooklets or scolices were found, it nevertheless seems most probable that this was a case of hydatid of the liver which suppurated. The transverse colon probably then became fixed to the under surface of the liver by adhesions and the pus was thus able to make its way out of the abscess cavity into the gut.

Remarks by Dr. SMITH.—This case is an excellent illustration of the occasional expulsion of a hydatid by nature's own means. We had fully made up our minds that surgery afforded the patient the best chance and were only waiting a favourable opportunity for operation when the rupture took place. There could be no room for doubt as to the nature of the membrane passed per rectum, as it was totally unlike any other possible membrane, though it is somewhat unfortunate that no hooklets were found. I can personally only call to mind one other case in which a hydatid has been naturally evacuated with such a happy result; that was in a boy who without any symptoms worth mentioning or physical signs either coughed up from the lung the whole of a hydatid cyst in large fragments. I showed the patient and the membrane at a meeting of the local branch of the British Medical Association in 1899. I cannot add anything of value to Dr. Clarke's account of the present case.

QUEEN'S HOSPITAL, BIRMINGHAM.

A CASE OF RUPTURE OF THE THORACIC AORTA.

(Under the care of Dr. O. J. KAUFFMANN.)

THE very interesting case recorded below is remarkable in that rupture of the aorta apparently occurred without any local disease and without any special strain. It must be remembered, however, that rupture of the aorta is often a slow process. A small fissure may appear in the internal layer of the wall as the result of some sudden increase of pressure, but the tearing of the other coats may proceed gradually and no violent symptoms may be produced until the outer coat gives way. As to the wall of the aorta in this case a microscopic examination would be needed to render it certain that no morbid change was present. Rupture of the aorta may occur in young adults, as in a case recorded by Dr. J. Lumsden,¹ where a young woman, 23 years of age, died from rupture of the aorta after lifting down a box from a shelf, and apparently, from the description, there was only a slight change in the endothelium and in the subendothelial tissue. Occasionally the aorta has been perforated by an exostosis from a rib, as in a case recorded by Mr. E. A. Lightburne,² where a boy, 12 years old, died from this cause with extensive hæmorrhage into the left pleural cavity. For the notes of the case we are indebted to Dr. A. Stanley Barnes, late house physician.

A groom, aged 19 years, was admitted into the Queen's Hospital, Birmingham, under the care of Dr. Kauffmann, on Oct. 11th, 1900. No history could then be obtained, as he was almost unconscious and quite unable to answer questions. The history obtained next day was as follows. He always was a "delicate" lad, but had never suffered from any illness sufficiently severe to keep him in bed. For a month before admission he had complained of an aching and sometimes shooting pain in the lower part of the chest behind (about the level of the eighth dorsal vertebra); this pain was worse on exertion, but was not accompanied by cough nor was it increased on deep inspiration. He consulted a medical man about it on Sept. 20th and was given a bottle of medicine, no physical examination being made. The pain was better on the next day and did not trouble him again for a fortnight, when on its recurrence the treatment was repeated, but this time without benefit. Despite his vague pain he kept at his work until Oct. 11th. On this day, at about 2.30 P.M.,

soon after his dinner, he felt very bad, the pain having suddenly become much worse. He ceased work at once and started for home, catching an omnibus a few yards away; he did not have to run for the omnibus nor had he undergone any other strain that day. Five minutes later the guard noticed that he looked very bad and called a medical man who found him in a collapsed and pulseless condition. He administered three minims of liquor strychniæ hypodermically and sent him at once to the hospital in charge of the police. Neither from the patient nor his parents could any history of an injury to the chest, either remote or recent, be obtained. He had had no cough and had never complained of indigestion. There was nothing in the family history to suggest either syphilis or tubercle; the lad himself denied exposure to syphilitic infection, and the father (with whom he lived) strongly supported his statement.

On admission the patient was a lightly-built lad with very little subcutaneous fat and looking somewhat wasted. He was very pale, the lips having a faint bluish tinge and the conjunctivæ being anæmic. He was partly conscious but could not answer questions and tossed about moaning restlessly on the couch. The pulse was rapid and very small. The abdomen was rigid but not distended nor was there any dullness in the flanks; there was considerable tenderness in the left hypochondriac region; no viscera could be palpated. The chest was slightly deformed, the sternum being too prominent. It was at once noticed that the left side of the chest hardly moved with respiration and that the palpable apex-beat of the heart was in the third interspace in the right vertical nipple line. The left axilla was dull up to the level of the fifth rib, and above this a low-pitched tympanitic percussion note was obtained, reaching as far to the right as the middle line and upwards almost to the clavicle. Behind, the lower part of the left chest was dull as high as the seventh rib, and above this the dullness shaded off gradually to a normal note at the apex. Over the very dull area no breath sounds could be heard and vocal fremitus was absent, in the hyper-resonant area faint breathing with prolonged expiration was heard, and in the partly dull area diminished rather harsh vesicular breathing was heard. Over the right lung no alteration of percussion note was observed and breath sounds though harsh were vesicular. The "anvil-sound" was well heard over the third and fourth left spaces in front. There were no heart murmurs. The pulse was 120, the respirations were 28, and the temperature was 98.2° F.

Morphia was given subcutaneously and the patient was put to bed. He slowly improved and by the next morning was able to give his part of the history recorded above, the rest of it being supplied by his mother and father. On Oct. 12th (the day after admission) he was still very weak and pale, but quite conscious and less restless. The abdomen was rigid and slightly retracted with inspiration, the recti abdominis hardening with expiration; a tympanitic note was obtained all over the abdomen. There was a little tenderness in the left hypochondrium but none elsewhere. The physical signs in the chest remained the same except that the hyper-resonant area extended to the right edge of the sternum and was now of a higher pitch than before.

As the patient was evidently suffering greatly from the increased pressure in the left side of the chest it was decided to attempt to remove some of the air and blood. A trocar and cannula were thrust through the fifth space in the left anterior axillary line—i.e., at about the upper limit of the dullness. The trocar was withdrawn and a water manometer about 10 inches high was attached to the cannula. On releasing the clamp between the cannula and the manometer the water was at once forced up to the top of the manometer tube and then overflowed. A mercury manometer was therefore substituted and indicated at once a positive pressure of 35 millimetres of mercury with expiration. About two ounces of air and a quantity estimated at from 10 to 12 drachms of dark fluid blood were then withdrawn and the intra-thoracic pressure fell considerably; as, however, the tubes became repeatedly blocked by clots of blood the cannula was removed and the wound was closed. For the same reason the final reading of the manometer, just before withdrawing the cannula, was untrustworthy. An examination of the chest immediately after the tapping showed that the hyper-resonant note was now much less high pitched and did not quite reach to the right edge of the sternum.

On Oct. 14th there was slight fever (the temperature being 99.6°) but the general condition was much improved. For the first time a heart murmur appeared, a soft whiff at the

¹ Brit. Med. Jour., Jan. 7th, 1899, p. 21.

² THE LANCET, Jan. 11th, 1896, p. 99.