have had an anæsthetic administered, but I find this is seldom necessary, as the pain is slight. The operation can now be proceeded with. The negative needle electrode is plunged into the root of the hair for about one-sixteenth of an inch, and the positive electrode sponge is applied in the immediate neighbourhood. The needle should be kept in for about the space of five seconds, then the sponge electrode should be removed and afterwards the needle electrode. To know that the operation is effectual the needle should produce slight frothing of the tissues. The hair destroyed can now be easily epilated with an ordinary pair of dressing forceps, and it should come out without the slightest adhesion. This operation applies more especially to hairs that are noticeable to the naked eye. Fine downy hairs can always be destroyed by the application of a properly made depilatory. A slight inflammation of temporary character occurs for an hour or two after the operation in the destroyed follicle; this can be controlled by the use of a soothing lotion. The operation, if carefully done with a battery in good working order, is invariably successful, especially if the hairs are few and of good size—from a quarter to half an inch long. If many hairs have to be removed, then several sittings will be required at intervals of about ten days. A hundred hairs can be removed at a sitting. The following cases may serve to illustrate the operation.

1. A young woman, aged thirty-two, consulted me on March 15th, 1884, with superfluous hair on the upper lip and chin; she had pulled them out herself, and there remained about a hundred short, dark, stumpy hairs. I removed in two sittings as many as I could see; for the removal of the remainder I prescribed a depilatory for her to use. I recently saw this patient, showing the result of the operation to be successful.

2. In August, 1884, a young lady, aged thirty, was sent to me from the north of England, who had had a superfluous growth of hair on the upper lip and chin for about five years. I removed all I could see in two sittings. I heard recently that the operation was successful.

3. I removed a patch of superfluous hair from the neck of a young lady last August, and I received a letter from her mother last week to say the operation was successful.

4. In October a young woman came to me to have a large quantity of hairs removed from the chin and upper lip. This was the worst case I have had, both from the numbers of the hairs, their depth of growth, and the extreme sensitiveness of the patient to electricity. I used the electrode with difficulty after painting the skin with a 5 per cent. solution of hydrochlorate of cocaine, and could only remove about fifty at a sitting. After seven sittings I have nearly accomplished their removal.

5. A gentleman consulted me in the summer of 1885 for a few superfluous hairs on the tip of the nose. I removed these at one sitting, the result being successful.

6. In July, 1885, I removed about fifty hairs an inch in length round the nipples of the breast of a young woman, also some from her upper lip. One operation proved successful.

Since I wrote this paper I had the honour to show my battery and electrodes in the museum, and demonstrate the process in the Surgical Section to the President and members, at the meeting of the British Medical Association in August last, in connexion with the destruction of the lupus nodule, moles, warts, and nævi, in a paper on the "Surgical Treatment of Lupus and Acne" I read at Brighton.

Sackville-street, W.

NECROPSY OF A CASE OF MYXEDEMA.

By GILBERT BARLING, M.B., F.R.C.S.

By the courtesy of Dr. Suckling I had the opportunity of examining the body of a patient of his suffering from myxœdema. A clinical note of the case has been reported by Dr. Suckling in THE LANCET of May 15th, 1886.

The necropsy was made thirty-six hours after death. External appearances: The body of a very stout old woman; the lower lip much everted and swollen; cheeks of a bright-red colour; a well-marked arcus senilis. The legs, face, and arms pitted on pressure, but no fluid could be made out in the abdomen. Incision into the tissues did not reveal the mucoid appearance mentioned in other cases. Head: The cranial bones were very thick, especially in front, there reaching to nearly a third of an inch. The brain and its

membranes presented no abnormal appearance. Weight of brain 2lb. 10 oz. The pituitary body was reserved for microscopic examination, but no change was subsequently found in it. The cervical sympathetic showed no change to the naked eye, but it was not examined with the microscope. The thyroid body was small, especially the right lobe, and it was very tough to the feel, and so intimately blended with the cervical fascia that it was difficult to separate the two exactly. The weight of the organ was 155 grains. Microscopic examination showed a simple atrophy of the gland structure—in some places complete, in others only slightly advanced. If a fairly plump lobule were examined, it showed at its periphery the remnants of atrophying capillaries; next to this a zone in which the capillaries had not undergone marked change, but the meshwork formed by them, enclosing the alveoli, had lost its epithelial lining, as well as the colloid material found inside. In the centre of the lobule, capillaries, epithelium, and colloid material appeared in the relative proportions found in health. There was nothing to indicate that the disappearance of gland tissue was due to a formation of granulation tissue resulting in a cirrhosis. The submaxillary glands weighed one ounce. No abnormal condition was found in them either by the unaided eye or with the microscope. Thorax: The lungs were crepitant throughout, and presented a few caseous and calcified nodules scattered sparsely through them. The heart weighed fourteen ounces, and there was marked hypertrophy of the left ventricle. The valves were fairly healthy, but there was considerable atheroma of the aorta. Abdomen: On opening this cavity several ounces of glairy but clear fluid were found. Some of this collected and boiled became practically solid in the test-tube. The liver was adherent to the stomach, its capsule was much thickened, and microscopic examination showed a slight cirrhosis in places advancing from the surface. The gall-bladder was distended with about four ounces of pus, the cystic duct being plugged with a calculus which was solitary. Kidneys: The pair weighed ten ounces; the capsule was thickened and adherent, and the cortex rather narrow. Microscopic examination showed a moderate amount of interstitial nephritis. The spleen was firm and dark, and weighed three ounces.

Birmingham.

A Mirror

OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

ST. BARTHOLOMEW'S HOSPITAL.

DISEASE OF THE KNEE-JOINT, SECONDARY TO SYNOVIAL CYST; AMPUTATION.

(Under the care of Mr. THOMAS SMITH.)

THE following case is one in which acute arthritis followed the formation of a large synovial cyst of the kind to which special attention has been drawn by Mr. Morratt Baker. Independently of the rarity of the case, the condition of the knee-joint, as found after removal, is of considerable clinical interest. Before amputation there was no swelling along the track of the cyst (which had been previously drained), the swelling of the joint was not excessive, and the temperature was normal. The extensive disease of the joint at the time of amputation was only foreshadowed by the excessive pain experienced by the patient on the slightest movement of the limb or even jarring of the bed; he was in a constant state of terror, unable to sleep, and begged for the removal of his limb. For the following notes we are indebted to Mr. H. E. Knight, dresser to the case:—

A. B—, aged forty-five, bootmaker, was admitted into Henry ward on August 2nd, suffering from a swollen left knee-joint. In June, 1885, he was in the hospital, and a synovial cyst of Baker (supposed before opening to be an abscess) was opened in the calf of his left leg; about one