septicemia is the cause of death rather than the meningitis. The clinical significance of these facts seems great, and the authors are seeking data to ascertain whether the withdrawal of cerebrospinal fluid during a septicemia and the subsequent development of a meningitis holds in the human subject also.

França, Carlos. CHEMICAL TREATMENT OF MENINGITIS. [Compt. Rend. Soc. de Biol., 1917, LXXX, p. 422.]

In the year 1902, França introduced the treatment of nontubercular meningitis by intraspinal injections of lysol solution. His method has proved very effectual in severe cases; the injections are well borne, and no bad effects follow. In cases of the epidemic form the lysol acts as an antiseptic agent, and the diplococci quickly disappear from the spinal fluid. The method he uses is as follows: After removal of 25 to 50 c.c. of spinal fluid by lumbar puncture, he injects intraspinaly a 1 in 100 lysol solution, from 12 to 20 c.c. for adults, and from 3 to 9 c.c. for children. If the patient's state be very grave, daily injections are given until the spinal fluid becomes sterile; this usually occurs rapidly. The only sequela seen is a yellowish tint of the palms and soles. In cases of purulent meningitis the lysol injection is preceded by lavage with normal saline solution. After the lysol injections the patient is put into an inclined position, with his head lowered. Urotropine is given as an adjuvant. França advises that his method should be used in cases of epidemic cerebro-spinal meningitis when, in spite of serum treatment, the course of the meningitis is slow and the meningococcus persists in the spinal fluid, and also in all bacterial forms of meningitis, with the exception of the pneumococcic and the tubercular. [Leonard J. Kidd (London, England).]

Netter, A. TARDY RELAPSES IN MENINGITIS. [Bull. d. l. Soc. Médéd. Hop., May 31, 1918.]

Netter discusses the probable causes for relapse in a number of meningitis patients who had apparently recovered, 1.14 per cent. out of 350 cases. Intercurrent measles was the cause in two instances. Anti-typhoid vaccination seems either to cause a reaction which simulates meningitis or which favors the proliferation of meningococci still present in the nasopharynx and otherwise harmless. In one case paratyphoid B followed soon upon meningitis and was in turn followed by meningitis and recovery then ensued under serotherapy. He says that Sainton reports a syncope after an intravenous injection of antimeningococcus serum, the patient being revived by artificial respiration.