GEOLOGISTS' ASSOCIATION JUBILEE VOLUME.

CHAPTER I.

MIDDLESEX AND HERTFORDSHIRE.*

By JOHN HOPKINSON, F.L.S., F.G.S., F.Z.S.

MORE than three centuries have elapsed since John Norden, in his *Speculum Britannia*, gave descriptions of "Middlesex and Hartfordshire,"† and he appears to have been the first to notice the difference in the soil of these two counties due to one being mostly on clay and the other on chalk. Of Middlesex he says that "The soyle is excellent, fat and fertile and full of profite. . . . Yet doth not this so fruitfull soyle yeelde comfort to the wayfaring man in the wintertime, by reason of the claiesh nature of soyle: which after it hath tasted the Autume showers, waxeth both dyrtie and deepe." And of Hertfordshire: "Concerning the Soyle: It is for the most part, chalkie, though the upper cruste in the South and West parts, be for the most part of redde earth mixed with gravell, which yet by reason of the white marle under it, yeeldeth good wheat and oates."

Middlesex is almost entirely on the London Clay, Hertfordshire chiefly on the Chalk, with the London Clay in the south-east, but in both counties these formations are much covered up by superficial deposits, mostly by gravel of varying origin, here and there by boulder-clay or brick-earth, while a great part of the Chalk is covered by clay-with-flints.

Upon the London Clay on certain hills there is Bagshot Sand, between the London Clay and the Chalk the Reading Beds come in, the Chalk is everywhere underlain by the Gault, a trace of the Upper Greensand coming in however here and there, and beneath the Gault on the extreme north and south is

* Including contiguous parts of Buckinghamshire, and excluding the north-east of Hertfordshire, for which see Cambridgeshire.

† "Middlesex" (1593) was the first part of the *Speculum Britannia*, *Hartfordshire* (1598), the second, and the two, with the "Preparative" (1596), were printed together in 1673, a third edition, here quoted from, being issued in 1723.

the Lower Greensand, which both ways soon thins out to nothing; then there is a wide gap in the downward succession, the Gault reposing in some places on Jurassic rocks, possibly in some places on Carboniferous, in some, both in Middlesex and Hertfordshire, on Devonian, and in the south-east of Hertfordshire on Silurian.

As the Gault and the strata above it dip to the south-east, and Hertfordshire is north of Middlesex, it has the older rocks on the surface, and therefore in taking the strata in the usual ascending order it will generally come first.

SILURIAN AND DEVONIAN.

Our knowledge of the presence of Silurian rocks under newer strata in the south-east of England is due to the *Wenlock Shale* having been met with in the boring of the New River Company at the Broad Mead Well between the Chadwell Springs and Ware. We have twice visited this spot: in June, 1878, when the boring was in operation, and in May, 1893, when water was being pumped from the well to the extent of about a million and a-half gallons a day. On the first occasion the diamond rock-boring drill was in operation, piercing the Chalk, and a depth of 250 ft. had been reached; and in less than a year the boring struck the Wenlock Shale at a depth of 796 ft. The strata passed through, after about a foot of made earth, etc., were River Gravel, 16 ft.; Chalk and Upper Greensand, 619 ft.; and Gault, 160 ft.; and the Wenlock Shale, with intercalated bands or lenticular masses of limestone and many fossils, was subsequently pierced for 36 ft.; the total depth reached being 832 ft.

That Palæozoic rocks were present under London was known before this boring was commenced, for in 1877 strata of *Upper Devonian* age were encountered at a depth of 1,064 ft. at Meux's Brewery at the corner of Tottenham Court Road and Oxford Street. Again, soon after the completion of the Ware boring, the Upper Devonian was reached at Turnford near Cheshunt at a depth of 980 ft. So early as 1855 old rocks had been proved to occur under London, but it is still uncertain whether these, the red rocks of the Kentish Town boring, 1,114 ft. beneath the surface, are of Old or New Red Sandstone age; or they may even be Carboniferous.

In commenting in 1879 on the discovery of the Wenlock Shale at Ware, the present writer remarked:

"We have, therefore, under the London Basin an axis of Palæozoic rocks, two divisions of which have now been discovered, namely the Upper Silurian and the Devonian. This axis has been inferred to be a prolongation eastwards of the

Palæozoic ridge which skirts the coal-basin of South Wales, then forms the Mendip range, disappears under Secondary rocks near Frome, and, after passing under London and the south-east of England generally, is possibly, under somewhat different conditions, the same ridge which is known as the axis of the Ardennes. The rocks of which it is composed are presumably, from their known character in their prolongation east and west, much contorted, and within the folds of the Devonian rocks it is *possible* there may be beds of coal, for elsewhere Carboniferous strata, with productive coal-measures, accompany them in their contortions, to the extent, as in the Auchy-au-Bois coal-field, of actually underlying them, the Devonian being folded over the Carboniferous strata. The strike of the old rocks, between London and the Mendip Hills, would according to this view be due east and west. The fossils from the Silurian rocks of the Ware boring are however typical of the Wenlock Shale as met with at Dudley and Wenlock Edge, seeming to show that the strike of the Silurian rocks at least is north-west and south-east, in which case we may have coal-beds on the northern flanks of this anticline as well as on the southern in the synclinal trough, or within folds in the Devonians between this axis and that of the Mendips and Ardennes to the south. The Ludlow rocks might be expected to occur under Hertford, and it would then become an interesting question as to whether the next series, occupying that part of the old land-surface lying between Hertford and Kentish Town, would be, as at Messrs. Meux and Co.'s boring, marine Devonians of Devonshire and Cornwall type, or "Old Red" beds similar to those which succeed the Ludlow rocks in Herefordshire. It appears more likely that a coal-basin would lie to the south-west if Old Red Sandstone beds of estuarine or lacustrine origin followed, than if the next series were of the marine Devonshire type." *

Since the above was written the Wenlock Shale in the Broad Mead Well has been determined to dip about 1° west of south at an angle of 41° with the horizon, and the Upper Devonian at Turnford about 17° west of south at an angle of 25° . †

JURASSIC.

Between the Gault and the Devonian rocks at Meux's Brewery, strata were met with, 64 ft. in thickness, which were at first considered to be Lower Greensand, but have since been determined by Prof. Judd to be *Great Oolite* (*Quart. Journ. Geol. Soc.*, vol. xl, p. 748). They thin out northwards, being 84 ft. at Richmond, and not being present at Kentish Town, probably ending not far south of the Kentish Town boring.

* *Trans. Watford Nat. Hist. Soc.*, vol. ii, part 7, p. 246 (1880).

† J. Francis, in *Rep. Brit. Assoc.* for 1895, pp. 451, 452 (1896).

CRETACEOUS.

The *Lower Greensand*, which comes to the surface in Bedfordshire, probably underlies the Gault throughout a good part of northern Hertfordshire. It has been proved to occur in the west of the county, at Long Marston, where it is 10 ft. thick under 215 ft. of Gault clay, but it is absent in the east, having thinned out to nothing before reaching Ware. It may be present also in the south of Middlesex, as it is 10 ft. thick at Richmond in Surrey, where it occurs 1,140 ft. beneath the surface, but it does not extend so far north as London.

The *Gault* underlies the Chalk throughout the whole of the two counties, except where it comes to the surface in two small areas in the west and north of Herts—west of Tring and north-west of Baldock. It varies from about 150 to 220 ft. in thickness in our area. It was seen in our excursion to Hitchin and Arlesey in May, 1900, the Upper Gault, unfossiliferous, being exposed to the depth of about 50 ft. at Messrs. Beart and Co.'s brickyard at Arlesey in Bedfordshire. On the eastern side of the large clay-pit a section was seen showing 10 ft. of the Chalk Marl and a thin band of the Cambridge Greensand reposing upon an uneven, evidently eroded, surface of the Gault. The pit is now worked only for the Gault clay, but formerly the Cambridge Greensand was extensively worked here, at Ashwell in Herts, and at other places in the neighbourhood, for the phosphatic nodules (coprolites) it contains, which were at one time used for making artificial manure. It contains two sets of fossils, the majority derived from the Gault, while some are indigenous, the latter being more closely related to the Chalk fauna than to that of the Gault. The bed passes up gradually into the Chalk Marl, and is regarded as the base of the Chalk.

The quarry of the Arlesey Cement Works was next visited. The upper part of the Chalk Marl, with the Totternhoe Stone and some 10 or 15 ft. of the Chalk above it, was exposed, and the Director, Mr. William Hill, pointed out that the Chalk Marl—the zone of *Ammonites varians*—was here about 70 ft. thick, the bluish-grey marl seen at the base of the quarry being the upward continuation of the whiter marl above the Cambridge Greensand. There is a considerable difference in the two marls, that just above the Greensand containing only 26 per cent. of argillaceous (insoluble) matter, while the bluish-grey marl contains 46 per cent. It was seen that this marl passed rapidly upward into a whiter and more calcareous deposit. The aspect of the Totternhoe Stone seen in the quarry was somewhat different from that at Totternhoe, where there is about 30 ft. of evenly-grained stone suitable for building purposes, while here the stone consists of two beds of rather rough and rugged (shelly) chalk, together about 12 ft. thick. The bed

forms the upper limit of the Chalk Marl, and above it is the zone of *Holaster sub-globosus*.

The succession of the strata is not here continuous, for the *Upper Greensand* usually comes in between the Gault and the base of the Chalk or Chalk Marl. It is of only local occurrence in this area. When present it is generally similar to the Cambridge Greensand in containing phosphatic nodules and fossils derived from the Gault, but its indigenous fossils are more nearly allied to those of the Lower Greensand than to those of the Chalk. In fact at Shenley Hill near Leighton Buzzard there is at the summit of the Lower Greensand a bed full of fossils which but for their position might be taken to be Upper Greensand species. The Gault and Upper Greensand constitute the Selbornian formation.

The succession of the various beds of the *Chalk* as they crop out in the escarpment on the north-western end of the Chiltern Hills has been well seen on our visits to the Dunstable Downs on several occasions (in 1881, 1889, 1903, and 1907).

Approaching this district from Tring Station, the village of Aldbury is seen to be picturesquely situated in a pretty valley cut out of the Chalk at the commencement of its escarpment by a stream no longer flowing. The Chalk Rock, dividing the Upper from the Middle Chalk, has here been cut through, cropping out near the middle of Moneybury hillside on the right, where the escarpment commences, and reappearing near the top of Aldbury Owers on the left, beyond which it ends, the hill on the right being much steeper than that on the left.

Beyond Aldbury the road leaves the valley to cross a spur of Pitstone Hill, where it passes over the lower portion of the escarpment, descending steeply to the plain beyond. Turning to the right just before reaching Ivinghoe, we are on the Icknield Way which leads to Dunstable, and at the corner we find a pit in the Lower Chalk, showing that in descending the hill the whole of the Middle Chalk has been traversed.

Proceeding along the Icknield Way, Ivinghoe Beacon Hill is seen on the right. The summit is 762 ft. above mean sea-level and about 300 ft. above the road where it is met by that from Ringshall. The hill is capped by an outlier of the Chalk Rock. The view from the summit in clear weather is very extensive and of much geological interest.

When on this hill at our cycling excursion on the 23rd of May, 1903, the view over the Gault plain was particularly clear, and the Director thus explained its geological features:*

"This extensive view embraces the whole of the Cretaceous rocks present on the northern margin of the London Basin

* In this and other similar reports of remarks a slight verbal alteration is occasionally made, and any error detected is either omitted or corrected. When no name is given, "the Director" implies the writer of this article.

except the higher beds of the Upper Chalk. We are standing on or just above the Chalk Rock, which forms the base of the Upper Chalk. On the summit of the hills behind are the lower beds of the Upper Chalk, the Ivinghoe Hills in the direction of Ringshall rising to 811 ft., the greatest elevation in this neighbourhood. For some miles along the Dunstable Downs, so far as the Five Knolls, four of which are distinctly visible, the Chalk Rock can be traced a little below the highest ground, its presence being indicated just beyond the upward termination of the steep face of the escarpment, nearly on level ground, by the resistance to denudation, due to its hardness, causing a slight ridge. Beyond the Five Knolls can be seen Totternhoe Castle Hill, the Melbourn Rock forming its summit, and just round the extreme point of that hill are the Totternhoe quarries (see Fig. 30, p. 163), where the Lower Chalk is now quarried for lime-burning, but the Totternhoe Stone at the base of the hill, for which the quarries were originally opened, is not now quarried as a building-stone. The position of the Melbourn Rock between Totternhoe Knoll and Ivinghoe Beacon cannot easily be traced, as it does not form a distinct feature in the landscape, but there is no difficulty in determining where the Totternhoe Stone occurs, for each little valley or coombe seen at frequent intervals all along the foot of the downs ends at its base, these valleys in some cases but not in all originating in a spring, owing to the water percolating through the Totternhoe Stone and being thrown out by the comparatively impervious Chalk Marl below it. This extends for some distance from the base of the escarpment, its surface being almost horizontal and its dip where it is exposed being less than when it passes under the great mass of the Chalk. On an isolated hill—an outlier of the Totternhoe Stone on the Chalk Marl—Eddlesborough Church is seen. The next village beyond is Eaton Bray, where a thin band of the Upper Greensand appears, and at a little greater distance in the same direction is Billington Hill with its church on the summit, that hill being an outlier of the Lower Chalk over the Gault of which the greater part of the plain below consists. Beyond this again the picturesque and well-wooded ridge of the Lower Greensand forms the horizon on the north from Leighton Buzzard for some distance across Bedfordshire towards Sandy and Potton. On the left, to the south-east, are the Tring Reservoirs which supply the Grand Junction Canal, the farthest being on the Gault and the other three on the Chalk Marl."

It was then shown, as pointed out by Mr. Whitaker long ago, that the Chalk escarpment is the result of sub-aërial denudation, not being a sea-cliff; and a view of an isolated terraced hill between the Downs and Eddlesborough, rising in a shallow valley which was traced up to a deep coombe, afforded an opportunity to broach a theory that some lynchets, as these terraces

are called, may be the remains of one side of a coombe, the scarps of lynchet and coombe being remarkably similar.

On this occasion, on descending the hill, a coombe just below it was examined, and its remarkably flat bottom, and regular, clearly-cut side-terraces were commented upon, a halt being made a little farther on to examine another, the bottom and sides of which are curved. Just beyond the point where the Icknield Way crosses the road from Hemel Hempstead to Leighton Buzzard, a closer inspection of one of these little valleys was made, but this is a true river-valley, not a dry coombe owing its origin to a time when the water-level in the Chalk was higher than it is now. At its head is a spring rising from beneath the Totternhoe Stone, and along its course other springs arise, for it flows nearly parallel with the strike of the rocks at the base of the Totternhoe Stone. Its sides are steep at first, then much less so, the ridge thus formed being believed to indicate the summit of the Totternhoe Stone, above which the Chalk, being softer, has a more gentle slope.

On our last visit to this district (15th June, 1907) the Beacon Hill was again ascended. The pit in the Lower Chalk near Ivinghoe mentioned above was then well worked for fossils, and several specimens of the zonal index-fossil, *Holaster sub-globosus*, were found, a few of *Terebratula semi-globosa*, and a "rag" bed full of fish-remains—teeth, scales, and pieces of bone. Its green-coated, brown phosphatic nodules mainly consist of material not differing much from the chalk in which the "rag" occurs, being much less phosphatic than those of the Cambridge Greensand. A local fault in this pit is well shown by the "rag" bed being thrown down to the right, and a slanting streak of crushed chalk in the face of the pit clearly indicates the line of fault.

On our return to Tring Station a chalk-pit under Pitstone Hill was examined. The junction of the Lower and Middle Chalk is here well shown, with the two *Belemnitella* marly bands separated by hard white chalk (about 3 ft. in all) forming the top of the Lower Chalk. In these beds the characteristic fossil, *Actinocamax plenus*, was found, and also *Ostrea vesicularis*. The hard, massive, nodular Melbourn Rock, 9 ft. or 10 ft. thick, forming the base of the Middle Chalk, was seen above the marly bands, and in and above the Melbourn Rock several specimens of the zonal index-fossil, *Rhynchonella cuvieri*, were found.

A little farther on, on the opposite (right-hand) side of the road, near Folly Farm, there is a pit in the *Rhynchonella cuvieri* zone, this species being common, and *Inoceramus mytiloides* being abundant, though mostly in fragments. Both these fossils were found on this visit, and Mr. Henry Kidner reported that he had also found in this pit *Conulus (Echinoconus) sub-rotundus*, *Discoidea dixonii*, *Inoceramus latus* (d'Orb.), *Terebratula bipectata*, *Camerospingia sub-rotunda* (Mant.), and *Ventriculites* sp.

It will thus be seen that the Dunstable Downs present a good field for the study of the physical features and for the examination of the zonal divisions of the Chalk from its base to the top of the Chalk Rock.

The Middle Chalk was formerly considered to terminate with the Chalk Rock, where present, but this—which is in the zone of *Holaster planus*—is now usually taken as the base of the Upper Chalk. A large chalk-pit on Rough Down, Boxmoor, visited in May, 1889, shows a thin band of the Chalk Rock near the top, most of the pit being in the higher portion of the Middle Chalk—the zone of *Terebratulina lata*,* which succeeds that of *Rhynchonella cuvieri*. On this occasion the Chalk Rock was described by Dr. Morison as “a cream-coloured rather rubbly limestone, jointed at right angles to the plane of bedding, breaking with a more or less conchoidal fracture, and weathering on exposure to the air. It contains,” he said, “numerous grains of glauconite and irregularly-cylindrical branching cavities filled in many cases with a ferruginous sandy clay or containing nodules of iron-pyrites. It varies in thickness from a few inches to 10 ft., but here it is only 18 ins. thick. In some places there are two bands a few feet apart.† The fauna of the Chalk Rock is peculiar, being notable for the number and variety of the Gastropoda; the Cephalopoda also are tolerably abundant; and the sponges are very numerous and varied. It contains more than twice as much magnesium-carbonate as ordinary soft chalk, nearly twice as much silica, more than three times as much sulphuric acid in the form mainly of sulphate of calcium, and a relatively very large amount of phosphoric acid in the form of phosphates. Examined microscopically it is seen to contain many small crystals, its hardness appearing to be mainly due to its crystals of calcium-carbonate, in fact to the conversion of the soft chalk into a crystalline limestone; sometimes large crystals of calcite are formed. It must have been deposited in a comparatively shallow sea, a considerable elevation of the sea-bottom taking place before its deposition.”

It may be added to Dr. Morison's remarks that a list of the Chalk Rock fossils found in this pit is given in Mr. Whitaker's *Geology of London*, vol. i, p. 73 (1889), and in Mr. Jukes-Browne's *Cretaceous Rocks of Britain*, vol. iii, p. 228 (1904), both being Memoirs of the Geological Survey.

The next higher zone—that of *Micraster cor-testudinarium*—comes on at the top of the Rough Down pit. Farther south the two railway tunnels between Hunton Bridge and Watford must be in it. In April, 1872, during the construction of the newer tunnel, we visited the cutting at its south-eastern end, and

* This species was formerly known as *T. gracilis*, var. *lata*.

† In that case the two hard beds with the loose rubbly chalk between them, forming together the zone of *Holaster planus*, may be from 15 to 20 ft. thick, as near Dagnall and Dunstable.

found the chalk to be deeply "piped" and most of its flints rough and hollow, their cavities containing sponge-spicules and other microscopic fossils. It would therefore appear to be in the upper part of this zone. An old chalk-pit in Berry Wood near Aldenham, which we visited in June, 1870, and May, 1875, is probably higher still in the same zone. In it we found the cavernous nature of the flints much more pronounced, the chalk being reported by the Director as abounding in the remains of sponges, surrounded by mere shells of flint, and containing spicules, Foraminifera, and Polyzoa, beautifully preserved.

Between Bushey Station and Watford, east of the line and well seen from it, is a chalk-pit in the next higher zone—that of *Micraster cor-angulum*. This was visited on the earlier of the above occasions, and also in May, 1879, and April, 1907. Its zonal position has only recently been determined with certainty by the finding in it of characteristic fossils by members of the Association. From the presence of *Micraster cor-testudinarium* it had been referred to the lower zone, but this species ranges into the higher zone whose index-fossil is also found in it, so in 1904 Mr. Jukes-Browne* suggested that it might be in the zone of *Micraster cor-angulum*, and this was confirmed on our last visit. Its flints, occurring in layers throughout, are mostly solid. The depth of the section is nearly 60 ft.; the lower part, in which nearly all the fossils occur, is white chalk; the higher part (about half) is yellow banded chalk.

The chalk-pits near Harefield, close to the River Colne, below Rickmansworth, expose as fine sections of the Upper Chalk as any to be met with on the northern side of the London Basin, and nowhere can better examples of pipes in the Chalk be seen than in one of these pits, while the absence of any vestige of a pipe in another affords an instructive lesson in physical geology. There are three large pits within a distance of a mile and a half, to one or other of which at least one visit has been paid (in 1882, 1894, and 1899). All are cut out of the hillside on the left-hand bank of the Colne, and may be best reached from Rickmansworth by the towing-path of the Grand Junction Canal.

The first pit, near Springwell Farm (Plate I, Fig. 1), presents a nearly unbroken surface of chalk of a pure white, relieved here and there by layers of flints, the clay-with-flints which comes on above having thrown off most of the rain-water falling upon it, thus preventing it from freely percolating into the chalk and forming large "pipes."

In the next pit, about a field's length farther south, near the Asbestos Mills (Plate I, Fig. 2), the whole section of the chalk, which presents an almost vertical cliff from 90 to 100 ft. high, is capped by gravel, from which pipes, often of most fan-

* *Mem. Geol. Surv.*, "Cretaceous Rocks of Britain," vol. iii, p. 231.

tastic shape, and of roughly-cylindrical form, extend downwards to distances varying usually from 30 to 70 ft. On our first visit to this pit Mr. Whitaker explained the mode of origin of these pipes, stating that they were caused by rain, which, in passing through air, absorbed carbonic acid gas and thus acquired the power of dissolving the hardest limestone. The rain-water with this gas in solution sank down through some line of weakness along which it gradually dissolved the chalk until the overlying gravel and sand sank into the cavity thus formed. Here and there a mass of gravel was seen which appeared to be unconnected with a pipe above it, this appearance being due to the chalk not having been worn away in a vertical line, so that the connecting link of the pipe was not to be seen. In these pipes there might frequently be seen angular chalk-flints, not worn at all, and nearly in the position they occupied before the chalk in which they were embedded was dissolved away.

The chalk in the third pit, that of the Harefield Brick and Cement Works, being capped by Tertiary strata, is not piped.

In all three pits the chalk seems to be mostly in the zone of *Micraster cor-anguinum*; the lower part of the first pit in that of *M. cor-testudinarium*; and the base of the second pit also in that zone. Flint frequently occurs in tabular layers. Of the ordinary flints most are solid, a few cavernous.

The following table showing the divisions of the Chalk in Middlesex and Hertfordshire may be useful for reference.* Of the Upper Chalk, the lower portion—that with *Micrasters*—only is included, as probably none of the upper portion—that with *Belemnites*—occurs in either county. The figures are the approximate thicknesses of the beds in this area.

DIVISION.	ZONE.	NATURE OF THE STRATA.
UPPER CHALK (Senonian), 180-280 ft.	<i>Micraster</i>	{ Firm yellow and white chalk with frequent layers of flints.
	<i>cor-anguinum.</i>	
	<i>Micraster</i>	{ Softer white chalk, with sparser layers of flints.
	<i>cor-testudinarium.</i>	
MIDDLE CHALK (Turonian), 220 ft.	<i>Holaster planus.</i>	{ Chalk Rock. Chiefly hard nodular chalk. 10-20 ft.
	<i>Terebratulina lata.</i>	{ White chalk with few flints, hard at base. 160 ft.
	<i>Rhynchonella cuvieri.</i>	{ Chalk without flints. 50 ft.
		{ Melbourn Rock. Hard nodular chalk. 8-10 ft.
LOWER CHALK (Cenomanian), 160-180 ft.		{ Belemnite (<i>Actinocamax plenus</i>) marl. 3 ft.
	<i>Holaster subglobosus.</i>	{ White and grey chalk. 55-75 ft.
		{ Totternhoe Stone. Chiefly hard sandy stone. 20-22 ft.
	<i>Ammonites varians.</i>	{ Grey chalk, chalk marl, chloritic marl, and coprolite bed. 80 ft.

* In the Geology of Herts. (1902) and Beds. (1904) in the *Victoria County History*, the writer included the Chalk Rock in the Middle Chalk; it is here placed in the Upper Chalk in accordance with the classification now adopted by the Geological Survey.

[Photo by J. Hopkinson.]

1.—CHALK-PIT NEAR SPRINGWELL FARM, HAREFIELD.

[Photo by J. Hopkinson.]

2.—CHALK-PIT NEAR THE ASBESTOS MILLS, HAREFIELD.

Phenomena connected with the flow of water in the Chalk have been studied in excursions to the Hertfordshire Bourne between Berkhamsted and Boxmoor, and to the swallow-holes in the bed of the River Colne from Mimms Hall near South Mimms, a little below the head of the river, to Potterell's Park near North Mimms.

The Hertfordshire Bourne is an intermittent stream only flowing after an exceptionally heavy rainfall. The years for which we have records of its flow are 1853, 1860, 1866, 1873, 1876, 1877, 1879, 1881, 1882, 1883, 1897, and 1903-4. It usually commences to flow early in spring and terminates in summer. It has never but once been known to flow when the mean rainfall in the county for the year ending 31st March has been less than 30 inches, the exception being in 1882 during a period of very heavy rainfall; and it has only twice for half-a-century at least failed to flow when the rainfall has exceeded 30 ins.; in fact we can only be certain that this has occurred once, and that was in 1892 when the year's rainfall (to 31st March) was 30.21 ins. Only on the last occasion did it begin to flow before the 1st of January, and that was in the wettest year of our Hertfordshire record.

The Bourne has been visited by the County Society nearly every time it has flowed since the year 1876, but only once (on 21st April, 1883) has the Geologists' Association visited it. The stream was then encountered where it is crossed by Harratt's End Lane, and where it had its source when visited by the Hertfordshire Natural History Society in 1881. On this occasion, however, it was flowing here in considerable volume, the field above the lane was in great part submerged, and for the source we had to walk about half-a-mile up the valley by the side of the lake thus temporarily formed. Here Sir John Evans explained the geological and meteorological conditions which give rise to this and other intermittent streams.*

The course of the Bourne was then followed to Bourne End, where the stream flows into the Bulbourne. In a portion of its course it was found to have taken temporary possession of the road, a *détour* having to be made into the adjoining fields to avoid it, and at Bourne End it was observed to be flowing through the usually dry culvert under the high road in a powerful stream.

The Colne swallow-holes were visited on the 19th of April, 1890. Although March was rather a wet month, February was very dry, and scarcely any rain fell in the early part of April, so these swallow-holes were not seen in operation, the bed of the river below Mimms Hall being dry. The largest cavities are in Potterell's Park, where the then dry bed of the stream is very devious, and a considerable area of the land has consequently a very uneven surface.

* *Proc. Geol. Assoc.*, vol. viii, p. 120.

In the present year (1908) swallow-holes were seen in action in the bed of the stream between Mimms Hall and Water End, above Potterell's Park, and it was noticed that after the water had flowed into the swallow-hole it reappeared lower down the river, showing that the plane of saturation of the Chalk was just about level with the bed of the stream. March was a very wet month, and our visit took place on the 11th of April. A few days earlier the river was flowing in force, and the road by its side was flooded near Water End.

Eocene.

Of the three divisions of the Lower London Tertiaries, the Thanet Sands, Woolwich and Reading Series, and Oldhaven Beds, the second only is present north of the Thames, except in part of South Essex. Although extremely variable in composition, the *Woolwich and Reading Series* is very persistent, resting, wherever the Thanet Sands are absent, directly on the Chalk. There are three types of this series, of which we have the Reading type only; the term Reading Beds will therefore alone be used. This type, unlike the Woolwich, has very few fossils.

In the absence, in our district, of the Oldhaven Beds, which are of only local occurrence, the Reading Beds are immediately succeeded by the *London Clay*, and as in almost every section both these formations are seen, they will be treated together, first along the outcrop of the Reading Beds from north-east to south-west, and then their outliers will be taken in the same direction. An occasional allusion to the underlying Chalk will necessarily also be made.

Easterly the Reading Beds are first seen in Hertfordshire at Bishop's Stortford, which was visited in July, 1897, and again, with Stansted in Essex, in July, 1905.

On our first visit the Rev. Dr. A. Irving pointed out that the pre-Tertiary plane of abrasion of the Chalk was shown by its outcrop on both sides of the Stort Valley, and by measurements taken in well-sections to the east and west, to be at about 190 ft. O.D. Well-sections in the valley show that along that line of erosion the Chalk underwent such extensive sub-aërial waste in late Tertiary and perhaps also in early Glacial times, as to be channelled to a depth of 190 ft., 170 ft. of which depth is now filled up with glacial detritus.

The Reading Beds are seen on both sides of the Stort Valley, where there are extensive brickyards. Green-coated flints are common in their basement-bed. In the Rye Street brickyards (on west side of river) these beds show signs of extensive erosion. On the slopes of the valley the London Clay is much obscured by drift. A good section of it was seen at the top of the field adjoining the brickyards of Mr. J. L. Glasscock. The workmen

assert that below the present bottom of the pit there is a bed of flint-pebbles. This bed occurs in some clay-pits at Birchanger, about half a mile distant, and there contains many oyster-shells, some of which are much worn by sand and water. Sharks' teeth occur at both localities.

On our second visit to this neighbourhood, in the old chalk-pit at the Castle Hill, Stansted, a fault was seen which had equally affected the Chalk, the Reading Beds, and the overlying gravels. It has a downthrow of about 6 ft. to the north. The high ground of Birchanger to the east is capped by London Clay of which bricks have been made. The sections are now obscure, but they formerly yielded sharks' teeth, septaria, etc. The base of the formation, where the sands of the Reading Beds were dug out, is marked by a band of flint-pebbles and oyster-shells. At Mr. Day's brickworks in Bishop's Stortford the clunchy character of the bed with green-coated flints reposing immediately upon the Chalk was noticed, and it was stated that specimens of *Micraster cor-anguinum* had been obtained from the pit, indicating the zone of the Upper Chalk upon which the Reading Beds lie.

The Reading Beds and basement-bed of the London Clay are well exposed in Mr. Lines's brickfields near Rush Green, Hertford, to which visits were paid in 1878 and 1885.

On our first visit Prof. John Morris called attention to the presence in a chalk-pit near the brickfields of a bed of green-coated flints, which he said was a proof that we had the true surface of the Chalk, this bed forming the base of the Tertiary Series. Sharks' teeth and oyster-shells (*Ostrea bellovacina*), Mr. Lines said, were frequently found in it.

Sections exposed in the brickfields were then examined, and Prof. Morris stated that the Reading Beds here, consisting of sands and clays, showed very different conditions from those south of London, where there were 30 or 40 ft. of ash-coloured sands. Here there were no freshwater shells, these deposits in the north and west being marine, though they were contemporaneous with the freshwater and estuarine beds south of London, which at Lewisham, etc., contained a great number of shells. The next higher bed seen here—the basement-bed of the London Clay—was of interest as representing an important change of conditions, its black flint-pebbles being derived from flints rolled on some sea-shore, giving evidence of a great depression over a large area extending between Marlborough, Hungerford, and Harwich.

On our second visit a more detailed examination of the same beds was made, and the section as then seen was compared bed by bed with its appearance when examined by Prestwich in 1854 and by Whitaker in 1869, their descriptions of it being given by the Director.

In 1854 the section, continued into the chalk-pit, was : *

		ft.
	Mixed clay and gravel	1
LONDON CLAY.	{ Basement-bed. Brown, sandy clay laminated with tougher clay, containing casts of <i>Panopæa</i> , etc.	
	{ Flint pebbles in clay at base (6 in)	6½
	{ Light greenish sand	4
READING BEDS.	{ White or ash-coloured sand	6
	{ Light, yellow sands	4
	{ Light-coloured mottled, red and grey clays	10½
	{ Yellow sand, with patches of <i>Ostrea bellovacina</i> in places at its base	6½
	{ Green-coated flints	1
		39½
CHALK		12

In 1869 the section was as in Fig. 1.

With the exception of a slight difference in the thickness of the highest layers of the Reading Beds, the only alteration since 1869 appeared to be that there was a continuous section from the London Clay downwards, the "smaller pits" being extended into the large pit in which the London Clay is exposed.

The Director prefaced these descriptions of the section with the following observations on the London Tertiary Basin :

"It may be well to explain that the term 'basin' is here used as a convenient expression for a series of beds originally deposited horizontally and subsequently let down in the centre by the sinking of the adjacent strata. The London Basin therefore is not a hollow in the Chalk in which the Tertiary strata were deposited, in which event the surface of the Chalk would be curved and the Tertiaries would be horizontal ; but it is a series of strata the members of which are conformable to, or parallel with, each other, and have only a very slight curvature caused by a folding or waving of the strata, which occurred after, and probably soon after, the deposition of the Eocene or Lower Tertiary beds. It must not however be inferred that the Chalk, after its quiet deposition by organic agencies in an ocean-bed, remained submerged throughout its extent until the lowest Tertiary beds, the Thanet Sands, were deposited upon it, for a considerable area was probably for some time land, and on rising from the bed of the ocean was subjected to denuding agencies, such as rain and rivers, similar to those now existing. In this neighbourhood there is evidence of a large amount of chalk having been washed away, and of the flints having been washed out of it, broken up, rolled into pebbles, and dispersed over wide areas, before it sank altogether beneath the ocean. The Thanet Sands are absent here ; the higher beds of the Upper

* Prestwich, in *Quart. Journ. Geol. Soc.*, vol. x, p. 92.

FIG. 1.—GENERAL SECTION AT MR. LINE'S BRICKYARD, NEAR HERTFORD.—*W. Whitaker.**

Chalk are also absent, and it is therefore probable that the Chalk was for a longer period a land-surface here, and on the north-western margin of the London Basin generally, than on the south-eastern margin, either from its greater elevation or later depression, being therefore subjected to denudation for a greater length of time. The lower and main portion of the Reading Beds here exposed consists almost entirely of fine-bedded sand,

* "Geol. London Basin, *Mem. Geol. Surv.*, vol. iv, p. 227.

showing that it was exposed to strong currents, and therefore furnishing additional evidence of the proximity of land."

Hatfield Park Kiln has twice been visited—in 1873 and 1876—by permission of the Marquess of Salisbury.

There are two pits, one at a higher level than the other, which together expose a section of the London Clay and Reading Beds from high-level post-Pliocene gravel to the Chalk. A description of the section is given in Fig. 2.*

The next exposure of the Reading Beds we have seen along their outcrop is one of great interest. In Newberries Park, Radlett, there are pebble-beds consolidated into a conglomerate known as the Hertfordshire conglomerate or "plum-pudding-stone." The exposure is in a pit on the hillside east of Radlett, and has been examined by us twice—in 1884 and 1890.

On the first occasion the section was far from clear, the pit being disused and much overgrown, but the conglomerate was seen in several places, with beautifully white, fine sand reposing upon an unconsolidated portion of the pebble-bed.

After giving a general account of the geology of the district, the Director said that the character of the Reading Beds here indicated a marine origin, partly at least littoral; and proceeded: "It is this shore-deposit that we have now before us, for the Hertfordshire conglomerate is merely a bed of the Reading Series—the shingle-bed of flint-pebbles—consolidated by the infiltration of silica. Blocks of this conglomerate occur here and there over a considerable portion of Hertfordshire, and beyond. We know that the Reading Beds once extended over the greater portion of the county, by the outliers which are seen at various places, and it seems more probable that the blocks we meet with are relics of these beds, the looser portions of which have been carried away by denudation, than that they have been transported in a northerly direction from the neighbourhood of Radlett. It is probable that the land over which the rivers flowed which deposited the clays, sands, and pebbles of the Woolwich and Reading Series was to the south of the London and Hampshire Tertiary basin, where the English Channel now is, and on this supposition we should expect to see the beds to the north more marine in character than the more southerly portion of the series. From the variable character of the whole of the series it may be inferred that in no locality where it occurs was it, or any portion of it, a deep-sea deposit. Littoral conditions may have prevailed over at least a considerable portion of Hertfordshire, so that there is no improbability in the shingle-bed having at one time been of considerable extent, and throughout this extent it may anywhere have been consolidated into a conglomerate. This is here seen to be split up into blocks, and the durability of the siliceous cement is

* *Mem. Geol. Surv.*, "Geology of London," vol. i, p. 205.

evidenced by the splitting of the pebbles. In some places the pebbles have become so soft that they can be cut with a knife, while the matrix preserves its hardness. Another point of interest is that here and there the surface of the conglomerate is seen

FIG. 2.—GENERAL SECTION OF THE TERTIARY BEDS AT HATFIELD PARK KILN.—*W. Whitaker.*

(Scale, 8ft. to an inch.)

to be smooth and rounded, as if it had been subjected to the action of ice. It is the only rock we have anywhere in this neighbourhood, and probably in our whole Tertiary basin, which is sufficiently durable to show any trace of glacial action."

FIG. 3.—SECTION FROM NORTH-WEST OF BUSHEY TO STANMORE HEATH.—*W. Whitaker.**Horizontal scale, 3 inches to a mile; vertical scale, twice as large (1 inch to 880 feet).*

The coming on of the London Clay over the Reading Beds has been made too abrupt, and the ground at and to the left of Bushey has been drawn rather too high, so that the slope on the right is hardly exaggerated.

On our second visit to this spot the pit had been opened up by Mr. Henry Lubbock in order that the section might be better seen. The conglomerate now appeared to be in large and small blocks embedded in clay, overlying a stratum of black-coated pebbles of irregular size intermingled with sand and clay.

Mr. Upfield Green, the Director on this occasion, said that he believed the Reading Beds were not here *in situ*, but were rearranged and intermixed with detritus of the London Clay washed down from the higher ground by the stream which now flows at a much lower level. The composition of the conglomerate varied greatly, some specimens partaking more of the nature of breccia than conglomerate, and while the inclusions in some consisted almost entirely of pebbles of the Reading Beds, in others angular and sub-angular fragments of flint and chert, with rolled pebbles of flint and clay silicified, were largely present. In some cases the silica cementing the sandy matrix had attacked and partially dissolved the pebbles. He expressed the opinion that the blocks were of fluvial origin and formed at the debouchure of streams flowing northward, which, draining the London Clay, cut through the Reading Beds, depositing the detritus either in subterranean streams or in the bed of a pre-Glacial river. The silica subsequently cementing these rearranged materials he considered to have been derived from

decomposition of the soluble silicates of the felspathic constituents of the Glacial drift.

The Hertfordshire conglomerate has also been examined in the neighbourhood of Watford, in one or other of the Bushey brickfields (see Fig. 3) to which visits were paid in 1870, 1872, 1899, and 1907. It was on the last occasion only that the brickfield in which this bed is exposed was visited, so for continuity of subject this will be taken first.

It is in Lower Paddock Road, and is known as Grover's Brickfield, but it has not been worked for some time. The exposure of Hertfordshire conglomerate, $2\frac{1}{2}$ to 3 ft. thick, is of interest as continuous with and merging into loose pebbles and sand, with pale sand below, seen for 3 or 4 ft. in depth, and containing a layer of flint-pebbles. Although the beds above could not be seen, the conglomerate is evidently *in situ*, forming an integral part of the undisturbed Reading Beds. On the east, level with the pebbles and sand, were seen the clays of the Reading Beds, and above them the basement-bed of the London Clay with its band of flint-pebbles. This appeared to indicate a line of fault running approximately north and south, and having a downthrow of about 7 or 8 ft. on the east, by which the calcareous clay had been brought against the pebbles and sand. This conclusion was confirmed by pieces of the conglomerate broken off near the fault, which were found to have been altered to "fault-breccia," the pebbles having been crushed and reduced to angular fragments in the faulting, while the siliceous cement showed a whitish crystalline surface due to crushing, instead of the usual smooth fracture. The fault must have originated in the Chalk below.

The Hertfordshire conglomerate has been seen in the same position in the Reading Beds on their outcrop near North Mimms and near Radlett Church on the opposite side of the valley to Newberries. It also occurs in the outlier of these beds at St. Albans. It therefore appears that the pebble-bed has been consolidated by the infiltration of silica in several places in the county whilst in its original stratigraphical position.

Bushey Kiln on the high road to London, and Watford Heath Kiln in Oxhey Lane on the road to Pinner, were visited in 1870, 1872, and 1899. Although these brickfields are only three-quarters of a mile apart, Bushey Kiln being due north of Watford Heath Kiln, there is a very great difference in the composition and thickness of both the basement-bed of the London Clay and the Reading Beds exposed at each place, as will be seen from the sections (Figs. 4 and 5) on pp. 20 and 21.*

The shells at the bottom of the bed *b* at Watford Heath are mostly in fragments, but several perfect specimens of the following, with a few undetermined species, were collected on our

* *Mem. Geol. Surv.*, "Geology of London," vol. i, pp. 201 and 202.

FIG. 4.—GENERAL SECTION AT BUSHEY KILN
NEAR WATFORD.—*W. Whitaker.*

(Scale, 8 feet to an inch.)

Drifted clay, in parts sandy, with flint-pebbles,
thickness variable.

			ft.
LONDON CLAY.	(a)	Brown Clay.	
	Basement Bed.	(c) Pebble-bed ; some of the pebbles large	$\frac{1}{2}$ (or more)
		(d) Brown sandy clay ; "large oyster-shells at the bottom" (R. Trench)...	about 6

READING BEDS, 35 or 36 feet.	(f ₁)	Green clay, with "race" (irregular cal- careous concretions) pass- ing into the bed below.	over 7
	(f ₂)	Greenish and brown clayey sand	
	(f ₃)	Clay.	
	At the eastern part of the yard these three beds are so full of race that they be- come a marl.....		
	(g)	Light - coloured false- bedded sand	about 12
	The beds below this were not to be seen, but Mr. Trench was told that they were as follows :—		
	(h)	Pebble-bed	about 5
	(i)	Sand	4
	(k)	Loam	5
	(l)	Flints in sand	2

(m) CHALK.

FIG. 5.—GENERAL SECTION AT WATFORD
HEATH KILN.—W. Whitaker.

(Scale, 8 ft. to an inch.)

FIG. 6.—SECTION ALONG THE RAILWAY FROM PINNER TO RICKMANSWORTH.—J. Hopkinson.

The level of the line is approximately (from aneroid measurements) 190 feet above ordnance datum at Pinner, 230 feet (30 feet below the original surface) at the highest point, near Grove Green, and 200 feet at Moor Farm.

first visit: *Aporrhais sowerbyi*, Mant., *Cyprina morisii*, Sby., *Cyrena cuneiformis*, Fér., *Cytherea obliqua*, Desh., *Natica labellata*, Lamk., *Nucula* (? *margaritacea*, Desh.), and *Panopaea intermedia*, Sby. A collection was also made from tabular masses, or concretions, of calcareous-argillaceous sandstone occurring at intervals in this bed, but not seen in place, the section from which they were obtained not having been worked for some time. In the bed *d* several rare fossils have been found, including bones of a turtle and a tooth of a mammal, *Hyracotherium leporinum*, Owen.

On our second visit to Watford Heath Kiln a well had been sunk through the Reading Beds to the Chalk, just on the edge of the mottled clay, which here ends abruptly, its place being taken by the bed of sand on which it rests, and the basement-bed of the London Clay reposing in one part of the pit immediately on mottled clay and in another part on sand. At Bushey Kiln a small section, exposing several layers of pebbles in sand, showed how the difference between the beds here and at Watford Heath might be partly accounted for. Not only were these layers irregularly distributed, but between each more or less horizontal one were to be seen thin connecting lines of pebbles passing through the sand at various angles, showing how, previously to the deposition of

each bed of sand, strong counter-currents tore up the underlying pebbles and redeposited them.

By the extension of the Metropolitan Railway beyond Pinner, sections were here and there exposed of the London Clay and its basement-bed, the Reading Beds, and the Chalk. Cuttings in progress between Pinner and Rickmansworth were examined in June, 1886. There is a shallow cutting between Pinner and the high road to Rickmansworth, and just beyond this point a much deeper cutting commences, both being in the London Clay. A little farther on the basement-bed is exposed, having the usual bed of flint-pebbles very largely developed, and then, near Goose Green, just beyond the highest point which is about 260 ft. above mean sea-level, the Reading Beds crop out under the escarpment of the London Clay. A small stream rises here and has been diverted almost from its source to allow of the line being carried in the direction of its bed, thus taking advantage of the lowest level for the least amount of excavation nearly to the point where the stream falls into the River Colne. The line then emerges from the cutting, being continued to Rickmansworth on an embankment.

The accompanying section (Fig. 6) has been drawn along the course of the line, and shows the slight upheaval of the Chalk, by which, and the lowering of the ground by the denuding action of a small stream, the course of which has probably been mainly determined by the line of weakness or breaking across of the strata caused by the elevation, the Reading Beds have been laid bare, forming an inlier surrounded by the London Clay. The relation which this inlier appears to bear to the outliers of these beds will be stated presently.

The next place where we have seen the Reading Beds along their outcrop is Woodcock Hill Kiln near Rickmansworth, on the road to Harefield and Ruislip. The section here exposed was examined in 1882, 1887, and 1899. The beds in the main upper pit are as follows:

Clayey gravel resting in hollows in the bed below.	
BASEMENT-BED OF THE	{ Brown and grey sandy clay with a ferruginous bed in which a cast of <i>Panopæa</i> was found, ? 5 ft. Layer of flint-pebbles. Brown loam, about 3 ft. Mottled plastic clays, thick.
LONDON CLAY.	
READING BEDS.	

A small, lower pit shows about 10 ft. of brown sand, with clay above, and some apparently in it, evidently dipping under the mottled clays. In the presence of a pebble-bed in the midst of the basement-bed of the London Clay, and the absence of one at the bottom, the section presents a rather unusual feature.

The Harefield Brick and Cement Works were visited in April, 1894, and June, 1899. Here there is a very fine section

from the lower beds of the London Clay through its basement-bed and the Reading Beds into the Chalk. It has been given in detail by Mr. Whitaker in his "Geology of London" (vol i, p. 196). The basement-bed of the London Clay consists of about 6 ft. of brown rather sandy clay with two layers of decomposed shells and a few pebbles and also calcareous concretions with shells which are better preserved. The Reading Beds contain much clay, the greater part of which is bedded and mottled, mainly grey with red, with some sandy clay, and at the base a bed about 2 ft. thick of green-coated flint-pebbles passing down into unworn green-coated flints in greenish sand. Leaves of *Salix* have been found in lenticular masses of clay directly above the bed of flint-pebbles.

Farther to the south-west the Reading Beds and London Clay have been seen in cuttings in progress for the Great Western and Great Central Railways in the vicinity of Gerrard's Cross, in June, 1903, and April, 1905. On the second occasion the long cutting at Gerrard's Cross, through the high ground to the west of the River Misbourne, showed the following general section from the east end to the centre of the cutting :

	ft.	ft.
1. Plateau Gravel (sand and gravel)	from 16 to	25
2. London Clay (dark grey and brown sandy clay)	about	5
3. Reading Beds (blue and mottled clay)		15

The section at the west end of the cutting was as follows :

	ft.	ft.
1. Gravel	5 to	15
2. White, brown, and green sands		12
3. Clay, with flints at base	about	1
4. Chalk (visible in bottom of cutting).		

Attention was drawn to the fact of the band of London Clay displaying abrupt changes in thickness in places, and to its not being quite continuous up to the centre of the cutting, as in one place the gravel was seen to rest directly on the Reading Beds. It was suggested that the London Clay was perhaps redeposited, not being strictly *in situ*. Near the west end of the cutting some Sarsen stones were examined, a few being in place. One measured 4 ft. by 2 ft. 3 ins. by 2 ft. They usually occur in the white and brown sand, and a few have been found in the gravel.

Beyond the very irregular line of outcrop of the Reading Beds there is a series of outliers of the same, some with London Clay also, extending from Albury on the north-east to beyond Rickmansworth on the south-west, right across Hertfordshire and into Buckinghamshire. Their average distance from the main mass is about five miles. About four miles beyond these a second line of three, not so well defined, occurs; and about seven miles farther, again in an almost straight line in the same direction, there is a third line of five small outliers extending from Caddington in Bedfordshire to Aldbury Common in Hert-

fordshire. Thus it is seen that the London Tertiary Basin has extended at least sixteen miles beyond its present margin in this area, and probably its extent was at one time far greater.

These outliers appear to show that the dip of the Chalk is not quite uniform, having a series of slight undulations which are maintained for a considerable distance along the strike. If the synclinal folds occurred where there are now these lines of outliers and the surface became reduced to a level by marine denudation, the Tertiary Beds would be preserved from erosion in the synclinal folds, and subsequently, by their clays protecting the underlying Chalk from the dissolving action of carbonated water, would form the highest ground. Local circumstances, such as the presence of protecting sheets of gravel, would determine the areas along each ridge where the Tertiaries would eventually be preserved.

Within the present margin of the London Clay there appears to be an indication of another flexure along the strike of the Chalk parallel with these outliers, in two inliers of the Reading Beds, one at Northaw, the other at Pinner. These inliers seem to be due to an anticline of the Chalk, which, increasing to the south-west, has resulted in the upheaval of this rock where it forms the hill on which Windsor Castle is situated.

Of the outliers in the first line, those of Datchworth, Ayot Green, St. Albans, Chorley Wood, Coleshill, and Penn have been visited.

In the Datchworth outlier the basement-bed of the London Clay overlies the Reading Beds, but only the gravel which caps the Tertiaries has been examined.

Interesting exposures of the Reading Beds and basement-bed of the London Clay have been seen on several occasions in the Ayot outlier (in 1886, 1898, 1902, and 1906), in four pits in a large brickfield adjoining Ayot Station. The section made up from these pits in 1864 was as follows:*

	Sandy contorted pebble-gravel (at highest parts), 8 or 10 ft. shown, but a greater thickness said to have been found.
LONDON CLAY.	<div> <div>Brown clay, sometimes laminated, 8 or 10 ft. shown, but as much as 14 ft. said to have been found.</div> <div>Basement bed. Brown loam, mostly with green grains; a bed of pebbles, about 3 ins. thick, at or near the bottom, 4 to 4½ ft.</div> </div>
READING BEDS.	<div> <div>Light-coloured sand, for the most part nearly white, but partly brown and buff, with thin layers of pale grey clay, false-bedded at one place. The uppermost 6 to 12 ins., sometimes clayey and ferruginous; the middle part sometimes with lines of pebbles "mostly well-rounded, but some not; mostly black, but some green-coated" (Topley); 14 ft. shown, but said to have been worked to a depth of 20 ft. without reaching the Chalk.</div> </div>

* Whitaker, "Geology of the London Basin," *Mem. Geol. Survey*, vol. iv (1872), p. 237.

At our first visit the section was clear down to the Chalk, a depth of about 24 ft. of sand being exposed. The false-bedding was very evident. On our second visit it was seen that in some places an appearance of false-bedding was given by the irregular dissolution of the chalk below having affected the sands, setting them down irregularly. This was on the north side of the brickfield. On our last visit the various pits, examined in detail, gave the following results:

In the first section, north-eastern end of the brickfield—

1. Boulder-clay, with stones, not well exposed.
2. Gravel of flint pebbles, quartz-pebbles, blocks of quartz, subangular flints, etc.

In the second section, a little south of the first—

2. Gravel as above.
3. London Clay. Tolerably stiff clay of a brown colour, partly laminated; only 6 ft. seen, but 14 ft. said to have been exposed.
4. Basement bed of the London Clay. Rather sandy brown clay, with a layer of black flint-pebbles at the bottom.

In the third section, a little west of the second—

4. The above layer of pebbles close to the top.
5. Reading Beds. Brown, yellow, and orange-coloured sand, much current-bedded in places, with a little iron sandstone in irregular layers. Near the top, in one place, about a foot of pale grey clay and a few irregular layers of clay.

In the fourth section, western end of the brickfield—

5. Reading Beds. Sands as above, with some much lighter in colour, in places nearly white, with two thin layers of flint-pebbles about 6 ft. apart.

The teeth of several species of shark have been found by the writer in the basement-bed of the London Clay—in the bottom layer of flint-pebbles in sand—and are now in the Hertfordshire County Museum at St. Albans. Mr. E. T. Newton has obtained from a workman some teeth of the sharks, *Odontaspis elegans*, *O. cuspidatus*, and *Lamna vincenti*, and also of a teleostean, *Phylloodus toliapicus*, which must have come from the same bed.

In the St. Albans outlier, which we visited in May, 1892, the Reading Beds only are present. They were first seen in the brickyards near Bernard's Heath, and then in the gravel-pits on Townsend Farm, where the Hertfordshire conglomerate appears to be in place. It was traced for about 20 ft. in one direction and 5 ft. in another at right angles, lying in a horizontal position under the gravel, and with sand beneath it.

At the Chorley Wood outlier near Rickmansworth, the former brickfield has been taken into the garden of a private house, the occupier of which allowed us, in June, 1888, to see the section exposed. It is nearly on the edge of the outlier, which

is conspicuous from the valley below as a well-wooded eminence. Its chief interest is in showing an unusual development of the bottom bed of flint-pebbles. This bed is 12 ft. thick, and the pebbles, some so large that they might almost be called boulders, are closely packed together, only the interstices left by their spherical shape being filled up with sand.

The following is Mr. Whitaker's description of the Reading Beds seen in this section in 1859, resting on the Chalk :*

Plastic clay ("paving clay").

Clayey sand, brown and light-coloured, 1½ ft.

Red-mottled plastic clay ("tile-clay").

Bluish grey sandy clay ("devil").

Green sandy clay ("fire-earth"), with an irregular line of ironstone nodules ("red nob"), below which it is more sandy, and passes into the next.

Green sand.

Bottom-bed. Flint-pebbles, some very large, in sand, greenish towards the top, brown and light-coloured lower down; about 12 ft.

The junction with the Chalk was seen in a pit close by, and at a little distance to the west there was evidence of the junction of the Reading Beds with the Chalk in a swallow-hole formed by water falling as rain on the Reading Beds and flowing over or percolating through them to the edge of the Chalk, on meeting which it sinks through and dissolves it. This swallow-hole then held water, having evidently been artificially plugged in order to form a pond.

The outliers at Coleshill and Penn, between Amersham and High Wycombe, are the last in this series which we have visited (in May, 1898). At Coleshill the Reading Beds consist of mottled and dark-blue clay, and are capped by the basement-bed of the London Clay. This is not present at Penn.

The second line of outliers seems to merge into the first on the north-east, where the outliers are large and near together, each covering several square miles. Pointing south-westwards from the one south of Bennington and through the northern part of the Datchworth outlier, two small ones may be seen, one at Leverstock Green, which has twice been visited (in May, 1889, and May, 1900), and the other at Tyler's Hill or Cowcroft near Chesham, to which three visits have been paid (in May, 1878, April, 1897, and May, 1902).

In the Leverstock Green outlier the Reading Beds are seen in the Bennett's End brickfields, of which there are several. The most interesting geological feature here is the presence of two faults running nearly north and south, by which the London Clay and Reading Beds are let down against the Chalk which is seen on either side of them. The eastern fault, in a pit in the rear of

* *Mem. Geol. Survey*, "Geology of London," vol. i, p. 193.

Tilekiln Farm, has not been seen clearly for a long time, but the following section (Fig. 7) shows what was to be seen in 1861 :*

FIG. 7.—DIAGRAM-SECTION OF A PIT NEAR BENNETT'S END.—
W. Whitaker.

- LONDON CLAY: { *a.* Roughly laminated brown clay, from 0 to about 6 ft.
 b. Brown loam, over 6 feet.
 Basement { *c.* Flint-pebbles and oyster shells, about } about 12 feet.
 Bed. 6 inches.
 d. Brown loam, like *b*, $5\frac{1}{2}$ feet.

e. Bluish-grey plastic clay of the Reading Beds, holding water.

f. Chalk abutting against the basement-bed of the London Clay.

x x. Line of clay along the fault. (The upper right-hand part of the section, lightly engraved, not actually seen.)

The western fault, seen in 1900, was described by Mr. Upfield Green as then appearing as "a ridge of chalk, its eastern side sloping to an angle of $75-80^{\circ}$. The surface of the slope," he said, "is hard, polished, striated, and covered with a layer of black clay. Against this 'wall' of chalk rests, horizontally, a bluish-grey plastic clay, part of the Reading Series, surmounted by about 12 ft. of the basement-bed of the London Clay. . . ." It would appear that the Reading Beds have maintained their horizontal position against the western fault, but have bent downwards against the eastern fault to an angle of about 30° .

The basement-bed of the London Clay has here yielded many fossils, a list of which is given on pages 264 and 265 of Mr. Whitaker's Memoir on the "Geology of London" (vol. 1).

Tyler's Hill, or Cowcroft, as it is sometimes called, is a mile and a-half to the east of Chesham, whence it is approached over a hill and across a valley in which lies Layhill Common. It is conspicuous for some distance by the dense wood which covers it, the presence of this wood being due to the existence of the outlier of which the hill is formed. A chalk-pit and brick-field expose a complete section from the London Clay to the Chalk.

The following is the description of this section as seen in

* Whitaker, "Geology of the London Basin," *Mem. Geol. Survey*, vol. iv (1872), p. 235.

1853 by Sir Joseph Prestwich, from his classical paper on "The Woolwich and Reading Series":*

		ft.
	Gravel, chiefly of flint-pebbles in clay, averages	4
LONDON CLAY.	{ Brown clay with a few nodular septaria . . .	10
	{ Layers of laminated grey and brown clay. . .	3
BASEMENT-BED	{ Layer of imperfect septaria full of fossils . . .	0½
OF THE LONDON	{ Light brown sandy clay . . .	2
CLAY, 3¼ ft.	{ Flint-pebbles in clay . . .	1
	{ Umber-coloured clay, in places slightly mottled	
	{ red and yellow . . .	2.
	{ Fine siliceous sand, in places very white . . .	3
	{ Light-coloured soft sandstone with an occasional pebble; variable. . .	1
	{ Light-coloured siliceous sands with a few seams of grey clay, the lower part coarser, yellow, and brown . . .	10
WOOLWICH	{ Laminated grey and yellow clay and sand, with an under-seam of pebbles . . .	1
AND READING	{ Yellow and ash-coloured sand with seams of grey clay . . .	8
SERIES, 31 ft.	{ Grey clay laminated with sand . . .	4
	{ Large unrolled flints, apparently white-coated . . .	2
		51½
CHALK.		25

In septaria in the basement-bed of the London Clay *Ditrupa plana* abounds, together with *Ostrea bellovacina*, a few *Natica glaucinoides*, a *Fusus*, and teeth of *Lamna*. (Prestwich.)

On our second visit sections of the London Clay and Reading Beds were examined on opposite sides of the brick-field—in the north-east and south-west corners. In the first section the whole of the Reading Beds, here about 25 ft. thick, were seen, the lower portion consisting of grey, mottled, laminated clay, with a layer of about 6 ins. of green-coated un-waterworn flints and pebbles at the base, reposing on a level surface of chalk; the upper portion of very white sand, indurated in parts; capped by the basement-bed of the London Clay, 3 to 5 ft. thick. In the opposite section about 15 ft. of the Reading Beds were seen, the lower portion not being exposed. They were found to consist of 7 to 8 ft. of buff-coloured sands with irregular pipe-clay partings, passing upwards into 4 ft. of bright-red mottled sands with 4 ft. of white sand above. On this was the basement-bed of the London Clay, consisting of a well-marked layer of flint-pebbles about 6 ins. thick, in which sharks' teeth were found, followed by about 12 ft. of evenly-stratified laminated sandy clay, with a more or less continuous layer of flint-pebbles, from which layer many large calcareous blocks containing fossils had been taken.

In some of these blocks the following fossils were found: *Ditrupa plana*, *Cardium laytoni*, *Cytherea* sp., *Glycimeris retupiensis*, *Nucula* sp. (?), *Panopæa intermedia*, *Pectunculus*

* *Quart. Journ. Geol. Soc.*, vol. x (1854), p. 90.

brevirostris, *Fusus ficulneus*, *Melania* sp., *Natica* sp., *Pyrula* sp. On our last visit, in the first section a layer of detached flint-pebbles, some of large size, was seen about 6 ft. above the base, and in the second section the white sand of the Reading Beds was exposed.

Of the third line of outliers of the Reading Beds, two were seen in June, 1907—one being on Ivinghoe Common, west of Ringshall, and the other on Aldbury Common. Doubts, however, have been expressed as to the beds being *in situ*. Undoubtedly at both places they are Reading Beds—on Ivinghoe Common, 700 ft. above mean sea-level, sand only, and on Aldbury Common, at a height of 650 ft., both sand and clay. There they are seen in a brickfield, worked chiefly for brick-earth. Along the top of the pit is a thick, irregular deposit of clay full of angular pieces of flint with flint-pebbles and a few very small ones of quartz. A hole recently dug in one corner had resulted in the discovery of a mass of clay 12 to 14 ft. thick, brilliantly coloured in various shades of purple, red, and green. Near the top was an oval-shaped mass of fine, pale sand, suggestive of its having been thrust into the clay while in a frozen condition. Large flints were found in the lower part of the clay, and extending through the mass in two places were irregularly-contorted layers of flint-pebbles, having the appearance of Tertiary pebble-beds squeezed by lateral pressure. The clay appeared to consist of one thick mass doubled up, or of two masses pressed against each other, the beds of one part seeming to be horizontal and of the other nearly vertical. Much sand was seen amongst the clay. It was considered that the whole mass was undoubtedly of Reading age, its position and features being in all probability due to glacial conditions. Is it not, however, more likely that the Reading Beds were originally in position here, and have subsequently been disturbed by glacial action?

There is not sufficient interest attaching to the very few occasions on which we have examined sections of the *London Clay* alone to warrant any account of them being given here. In the early days of the Association cuttings for new lines of railway or for the widening of old lines out of London were visited, and many fossils were collected. A few found by the writer on our excursions to Finchley in 1865 and to Highgate in 1866 are in the Hertfordshire County Museum.

Overlying the London Clay on the hills of Harrow, Hampstead, and Highgate are the *Bagshot Sands*—distant outliers of the widely-spread sands that give such a distinctive feature to the tract over which they occur south of the Thames, a feature seen on a small scale on Hampstead Heath. Only the lower portion of the sands occurs in these outliers, and it is only at Hampstead that we have examined them, the outliers of Harrow and Highgate not having been visited. The junction of the London Clay

FIG. 8.—SECTION FROM PINNER TO HAMPSTEAD.—J. Logan Lobley.

with the Bagshot Sands has been seen at Hampstead on seven occasions—in 1870, 1872, 1874, 1877, 1887, 1895, and 1905.

Of the London Clay the only features recorded in our reports are the presence of bands of septaria and the fact of the upper and more sandy portion of the clay being more fossiliferous than the lower, which is mostly a stiff clay of a bluish tinge on a freshly-cut surface. It contains selenite and iron-pyrites, etc. Of its abundant fauna the only named *species* which we appear to have recorded, in addition to those from its basement-bed already mentioned (pp. 22, 26, 28, and 29), are the Mollusca *Corbula globosa*, *Modiola elegans*, and *Nautilus imperialis* from the middle zone at Ealing, and *Pectunculus decussatus* and *Voluta nodosa* from the top sandy clay at Hampstead. To these may be added numerous *genera* recorded from Ealing and Elstree.*

At one of our earlier excursions to Hampstead (27th April, 1872) it was seen that the water which percolated through the porous Bagshot Sand which capped the hill, being upheld by the impervious London Clay below, appeared as springs on the side of the hill. The "Pectunculus bed" in sandy clay, just below the Bagshot Sand (see Fig. 8), was found to be about 50 ft. thick. Drains

* *Proc. Geol. Assoc.*, vol. x. p. 181
vol. xiii, p. 387.

in course of construction in Greenhill Road exposed a good section of the sand, and the line of junction of sand and clay was followed as nearly as possible, being indicated by pools of water in Gayton Road, by the chalybeate spring in Well Walk, and by springs near the pond in the Vale of Health. In a large sand-pit on the top of Hampstead Heath attention was particularly directed to the gullies or small ravines which had been excavated by the action of rain falling on the surface and percolating through the porous deposit until checked by the thin bands of sandy clay which are interstratified with the sand. These ravines furnish an illustration of the process by which the diverging valleys around Hampstead may have been formed. From the Mount near North End the geological character of the country to the north was explained, the high range of hills by Bushey and Elstree forming the northern escarpment of the London Clay, and the intervening range of hills by Hendon and Finchley being capped by Glacial clays and gravels resting on an eroded surface of the London Clay. On the West Heath the position of the junction of the Bagshot Sand and the London Clay was again seen, and the spot by the Leg of Mutton Pond, where *Pectunculus decussatus*, etc., had been obtained during drainage operations, was pointed out.

At our last visit to Hampstead, at Kenwood Farm, near the "Spaniards," by the courtesy of the Express Dairy Company, a fine cliff of Bagshot Sand, showing cross-bedding and seams of clay, was examined, and the Director, Mr. G. Humphreys, contributed some notes on the geology of Hampstead, of which the following is a slightly condensed version:

"The sandy hill of Hampstead with its radiating valleys has been formed by circumdenudation, the sterility of the soil accounting for its having remained so long unenclosed prior to its purchase as a public park. The area of the Heath is 550 acres, the highest point being about 450 ft. above Ordnance Datum. The London Clay, on which the Bagshot Sands rest, is a marine deposit, blue in colour, but rusting on exposure to the air. It contains embedded masses, such as septaria, crystals of selenite, iron-pyrites, and in certain zones many fossils.

"This section of Bagshot Sand is in an old sand-pit. About midway up the cliff are some thin clayey seams interstratified with the sands. Ferruginous bands of indurated sandstone also occur, and on digging at the base we again encounter the 'pan.' The percolation of water through this highly ferruginous sand is the cause of the chalybeate wells at the junction with the impervious London Clay. The chalybeate spring in Well Walk attracted much attention in the past, and during the first forty years of the eighteenth century Hampstead became a flourishing spa, but ultimately fell into disrepute, morals and water both becoming contaminated. Mr. Allen Dick's analysis of Bag-

shot Sand from the Hampstead outlier is as follows: Quartz, 75 per cent.; felspar, 20; grains attracted by magnet, 2; clay, 1; zircons (in recognisable crystals), $\frac{1}{2}$; rutiles, 1-5; tourmalines, 1-10. The grains are sub-angular, giving evidence of wearing action on a sea-shore. A fragment of a fossil shell was stated to have been found in a ferruginous concretion, but no fossils are authoritatively recorded from the Hampstead Bagshots.

"The greatest thickness of Bagshot Sand under Hampstead Heath is 80 ft. This overlies 400 ft. of London Clay. Enormous quantities of sand have been removed from the heath."

The Bagshot Sands are the most recent regularly-stratified deposits in the area under consideration. A very long interval followed their deposition, during which the rest of the Eocene strata, and the Oligocene, Miocene, and Pliocene were deposited elsewhere. Here this was probably a period of extensive denudation, during which most of the great sheet of Bagshot Sand was swept away, the outliers of the hills of Harrow, Hampstead, and Highgate alone being left as witnesses of its former extent.

POST-PLIOCENE.

We cannot with certainty take the superficial deposits, most of which may be included under the comprehensive term "Drift," in strict stratigraphical succession, but the sequence followed will be: Clay-with-flints, the Plateau Gravel and Westleton Shingle, Glacial gravel and sand, Boulder-clay, River-drift and Alluvium (lacustrine as well as fluvatile).

Nor will space allow us to treat these deposits so fully as the older strata, or to refer to every notice of each deposit seen. It must suffice to state that every locality mentioned has been visited by the Association.

Clay-with-flints occurs only on the surface of the Chalk, of which it covers a considerable area in Hertfordshire. It is a stiff brown or reddish clay with large unworn flints. It has been formed in much the same way as the pipes in the Chalk—by dissolution of the carbonate of lime by carbonated water, the alluminous matter being left as a matrix for the indestructible flints. Perhaps also some waste of the Tertiary clays sometimes enters into its composition. Its formation probably dates from the time when the Chalk was first subjected to sub-aërial action, and is still going on. Associated with it is brick-earth, which appears, where directly lying upon clay-with-flints, to be a reconstructed deposit of Tertiary clays and sands.

The clay-with-flints varies much in thickness, from several feet when lying in hollows in the Chalk to only a few inches when on a raised surface. The brick-earth is a much thicker deposit. We have seen it in the brickfields near Leverstock

Green as a variously-coloured sandy clay from 20 to upwards of 50 ft. in thickness; and at Tyler's Hill, in the south-east corner of the brickfield, a section shows the Chalk fretted away in pinnacles which are invested by a layer of unworn flints in clay mingled with pebbles and broken shells (clay-with-flints), surmounted by brick-earth which thickens towards the south, extending downwards far below the level of the base of the pinnacles. In the upper portion of the brick-earth masses of conglomerate and irregular layers of flint- and quartz-pebbles are found, the latter parallel to the top of the Chalk, though not resting upon it.

The Plateau Gravel, which Professor Hughes designated "gravel of the higher plain," is intermediate in age between the Tertiary and Glacial periods, and that which he called the "gravel of the lower plain" is mostly of Glacial age. The gravels which Sir Joseph Prestwich has described as Westleton Shingle, Southern Drift, etc., are gravels of the higher plain.

Westleton Shingle occurs on Tertiary strata, usually on high ground, and as a rule at a greater elevation on the west than on the east. At Bayford we have seen it at 300 ft. O.D.; at Ayot and on Bernard's Heath, St. Albans, at about 420 ft.; on Tyler's Hill and Penn Common at about 550 ft. At Bayford it has flint- and quartz-pebbles in plenty, with radiolarian chert, jasper, schorl rock, and other foreign pebbles. On the Tertiary outlier of Ayot we have seen in Griggs Wood a more typical exposure on the surface of rather higher ground than in the brickfield, where it underlies boulder-clay.

At Bishop's Stortford, at about 220 ft. O.D., capping low hills on both sides of the Stort Valley, a plateau gravel has been seen which is not nearly so simple in its constituents as typical Westleton Shingle. This and other somewhat similar gravels in that neighbourhood are fully described by Dr. A. Irving in his report of the excursion of 17th July, 1897.*

The shingle-gravel on Hampstead Heath (450 ft.) and on Stanmore Common (500 ft.) is of pre-Glacial age. Prestwich thought that it bore more analogy to his Southern Drift than to the Westleton Shingle. It may be a reconstructed Bagshot pebble-bed.

Plateau-gravel, with sand, has been seen on the London Clay at Gerrard's Cross, in the long railway-cutting through the high ground to the west of the River Misbourne, its thickness varying from 5 ft. at the western end to 25 ft. at the eastern end.

Glacial gravel and sand occur almost all over our district. Amongst the localities at or in the vicinity of which they have been more particularly noticed may be mentioned, in Hertfordshire, Bishop's Stortford, Hertford, Hatfield, Datchworth, Welwyn, and Watford; in Middlesex, Whetstone, Finchley, and Hendon; and in Buckinghamshire, Loudwater.

* *Proc. Geol. Assoc.*, vol. xv, p. 193.

The Datchworth Tertiary outlier extends over Harmer Green and Burnham Green, and much of the greater part of it is

capped by Glacial gravel and sand, which has tended to its preservation. We examined the Glacial drift under Dr. A. E. Salter's guidance in 1905. It lies at about 400 ft. above mean sea-level. In a pit near Harmer Green a stratified gravel with false-bedded sand was seen. The gravel is made up of Tertiary flint-pebbles, sub-angular flints, quartz-pebbles, rough quartz-blocks (up to 4 ins. long), light-coloured quartzites, banded rhyolite, jasper, grits, dark chert, Herts conglomerate, etc. The non-local material is chiefly of a Palæozoic character. In the larger pit north of Datchworth Church sandy and clayey patches are frequent in the gravel, which shows lines of stratification well, except where it is disturbed owing to piping in the Chalk below. It consists chiefly of flints (some very large), Bunter quartzites, quartz, Herts conglomerate, and sandstone, and differs from the gravel at Harmer Green in containing much Triassic material. At Burnham Green boulders of Sarsen stone and Herts conglomerate were noticed.

At and around Watford the Chalk is covered by Glacial gravel. It is very irregular in thickness, as was well shown in 1872 when the cutting on the

FIG. 9.—DIAGRAM-SECTION THROUGH WATFORD AND BUSHEY.

Horizontal scale : 1 inch = 1 mile. Vertical scale exaggerated.

main line of railway was being made towards the new tunnel. The gravel here consists of sub-angular flints derived from the Chalk, of rounded flint-pebbles also originally derived from the same formation but owing their present form to the attrition they have undergone in the older Tertiary rivers and seas, and of pebbles of very ancient rocks. Here and there also there are in the gravel more or less regularly stratified beds of sand divided by layers of pebbles.

A few years later, in 1879, Glacial gravel was again examined at Watford, first in the Colney Butts gravel-pits and then in the Hagden Lane gravel-pits. At neither place is there any sign of bedding. Numerous drifted fossils have been found in these pits, some derived from the Chalk, others from Jurassic rocks. On the opposite (south-east) side of the Colne the gravel is, as shown in the section (Fig. 9), replaced by brick-earth.

In the excursion to Penn and Coleshill, Glacial drift was seen on the slopes of the hills on both sides of the Wycombe stream at Loudwater, at about 350 ft. above O.D. On the south-west it consists of clayey gravel made up of sub-angular flints, flint-pebbles from Eocene pebble-beds, and a few boulders of quartz and quartzite, one quartz boulder measuring 2 ft. by 14 ins. by 12 ins. On the north-east it is stratified gravel and sand, the gravel being mostly weather-worn material of all sizes, with large pebbles of quartz and quartzite, and a few pebbles of black chert, and the sand having thin seams of clay in places.

There may be some doubt about the age of drift gravels where not associated with boulder-clay, but where, as between Hatfield and Ayot Green, *boulder-clay* occurs in and upon the gravel there can be no doubt as to its age being Middle Glacial. The large gravel-pit by the side of the Great Northern Railway near Woodhall Lodge Farm for long exposed a very instructive section, but the pit is now being filled in. The section was very clear on our visit in May, 1898, when it was as follows:

		ft
MIDDLE GLACIAL.	Boulder-clay, containing much chalk in its lower part, and <i>débris</i> from the Midlands, <i>e.g.</i> , various fossils and Triassic and other pebbles.	12
	Sands, with patches of gravel containing similar <i>débris</i> about	15
	Boulder-clay similar in character to the higher bed	1
	Ochreous flint-gravel and sands with Triassic quartzites about	15
CHALK, disintegrated.		

Sometimes, as at Bishop's Stortford, boulder-clay and gravel are much mixed up, or occur close together on the same level. There, within such a small area as the churchyard of Hockerill

Parish, we find in the graves such varieties as an almost pure (reconstructed) chalk with angular black flints, a stiff tenacious clay, a fine sandy gravel with small pebbles and chalk detritus, and an iron sandy gravel with no chalk at all. Jurassic fossils derived from the Mercian Jurassic rocks are of quite common occurrence.

The boulder-clay of Bricket Wood, between St. Albans and Watford, has not been visited by the Association; but the next and most southern patch, that of Finchley, has frequently been seen with the underlying Glacial gravel in excursions to Whetstone, Finchley, and Hendon. The boulder-clay here covers a triangular area of about three square miles, the gravel beneath it enlarging the area of the Glacial deposits to about five square miles. The apex of the triangle is at Whetstone, the base extending from Hendon nearly to Muswell Hill; Finchley is about in the centre. The most complete account of the deposits is in the report of the fourth visit (1892), by the Director, Dr. Henry Hicks. Almost the whole of this report is here reproduced, with two of the illustrations which accompanied it.

From the terrace of Hendon Grove, which commands an extensive view over the undulating plain of north-west Middlesex to Stanmore Common and the Hertfordshire Hills, Dr. Hicks pointed out the physical features of the district, which, he said, in the main dated back to a very early Glacial, if not to a pre-Glacial period. The hills were more or less mantled by glacial deposits, which have either been removed or rearranged along the lower horizons and in the valleys, but occur undisturbed on nearly all the heights and along the slopes of the hills. At the base, resting on an uneven floor of London Clay, there is usually a rough pebbly gravel, mainly of flint-pebbles, but containing also large and but slightly-worn flints, masses of Sarsen stones, and a few pebbles of quartz and quartzite. At Hendon and Finchley angular and sub-angular fragments of chert and ragstone are also found in it in considerable abundance. The pebbly gravel which occurs on Stanmore Common, Bushey Heath, and at Elstree, etc., at heights of over 400 ft., resembles this lower gravel at Hendon in a remarkable degree, except that it does not often contain chert and ragstone. On Bushey Heath it is overlain by the so-called quartz gravel of the Upper Plain.

In a field at Hendon Grove a portion of what had been an extensive section was specially exposed, and it was explained that at the south corner a mass of yellowish clay filled a channel in the gravel, passing through it to reach the London Clay below. In this clay several distinct patches of gravel were noticed, which Dr. Hicks said could hardly have been deposited except in a frozen condition. The gravel in these patches usually differed somewhat in character from the lowest pebbly gravel,

resembling the quartzose gravels which in many parts of the district overlay the pebbly gravel. The section is as under:

- a.* Surface soil.
- a'*. Gravelly soil (with flints). Irregular.
- b.* Yellowish-brown clay, in places passing downwards through the underlying sands and gravels. 2 to 7 ft.
- d.* Loamy sand with ochreous band. 1 ft.
- e.* Gravel. 1 ft.
- f.* Yellowish sand, sometimes showing current-bedding. 2 ft.
- g.* Ochreous gravel, with large pieces of Sarsen stone, rounded and sub-angular flints, chert, quartzite, quartz, ironstone, etc. 2 to 4 ft.
- h.* London Clay much disturbed, of a brown colour, and forming a very uneven floor under the glacial deposits.

The West View pit was next examined, and the section exposed (Fig. 10) was explained.

It was pointed out that here, as at Hendon Grove, the brown clay passes in places through the sand and gravel to reach the London Clay floor, and that it attains a thickness at those points of 16 or 17 ft. The gravel overlying the brown clay was shown to contain a larger proportion of angular, white-coated, and unworn flints than was found in the gravels below it. It was mentioned that the sections here and at Hendon Grove exhibited the conditions not only to be witnessed everywhere over the Hendon plateau, but also in the Finchley and Totteridge districts wherever sections showing the deposits below the chalky boulder-clay were to be seen. In the West View pit, as at some other points in Hendon, there were indications of a slight spread of chalky boulder-clay over the brown clay, but it was separated from the latter by a varying thickness of gravel.

In a section near Golder's Hill, on the Finchley side of the Brent Valley, a similar succession of beds to that found on the Hendon plateau was observed. The deposits reach down on that side of the Brent Valley to within a short distance of the level of the present stream, being here well below the 200 ft. contour-line. There is therefore a rise of over 100 ft. in the floor of London Clay from this spot to where it is found near the higher levels at Finchley under similar glacial deposits, the highest point there being over 300 ft. above Ordnance Datum.

A section in a field laid out for building purposes, between Finchley Road and the Manor House, Finchley, did not indicate, Dr. Hicks stated, the complete succession of the beds, which had been better exposed at some other points (see Fig. 11).

In nearly all the sections exposed since, a much greater thickness of brown clay occurs between *d* and *e*.

FIG. 10.—SECTION IN WEST VIEW GRAVEL-PIT, HENDON.—*H. Hicks.*

(About 20 feet in length.)

<i>a.</i> Surface soil.	<i>f.</i> Yellowish sand with seams of clay. 2 ft.
<i>a'.</i> Gravelly soil.	
<i>b.</i> Brown clay, with a few flints, patches of gravel, and in places much "race." 6 ft.	<i>g.</i> Gravel with angular masses of Sarsen stone, sub-angular flints and flint pebbles, chert, quartzite, etc. 2 to 3 ft.
<i>c.</i> Laminated sandy clay. 6 ins.	
<i>d.</i> Grey sandy gravel.	<i>h.</i> London Clay.

FIG. 11.—SECTION IN FIELD OPPOSITE THE QUEEN'S HEAD, FINCHLEY.
—*H. Hicks.*

(March 8th, 1892.)

<i>a.</i> Surface soil.	<i>d.</i> Sandy gravel, with thin seams of brown clay. 2 ft. 6 ins.
<i>b.</i> Chalky boulder-clay. 2 to 4 ft.	
<i>c.</i> Brown sandy clay, with sub-angular flints, etc. 1 ft.	<i>e.</i> Yellow sand. 2 ft. 6 ins.
	<i>f.</i> Rough pebbly gravel. 3 to 5 ft.
	<i>g.</i> London Clay.

The section in a deep well on the east side of the road was :

Soil.
Chalky boulder-clay. 3 ft. 6 ins.
Sub-angular gravel. 1 ft.
Sandy gravel. 5 ft.
Brown clay with much "race" and flints. 6 ft.
Loamy sand. 1 ft.
Pebbly gravel. 4 ft. 6 ins.
London Clay.

The brown clay which separates the pebbly gravel from the chalky boulder-clay is evidently on the same horizon as the brown clay with "race" at Hendon, and is undoubtedly an earlier deposit than the chalky boulder-clay in the above section. In addition to the patches of gravel which it frequently contains, scattered flints are found in it, but no rocks which need have travelled farther than from the Midlands, or the south of England, have yet been detected in it.

At an earlier visit to Whetstone and Finchley, in 1887, remarks were made by the Directors, Mr. H. B. Woodward and Mr. J. G. Goodchild, on the origin of the gravels and boulder-clay of the district. In a pit south of Whetstone, where boulder-clay was seen resting on Middle Glacial sands and gravels, Mr. Woodward pointed out that the clay here was part of the chalky boulder-clay which covers a large part of the Eastern Counties. It was composed of chalk, flints, and Jurassic clays. In his opinion it was a product of land-ice. The streaky structure which in places simulates stratification was due to pressure. As a rule the beds beneath the boulder-clay in the Eastern Counties were more or less contorted, and in this pit there was evidence of some disturbance in the underlying beds.

In a pit in Somers Lane, where the clay is being worked for bricks, a continuous face of it, about 30 yds. long and 6 ft. high, was exposed. In the middle of the section the upper 4 ft. consisted of ordinary boulder-clay; beneath it, for a length of 20 yds. or more, were bands of striped loams, loamy clays, and sands, alternating with and passing horizontally into boulder-clay like that above. In the boulder-clay were found red chalk, carstone, Jurassic rocks, New Red Sandstone and breccias, rocks of Carboniferous age, rocks from Charnwood Forest, granitic and metamorphic rocks, and basic rocks, some similar to those of the Whin Sill of the north of England, and others regarded as of Scandinavian origin. Here Mr. Goodchild said that he believed the Finchley boulder-clay marked very nearly the southern limit of the ice-sheet, the rate of melting of the ice-fringe hereabouts having exactly balanced the supply. He considered it as approximately of the same age as that originating when Shap granite, etc., was transported inland and uphill over Stainmoor, that is as contemporaneous with the period of maximum glaciation, when Scandinavian land-ice on

one side and the North Atlantic ice on the other reinforced our glaciers and hemmed in our local ice, so that it was compelled to flow southwards beyond the limit it would otherwise have attained. He regarded it, and all the various stratified beds associated with it, as the sediment of the mud and stones which had worked up into the ice from various parts of the areas its feeders had traversed, and which had been liberated by the melting of the ice under sub-aërial conditions immediately after the climax of the Glacial Period.

Amongst the drifted fossils observed were *Gryphaa dilatata*, *Belemnites abbreviatus*, *Lima gigantea*, and *Discina latissima*.

At our last visit to this district a discussion occurred on the origin of the Finchley Drifts. It is fully reported in our PROCEEDINGS (vol. xix, p. 314).

We cannot always differentiate between *river-drift* (Pleistocene), by which is meant the sediment of rivers flowing at a higher level than our existing streams, and *river-alluvium* (Holocene), which is the sediment of our present rivers, for they pass insensibly into each other, and the only lake sediment we have to notice which comes under the term alluvium is of Pleistocene age. When terraces of gravel occur, the highest terrace is the oldest, for all our rivers, whilst depositing gravel, have been cutting down their beds. On the other hand, when rivers have attained their base-level and are depositing mud, they are usually silting up their beds; then the lower stratum is the older, and it is the same with lacustrine alluvium. It will therefore be most convenient to take these deposits in topographical order, commencing with the Hitchin alluvial deposits.

The Hitchin fresh-water bed occurs between brick-earth from 10 to 30 ft. in thickness, and, so far as known except in one place, gravel and sand presumed to be of Glacial age. The exception is where beneath it has been found chalky boulder-clay with a base of loamy chalky gravel reposing on gravelly sand. When examined by the Association in Mr. Jeeves' brickfield in June, 1896, the following beds (in descending order) were exposed:

1. Brick-earth of a reddish colour with a few stones scattered through it, and containing numerous flint-implements.
2. The freshwater bed, consisting of soft calcareous loam of a light colour, from brown to almost white, passing down in some parts of the brickfield into a dark grey or almost black deposit; fairly evenly stratified.
3. Gravel composed of flints, chalk, pebbles, and a great variety of rocks all more or less rolled and water-worn; well stratified and often current-bedded.

Mr. H. W. Monckton observed that in the Geological Survey Map, 46 N.E., Drift edition, eight divisions of the Drift were

given, two of which, "clay-with-flints" and the "loam" associated with it, did not appear near the present locality. The other divisions were, beginning with the newest: 1, Alluvium; 2, Loam; 3, River-gravel; 4, Glacial loam; 5, Glacial boulder-clay; 6, Glacial gravel. According to the map, the brick-earth was classed with the loam of Glacial age (No. 4), but he thought it was an open question whether it and the fresh-water bed ought not to be classed with the "Loam" No. 2 instead of with the Glacial loam. The celebrated fresh-water deposit at Hoxne, he said, bears some resemblance to the Hitchin bed: at one time classed with the Glacial Series it is now considered by Messrs. Whitaker and Dalton to be post-Glacial. Subsequent researches by Mr. Clement Reid have borne out Mr. Monckton's views.

It would be useless to repeat the list of fossils found in this bed given in our PROCEEDINGS, as it has since been greatly augmented. The fullest lists are given in the *Transactions of the Hertfordshire Natural History Society*—of the fossils generally by Mr. Clement Reid (vol. x, pp. 18, 19; vol. xi, p. 64), and of the Ostracoda by Mr. Frederick Chapman (vol. xi, pp. 61-62). The following groups are represented: Characeæ, Mosses, Flowering plants, Polyzoa, Entomostraca, Mollusca, Insecta, and Mammalia. The more recent finds include statoblasts of the fresh-water polyzoon *Cristatella mucedo*, winter-eggs of the entomostracan *Daphnia*, and the teeth of a vole, probably *Microtus arvalis* (*loc. cit.*, vol. xi, p. 64). Up to April, 1901, twenty-nine species of flowering plants had been identified by Mr. Reid, all of existing British species, and including *Najas marina*, now only found in Britain in two of the Norfolk Broad. All the Mammalia are from the top of the bed, just beneath the brick-earth.

The Hitchin fresh-water bed probably represents, as at Hoxne, "a silted-up river-channel, which remained for a time in the state of a lake or mere." *

The district of most interest in connection with our examination of Thames valley-gravel is that extending from the Colne at West Drayton, through Dawley, Hanwell, and Ealing, to the Thames at Brentford. Excursions to this district have been made in 1887, 1895, and 1904.

At West Drayton sections of gravel (with flint implements), sand, and clay have been examined. At Dawley three sets of pits have been visited. In Mr. Odell's pits the implementiferous lower bed of stratified gravel was found to be surmounted by a thick, irregular deposit of sharp white sand, with unstratified gravel and clay above it, which filled a long, shallow depression, and extended on both sides of it. In Mr. Maynard's pits a very fine exposure of stratified drift about 5 or 6 ft. thick, with enclosures of sand and clay, and brick-earth above it, was seen to rest

* Lyell, "Antiquity of Man," Ed. 4, p. 268 (1873).

on well-defined stratified gravel. And in Mr. Pipkin's pit very remarkable wave-like forms of chalky matter, clay, and stones were seen overlying contorted sandy deposits and unstratified gravel, and underlying dense, undisturbed, brown brick-earth. Near Hanwell a section 20 ft. in depth has been seen, exposing fine sections of contorted and also unstratified deposits above horizontally-bedded drift. In the unstratified gravel were enclosures of bluish clay, sand, etc., and a huge furrow reaching from the top to the bottom of the excavation was exposed. This furrow is within six or eight feet of a similar deep, unfilled furrow, and the gravel in the space between them was seen to be contorted and squeezed into arch-like forms.

The report of an excursion on the 18th of June, 1887, to the Mount, Ealing, and Horsington Hill, occupies more than nine pages of our PROCEEDINGS (vol. x, pp. 172-181). A better idea of the principal results of the investigations of Mr. J. Allen Brown, the Director on that occasion, will perhaps be derived from the following extracts from his work, *Palaolithic Man in N.W. Middlesex* (1887), than would be gained from an abstract of that report:

"The gravel and brick-earths in this locality [Ealing] extend from the margin of the Thames, say, to the top of Castlebar Hill, a distance of about three miles. Castlebar Hill, the Mount, Hanger Hill, and other high ground extends as a ridge, dividing the Thames Valley, or alluvial, deposits from the wider extension of the valley to the north. . . . The ground dips gently to the river, but with three very distinct inclines; they are comparatively sharp descents, forming benches or terraces, marking the elevations at which the greater river of the Pleistocene period formerly flowed at different times; they are geological bench-marks showing that the ground has thus been successively eroded until the river has worn itself a channel more than 140 ft. below its first level. . . . Prestwich (in *The Ground Beneath Us*) . . . showed that these river-drift gravels are but re-deposits at lower levels of yet older marine gravel, called the Plateau gravel, which is now found covering the Chalk hills of Hertfordshire and Bucks, the latter being much higher than the London Clay hills, with their coating of gravel and alluvium, lying between the Chalk ranges of Herts and Surrey.

"Mr. Whitaker . . . has divided the river-drift deposits into three distinct terraces or benches up to the 100 ft. contour. . . . There is, first, a low terrace occupying the low ground in the salient bends of the present Thames at an average height of 10 to 20 ft. above the present stream; second, a mid terrace ranging from 20 to 30 ft. above the bottom of the valley; and, third, a high terrace gravel with its covering of brick-earth at a level of from 50 to 100 ft. above the Ordnance Datum; but,

as he believed, the series should be extended to a higher level.

"My investigations have shown that this high-terrace gravel reaches, in the vicinity of Ealing, to a much greater height, since I have found it to occur not only on the shoulders of Castlebar Hill, say, to 130 ft. O.D., and continuous with the other high-terrace gravels; but . . . actually united with the thick beds of gravel along the summit of the hill itself. . . .

"The height of these deposits is from 150 to 160 ft., the highest part of the hill being 167 ft. O.D. But there is a yet higher deposit of gravel to be described, which however tells of another condition of the region we are noticing, and which is probably much older than those already mentioned. It is met with on the tops of the highest hills of the ridge, *i.e.*, on the Mount, Hanger Hill, as well as on Horsington Hill on the north. On the Mount, . . . 204 ft. O.D., and nearly 40 ft. above Castlebar Hill, I found the gravel was associated with blocks and boulders of rocks entirely foreign, or not found *in situ* in the Thames Valley. . . . The whole presents us with all the elements and conditions of a glacial formation." (pp. 43-46.)

"Professor Prestwich had shown us that the high level and lower level gravels are but extremes of a series marking a long period of time, and probably formed under analogous but not identical conditions. The high-level gravels are formed in the estuaries dating after the boulder-clay deposits and the great extension of ice. The high-terrace gravels throughout the Thames Valley have yielded the flint-implements, worked flakes, and waste flakes of Palæolithic man. I have found them at Dawley, . . . Hanwell, Ealing, Acton, Brentford, Turnham Green, etc." (*Loc. cit.*, pp. 30, 31.)

The section (Fig. 12) represents contorted ice-drift at the Mount, Ealing. Long, tongue-like masses of sand and gravel are shown, the gravel enclosing masses of sand and deposits of purple clay, the whole extending like irregular furrows into the midst of the stratified beds. The pressure exercised by masses of ice, pushing before them, contorting, and churning up the laminated beds into these remarkable *flexures*, seems, Mr. Brown said, to be the only means which could have brought about such results.

In 1875 excavations on the site of the Natural History Museum at South Kensington were examined, and the Thames Valley gravels were seen to have, within the area, a maximum thickness of about 20 ft. towards the eastern end, whilst on the west they were only 7 ft. thick. A tooth of an elephant, *Elephas primigenius*, had been found, resting almost directly on the London Clay.

In 1906 the Hertfordshire Natural History Society visited Long Valley Wood, near Croxley Green, Rickmansworth, and

the extensive sections of fluvial gravel there exposed were considered to be of sufficient interest for examination by members of the Geologists' Association. Accordingly, in the following year, after sections of the Chalk and Tertiaries had been seen at Bushey, these Drift deposits were examined.

The gravel here is or has been worked for a distance of about half-a-mile in a north-easterly direction. Layers of sand and clay are interstratified with it, and the total thickness of the alluvial deposits is from 20 to 30 ft. They rest on the Chalk in a very uneven manner, so that it is difficult to determine the height of their base, but it appears to average about 30 to 40 ft. above the level of the existing River Gade, on the right bank of which they are situated. The Gade is a tributary of the Colne, which is from a quarter to half-a-mile distant, and this gravel should perhaps be considered as part of the alluvial deposit of the Colne in Pleistocene times rather than that of the Gade.

The layers of sand and clay and some peculiar black bands give to the gravel in many places a stratified appearance, but nowhere horizontal for any considerable distance, this conformation being due in

FIG. 12.—SECTION AT THE MOUNT, EALING. (About 50 feet long).—J. Allen Brown.

* Included mass of sand. ** Included deposits of purple clay.

places to cross-bedding, but mostly to the irregular dissolution of the underlying Chalk.

The gravel consists for the most part of water-worn flints, some completely rounded, others sub-angular, and also contains quartz and quartzose-pebbles. Its most interesting feature is, however, the large number of Palæolithic implements which have been found in it, most numerous near its base. A molar of an elephant, most probably *Elephas antiquus*, has been found at a depth of 20 ft. from the surface, and also a tusk, believed to be of the same species, but this crumbled away when touched.*

In 1906 a mass of sand and clay, with Sarsen stones, was seen in the gravel, and then determined to belong to the Reading Beds. Its appearance at that time is shown in Plate II, Fig. 1. It had been preserved intact, so far as possible, for the present visit, but the large Sarsens which were on the top had fallen down, and the clay above the sand had shifted from the horizontal position in which it was first seen. One of these Sarsens is represented in Plate II, Fig. 2. It is about 5 ft. long and 2 ft. 6 ins. wide.

The preservation, in gravel formed by a river, of soft mottled clay and loose white and yellow sand apparently in the position in which they were originally deposited is difficult to account for. The mass when first seen was about 6 ft. in height and 10 in width, and the gravel was distinctly arched above it. The sand was stated by the foreman of the gravel-pits to go down at least 6 ft. below the present level of the surface, which is here close to the Chalk. This seemed to indicate a "pipe" in the Chalk. The Sarsens were found to be soft in places; one was sharply split across, another was deeply mamillated.

A recent visit by the writer (1908) showed that "piping" was the cause of the preservation of this patch of Reading Beds, and of the curvature of the stratum of gravel above it. It had been in a "pipe," but the chalk around it has been dissolved away by carbonated water, which has not affected the Tertiary material. The gravel above that has therefore remained in the position in which it was deposited, but has sunk around it where resting on the Chalk, thus assuming an arched form. This explanation implies an enormous lapse of time since the deposition of the Palæolithic gravel—a sufficiently long period for a depth of 6 ft. of chalk to be removed by chemical dissolution.

Between Harefield and Colnbrook the River Colne runs over a deposit of gravel probably from 10 to 15 ft. in thickness, and lying upon it there is usually a layer of peat or peaty soil, with occasional patches of shelly marl. In 1903† this deposit was examined near Denham, and the following species of Mollusca obtained from the marl were identified by Mr. A. S. Kennard: *Vitrea crystallina*, *nitida*, and *radiatula*, *Arion ater*, *Vallonia*

* Sir John Evans, in *Trans. Herts. Nat. Hist. Soc.*, vol. xiii, p. 66.

† *Proc. Geol. Assoc.*, vol. xviii (1903), p. 188.

[Photo by J. Hopkinson.]

1.—MASS OF READING BEDS IN PALÆOLITHIC GRAVEL AT LONG VALLEY WOOD, NEAR RICKMANSWORTH.

[Photo by J. Hopkinson.]

2.—SARSEN STONE FOUND IN READING BEDS AT LONG VALLEY WOOD, NEAR RICKMANSWORTH.

pulchella, *Acanthinula aculeata*, *Pyramidula rotundata*, *Hygromia granulata* and *hispidula*, *Helix aspersa*, *nemoralis*, and *arbus-torum*, *Clausilia bidentata* and *laminata*, *Cochlicopa lubrica*, *Pupa muscorum*, *Succinea elegans*, *Ancylus fluviatilis*, *Velletia lacustris*, *Bithynia tentaculata*, *Valvata piscinalis* and *cristata*, *Limnaea stagnalis*, *pereger*, *palustris*, and *truncatula*, *Planorbis albus*, *spirorbis*, and *contortus*, and a *Pisidia*. Mr. Kennard considered the most noteworthy species to be *Helix aspersa* and *Acanthinula aculeata*, the former having for long been believed to be a Roman introduction into these islands, and the latter being rather rare in deposits of any age. Flint-flakes and old cores were found scattered over the recently-exposed gravel, and Mr. Kennard said that the majority of the worked flints were obtained direct from the Chalk, the nearest locality being about two miles away. The implements are nearly all well-known cave types. Their exact age is doubtful. They must be newer than the third terrace, on which they rest, and older than the alluvium, but whether they belong to the last stage of the Pleistocene or to the early part of the Holocene is doubtful, though, judging from the character of the implements, they appear to belong to the former.

The gravels of the Brent Valley were examined in 1887, when the Director, Mr. J. Logan Lobley, said that they formed part of the alluvial deposits brought down by the Brent in the past, and were not confined to the lowest level of the valley, being also seen on the level of Kingsbury Church, at an elevation of 60 ft. above the stream, giving a low-level and a high-level gravel. In this respect the Brent Valley is comparable with the main Thames Valley, which displays gravels at different heights. The Thames gravels consist of sub-angular flint fragments derived from the erosion of chalk lands, while the Brent gravels consist of whole or broken rounded pebbles of flint, with some of quartz and fragments of other rocks.

The Drifts of the Valley of the Lea have been examined in the vicinity of Tottenham and Homerton. On one of our early excursions (7th August, 1869) some excavations made in this valley near Walthamstow for reservoirs of the East London Waterworks were inspected. Prof. John Morris then pointed out that one of the most important features to observe was that the beds occurred irregularly, or more or less in basins, varying much in thickness. The gravel forming the bottom of the reservoir was an ordinary valley-gravel overlying London Clay, and contained remains of the elephant and of *Bos primigenius*; on this was a shelly marl covered by a bed of very black peat, this deposit containing a large number of land-shells, drifted and fallen trees and roots of trees, and many hazel nuts. This stratum is covered by a grey clay, which frequently occurs in thin beds overlain by a dark or brown-coloured

clay, in which can be observed the stems and roots of plants undergoing decomposition and encased by a kind of bog-iron-ore which also occurred in irregular masses.

Professor Morris said that the conditions here seen might in a great degree be explained by facts within the range of human history. In the first and even middle periods of the eighteenth century the tract between the Roding and the Lea and the adjacent country was forest-land to a large extent, with plenty of game and herds of deer. And just as we find in the lower beds—the old valley-gravels—the elephant and *Bos primigenius*, so in the higher beds would be found the goat, horse, and ox. There had been found three species of oxen, the red deer, the roebuck, the fallow and moose deer, the wild boar, the wolf, the fox, and the beaver. There had also been found the remains of man, and implements used by him, including a bronze spear and arrow-heads, a Celtic sword, part of the bronze sheath of the Celtic dagger, a bone knife, a pierced stag's horn axe-head, a stag's horn club, earthen pots, and an armlet of Kimeridge coal.

The general character of the deposits indicated the former and far greater extent of marsh and forest-land, as well as the course of the ancient river, its unequal depth, and the slow deposition of clayey, sandy, and shelly beds, many of the shells being fluviatile forms, such as *Unio*, *Planorbis*, *Limnæa*, *Cyclas*, and *Bithynia*, associated with marsh and land genera, such as *Succinea* and *Helix*, probably washed in during floods and deposited with the fresh-water species; while the numerous remains of animals entombed with them showed the abundance of those forms which roamed among the forests or frequented the marshes during the period in which the river rolled its onward course and formed the deposits now exposed. Nor was man absent from this area, as is fully testified by his implements.

Thirty-two years later, excavations were again being made in the neighbourhood of Tottenham for reservoirs of the East London Water Company, and we visited them in April, 1901. The works in Woodford Road were first seen. The most important find here is a fine example of a "dug-out" canoe made from the trunk of a large oak. Though usually considered to be of pre-Roman age, part of a Roman amphora was found at the same level, and only a few feet away. Other relics comprised a Saxon spear-head, two wooden implements, probably pile-drivers or mallets, two human skulls, an antler of red deer, bones of *Bos longifrons*, horse, and sheep, fragments of a Viking ship, and other objects of Roman, mediæval, and modern age.

The most interesting sections were seen at the intake of the works near Chingford Mills. The sections vary greatly, but the sequence generally is about 10 ft. of peaty alluvium resting on blackish-grey gravel, and that again on red gravel. In places

the alluvium was seen to contain lenticular patches of shell-marl, and in one place similar shells were found in the blackish-grey gravel, the species proving its Holocene age. The red gravel is probably of Pleistocene age.

Collections were made from the shell-marl, and bones of horse, sheep, red deer, pig, *Bos longifrons*, and dog or wolf were also obtained. In former excavations the remains of elk, bison, *Bos primigenius*, fox, goat, beaver, roe deer, fallow deer, reindeer, and great Irish deer have been found, besides a single bone of the sea-eagle. From these beds fifty-two species of Mollusca have been collected, the most interesting of which is *Acanthinula lamellata*, a species not known living south of Staffordshire.

The last locality to be mentioned in which river-gravels and alluvial deposits have been examined is Homerton and its neighbourhood, lower down the Lea Valley, visited in 1883.

The river-terrace here is 33 ft. above the level of the Lea, and the opposite terrace on the eastern side of the Lea in Essex is exactly a mile and a half away. The Director, Mr. Worthington G. Smith, explained that other terraces were more than 90 ft. higher than the Lea and correspondingly wider apart, showing the great change which has taken place since Palæolithic times in the excavation of this wide and deep valley. Very few implements were found in the valley lower than the 33 ft. terrace, proving that there was a break in the continuity of the Palæolithic age at the time of the formation of this terrace. The few implements found in it were as a rule abraded and ochreous. The gravel and sand seen in the open excavations were of various depths, from 12 to 30 ft., and rested on London Clay. Mr. Smith then called attention to the "warp and trail," ice-deposited material, which he said covered nearly all the north-east of London. It does not, however, go under the Lea, gradually thinning off farther down in the valley, and by its deposition it seems to indicate that it followed a natural depression of an old surface along which the Lea afterwards ran. All the implements are older than the deposition of the "warp and trail," and enormously older than the Lea as known in Neolithic and modern times. As a proof that the valley has not materially changed since Neolithic times he instanced the finding of Neolithic celts and other tools just beneath the surface *humus* at the bottom of the present valley.

The brick-earth, rich in Mammalian remains, was seen in a brick-pit at Upper Clapton, 44 ft. above the Lea. Many fine examples from here are now in the Natural History Museum at South Kensington.

Stoke Newington Common was then visited. Here Mr. Smith has lighted on an undoubted Palæolithic settlement of great extent, the sandy floor or working-place remaining just as it was

left by the men who lived on the spot before (in his opinion) the thick, tenacious, ice-laid deposit, known as "warp and trail," was deposited. The section shows, in ascending order: Thames gravel containing slightly-abraded lustrous implements; fine sand often full of fresh-water shells, including *Corbicula fluminalis* and *Hydrobia marginata*; the "Palæolithic Floor," or working-place, on which rest hundreds of implements and flakes as sharp as on the day they were made, quartzite hammer-stones with ends abraded off, fossil bones, antlers, teeth, etc.; and on the floor sandy loam, loamy sand, and "warp and trail" where it has not been removed for brick-making. On the top is surface *humus*, containing Neolithic implements, bone tools, British and Roman pottery, Roman coins, and other objects of recent date.

We have now traced the history of Middlesex and Hertfordshire, so far as revealed in excursions of the Geologists' Association, from Palæozoic times to the advent of Man, whose early progress towards civilisation, as given by Lucretius more than half-a-century before the Christian era, has been confirmed by recent researches. In the following extracts from Creech's edition (1714) of *De Rerum Natura* (Book V.) the word he translated "brass" is (correctly) named bronze.

"They knew no use of Fire to dress their Food;
No Cloaths, but wander'd naked in the Wood;
They liv'd to shady Groves, and Caves confin'd,
Meer shelter from the Cold, the Heat, the Wind."

"Whilst Cruelty was not improv'd by Art,
And Rage not furnish'd yet with Sword nor Dart;
With Fists, or Boughs, or Stones the Warriours fought;
These were the only Weapons Nature taught:
But when Flames burnt the Trees, and scorch'd the Ground,
Then Bronze appear'd, and Iron fit to wound.
Bronze first was us'd, because the softer Ore,
And Earth's cold Veins contain'd a greater Store."