LARYNGEAL URTICARIA, REPORT OF A CASE.

BY C. E. COOPER, A.B., M.D., DENVER, COLO.

The infrequent occurrence of laryngeal urticaria forms the chief motive for this report. Unfortunately the actual lesion was not visible, but sufficient clinical and objective evidence did exist to justify the belief in the presence of a wheal acting as an obstruction to respiration and located somewhere below the epiglottis.

Miss B., age 20 years, single and a nurse by occupation. Family history good. Was operated upon several months ago for appendicitis. Made an excellent recovery with evident general benefit to her health. For two weeks has been complaining slightly with rheumatism (articular) and has been taking aspirin and salicin in moderate doses, at the same time continuing her work.

On the evening of July 11th, after a meal of meat, potatoes, bread and butter and raspberries she was suddenly seized with a marked dyspnoea, choking sensations and cyanosis. She sought relief from another nurse who summoned an interne. He administered morphine, atrophine, nitroglycerine and amyl nitrate in an endeavor to relieve the dyspnoea. About one half hour afterward I saw her. She was extremely nervous, respirations about sixty and difficult, color good and pulse good. It fell to my lot to see her, not because any throat lesion was suspected, but because I happened to be in the hospital at the time and the sister in charge asked me to do something for her intense itching. From head to foot she was one mass of wheals and was evidently suffering extremely. In an endeavor to explain the marked dyspnoea and cyanosis of but a short time before, I examined the heart and lungs and found both normal. I then examined the throat; the pharynx as far down as I could see showed several wheals. The uvula was oedematous on the right side only, the situation of another wheal. Unfortunately, I had neither head mirror nor laryngeal mirror with me, but it is only fair to infer from the general manifestations of urticaria as well as the local ones in the pharynx and uvula, that the dyspnoea, cyanosis and prostration were produced by a wheal situated upon the epiglottis or within the larynx or trachea. Additional evidence rests in the sudden onset and simultaneous appearance of the urticarial rash with the dyspnoeic attack.

Adnephryn at once suggested itself as the remedy par excellence, but was not used because both the color and pulse were good and the patient evidently getting enough air, even though its passage was obstructed. Then, too, I feared that the irritation incident to any manipulation of the laryngeal structures might produce other wheals. A brisk purge of calomel and salts preceded by an enema was ordered. For the intense itching a pack of equal parts of milk and water with bicarbonate of soda and carbolic acid was applied, resulting in immediate relief.

I saw the patient an hour afterwards, the color was good, respirations about normal, pulse good, itching gone, but she was still quite nervous. The pharynx still contained a few wheals. The following morning the patient was normal except for a slight weakness incident to the catharsis.

California Bldg.

The Gates of Infection in Epidemic Cerebrospinal Meningitis.

N. Y. Med. Journ., Aug. 26, 1905.

This editorial calls attention to the interesting researches conducted recently by Westenhoeffer under the auspices of the German government. The results of twenty-nine autopsies made in the industrial district of Silesia, during an epidemic of considerably virulence.

The above observer finds that the invasion of the infections is through the pharyngeal tonsil. The latter was found enlarged, swollen, congested and oedematous in all the cases. The naso-pharynx was filled with a mass of muco-purulent material and the redness and swelling also extruded to the pharyngeal mucous membrane. The anterior portions of the nose was seldom involved. Either one or both middle ears were the seat of catarrhal or suppurative changes in 65½ per cent of all the cases. In 34 5/10 per cent of the cases the spheroidal sinuses were affected and in 27 6/10 per cent the maxillary sinus. In not a simple case did the infection find an entrance through the cribriform plate.

The moral of the above important observations is to clear the naso-pharynx of lymphoid overgrowth and to direct an asepsis to the nose and naso-pharynx during an epidemic of cerebro-spinal meningitis.

LEDERMAN.