that which removed the pain. It is worthy of note that, when pain was present, a tender spot was always recognised at the left side of the anterior cervical wall, and this tenderness disappeared on pressure being made upwards. My impression was that at this spot there was rigidity, compression, and consequent irritation of nerves. As the expansion of the uterus, assisted in the way described above, proceeded, it may be presumed that the nerve compression became less and less.

Cases of chronic ante flexion, with induration and compression of tissues round the cervix, are associated with abortion in not a few instances. They are amenable to treatment conducted on the principle of giving assistance to the unbending of the uterus in the early months. The cases in question appear to be not uncommonly overlooked, because the ante flexion is too generally considered as the natural condition of the uterus in the majority of cases. There is, however, a wide difference between ante flexion in gravid cases in regard to the condition of the tissues. If the tissues of the cervix are soft and natural, the ante flexion is possibly of little moment, as the obstacle to elevation of the fundus and unfolding of the uterine walls is not insurmountable. But when the ante flexion is coupled with induration, the raising of the uterus and the unbending of the organ are rendered difficult. The result is liability to abortion. The presence of rigidity and immobility is generally easily recognisable to the touch, and constitutes an indication of the necessity or otherwise for special treatment.

So far as interference with uterine expansion is concerned, there is in principle little difference between cases of retro flexion and ante flexion. The remarks above made have been limited to the case of ante flexion. There is the less necessity for advert ing to analogous difficulties in cases where, the uterus being in a chronic state of retro flexion, a tendency to abortion arises, because the nature of these latter cases is better and more widely understood, and because the utility of raising the fundus in such cases is pretty generally admitted.

Berkeley-square, W.

ERYTHRO-MELALGIA.

By JOHN E. MORGAN, M.A., M.D. OXON.,

PROFESSOR OF MEDICINE IN THE VICTORIA UNIVERSITY; CONSULTING PHYSICIAN TO THE MANCHESTER ROYAL INFIRMARY.

THE *American Journal of the Medical Sciences* for July, 1878, contains a paper by Dr. S. Weir Mitchell, in which he describes a rare vaso-motor neurosis of the extremities. To this disorder Dr. Mitchell has given the name of "Erythromelalgia," a Greek compound signifying an affection of the extremities, accompanied by redness and pain. The symptoms are characteristic, and usually commence with pain in one or both feet—often in the ball of the foot or of the great toe, or in the heel. Occasionally this pain involves the dorsum of the foot, and in rare cases the leg; more often it is confined to one or both soles. At first it is usually induced by any excessive exertion in standing or walking; but after a time it is experienced whenever the upright posture is assumed, or even when the feet merely hang down. The pain is usually of a burning character, and as the disease advances is described as intensely acute, often coming on in agonising paroxysms. It is relieved by the horizontal position and by cold, but is invariably aggravated by standing and walking, or by warmth, as by sitting near the fire, or even by the heat of the bedclothes. It is worse in summer than in winter. Pain, however, is not the only symptom referable to the feet; whenever any walking exercise is taken the parts flush, and in severe cases assume a dusky-red colour. Dr. Mitchell remarks that he has seen two cases in which the disorder, in its later stages, was associated with distinct evidence of spinal disease, while in one the hands were similarly affected. Dr. Sturge likewise, some years ago, reported an interesting case of this affection to the Clinical Society of London, in which, in addition to the foot symptoms, there was, after a time, wasting of the right leg and thigh. Dr. Mitchell looks upon the disease as peculiarly intractable, and seldom amenable to any form of treatment. In many cases the symptoms remain unchanged for years, and in rare

instances become gradually worse. In the paper to which I have referred, Dr. Mitchell has collected sixteen cases which correspond more or less closely with the description here given. Twelve of the sufferers were men and four women. The average ages of the men, where age is stated, was thirty-two, and of the women forty. In the great majority of the cases, the symptoms seemed to follow either some lowering constitutional disease, or they were ascribed by the patients to long-continued exertion in walking or standing, or to prolonged exposure to wet and cold. In addition to these cases collected by Dr. Mitchell, well-marked examples of the disease may be found in English and foreign medical journals and periodicals, though they do not materially extend our knowledge either of its etiology, its symptoms, or its treatment. During the last three years, five cases closely answering to the description given by Dr. Mitchell have come under my observation. Four of the patients were inmates of the Manchester Royal Infirmary and one was seen in private practice. Two of these cases I propose to describe somewhat in detail, as the symptoms in each were highly characteristic, and appear to me to shed some light on the true nature of the disease.

John C—, aged thirty-two, entered the Manchester Infirmary as an in-patient on Feb. 11th, 1888. The muscular system was well developed, and the man looked healthy and well nourished. He had a somewhat anxious, careworn expression, and though he called himself thirty-two, he looked fully forty years of age. He was an intelligent man, and graphically described the severity of his symptoms. By birth a Scotchman, he had, since the age of seventeen, been employed as a cowboy on a ranche in Texas, and in some of the Western States of America. He had led a healthy and active out-door life, and had been tolerably temperate in his habits. In the spring of 1886 he came to England as an *employé* and partner in an American novelty exhibition. While employed in the capacity of a showman he was constantly on his feet; indeed, he was in the habit of standing for twelve or fourteen hours daily. Some six months after landing in England he contracted gonorrhœa; the disease was accompanied by unusually severe symptoms, and he was more or less invalided for eight or nine weeks. On recovering and returning to his occupation, he found the soles of his feet very painful after standing or walking. The pain was of a peculiar burning character, and if he persisted in standing, or walked even a short distance, his feet would swell and assume a bluish-red colour. These symptoms became more and more severe, until at length he was altogether incapacitated from following his employment as showman. He was also unable to ride on horseback, as the pressure of his feet on the stirrups gave rise to acute paroxysms of pain. Moreover, as he could neither stand nor walk, he frequently relieved his feet by crawling on all fours along the ground. He stated that he had consulted numerous doctors in different parts of the country, but had derived little or no benefit from their treatment. Since the commencement of his illness he had lost considerably in weight. Although usually comfortable in the horizontal posture, he sometimes found his sleep disturbed at night, owing to the pain in his feet occasioned by the warmth of the bedclothes. He was then in the habit of protruding them beyond the bottom of the bed, in order that he might be relieved from the pressure of the blankets and coverlid. On examining his feet, it was found that sensation was unimpaired; there was neither hyperæsthesia nor paræsthesia. There was no loss of motion or wasting of the muscles of the feet or legs; neither was there any pain or tenderness along the spine, or retention or incontinence of urine or feces. The electrical reactions to the constant and interrupted current were normal. The external and deep reflexes were readily elicited. The feet were short, and somewhat thick and stumpy. All the flexor tendons over the dorsum of the foot, with the exception of that passing to the great toe, stood out prominently, and were somewhat tightly contracted. This toe was drawn downwards towards the plantar surface, and at the same time turned outwards beneath the second and third toes. There was nothing abnormal in the growth or conformation of the nails. The soles of the feet were often bathed in sour and somewhat offensive perspiration. On suspending the legs over the side of the bed, the feet almost immediately grew puffy and swelled, though there was no pitting on pressure; this was followed by a dusky bluish-red blush suffused over the toes, the soles, and the borders

of the feet. At the same time a burning, tingling sensation came on, accompanied after a time by intense pain. When he was directed to stand, and still more after he had walked but a few paces, all these symptoms were aggravated. In moving along the floor of the ward his gait was characteristic. With a view of removing, as far as possible, all pressure from the inside of his foot, he rested his weight on the outer border of each sole, shuffling over the ground; the two plantar surfaces facing each other during locomotion. On his return to bed, the pain, the congestion, and swelling soon passed off. This patient, who (as before stated) was admitted into the infirmary on Feb. 11th, remained as an in-patient till the 2nd of May following. Various local and general remedies were tried, some of which seemed rather to aggravate his symptoms than to afford him any relief. Among these I would mention bandaging and massage, both of which were persistently carried out, though both proved altogether useless. Electricity also, both the constant and interrupted currents, were prescribed for two or three weeks, but afforded no relief. Numerous remedies taken by the mouth exercised a merely palliative influence over his sufferings. As a last resource, the subcutaneous injection of morphia and atropia was tried, and proved highly efficacious, beneficial effects manifesting themselves almost at once. The needle of the syringe was passed directly into the sole of the foot on the inner side. This procedure was practised twice daily for about three weeks. Under this treatment a speedy improvement took place in all his foot symptoms, and on leaving the hospital, nearly three months after his admission, he was able to walk with freedom and comparative comfort. At that time all swelling and redness of the feet after exertion had entirely disappeared, and the pain experienced, even after considerable exertion, was altogether trifling. Five or six weeks after his departure from the hospital I received a letter from this patient, informing me that at that time he considered his recovery in all respects wellnigh complete.

Soon after the above case was admitted into the infirmary, a patient very similarly affected was received as an in-patient under the care of my friend and colleague, Dr. Dreschfeld. To his kindness I am indebted for the notes of this case. The man was admitted on March 24th, 1888. Two years before this the patient had occupied one of my beds, and at that time was suffering from chronic rheumatism. He was a collier by occupation, and necessarily much exposed to wet and to cold, being compelled at times to work for ten hours at a stretch in two feet of water. When a boy he suffered from typhoid fever, and had gonorrhœa some fifteen years ago. His present illness commenced twelve months before his admission. He was then attacked with violent pain in his feet, especially severe along the inner portion of the sole of each foot. Six or eight weeks after the pain he noticed a swelling and distinct blush about his feet. These symptoms were much aggravated by warmth or walking exercise, or even by allowing his feet to hang over the side of the bed. It was observed that the redness of the integument invariably followed any exertion which brought on pain. The pain was entirely limited to the internal plantar nerve and the parts supplied by it. This patient stated that he was always better in winter than during the summer months. When the warmth of the bedclothes induced pain in his feet, it was his practice to allow his uncovered legs to protrude beyond the bottom of the bed, with a view of keeping them cool. At times also he obtained great relief from enveloping his feet in cold wet cloths, or by turning upon them a cold-water tap. He had long noticed that the soles of his boots were more worn on the outsides than on the insides. In this case the knee jerk was normal. The man left the infirmary on May 2nd, considerably improved. At that time the blush, though present to a small extent, was not increased when the foot was in a dependent position, and though the painful spots on the plantar aspects of the soles were still present, the pain generally was far less severe than at the time of his admission. This patient returned to the infirmary on July 21st, eleven weeks after his discharge. At this time his symptoms were much aggravated. He suffered pain not only in both feet and in his legs and back, but also in his right arm and hand. It was found also that the knee reflexes were absent. In the month of April these reflexes were normal. The plantar reflexes also were increased, and there was general hyperæsthesia in both feet, as well as some

swelling and redness of the legs from the knees downwards. After some weeks he again left the hospital somewhat relieved, and has not since come under observation. Here we find that the symptoms, which in the early part of the year were almost entirely confined to the feet, had now extended up the leg as far as the knee. At the same time the knee jerk had disappeared, while the plantar reflexes were increased. The man had a pain also in his right arm and hand, and this pain was very much more marked along the area of distribution of the median nerve than of the ulnar.

In addition to these cases, another patient, Eliza H—, aged twenty-one, was admitted into the infirmary under my care on April 26th, 1888. She was a delicate girl, who had worked as a mill hand in a very hot room, and was compelled to stand for many hours of the day. She had never suffered from any serious disorder, but had frequently experienced distressing dyspeptic symptoms. In describing her ailments, she remarked "my feet feel quite comfortable while I lie in bed, but as soon as I get up and walk about, even for a short distance, I experience great pain in the soles of my feet, which look swelled, and feel hot and burning." She further stated that in walking she could not implant her right foot flat on the sole, but had to rest her weight on the outer side of the foot. The left foot she could place firmly on the ground without experiencing any pain. There were several painful spots on the right foot—one at the bifurcation of the posterior tibial nerve into the external and internal plantars, and others at the bifurcation of the digital nerves of the internal plantar in the cleft between the toes. The pain between the great toe and its fellow was, on pressure being exerted on this spot, very acute, while there was but little pain on pressure between the fourth and fifth toes. In this case the plantar reflex was well marked. This patient left the hospital early in May, considerably relieved. Here it will be observed that the affection was confined to one foot, the right; the left being comparatively unaffected.

In addition to the three cases here recorded, I have for several years had an opportunity of watching foot symptoms of a very similar character. The patient is a lady about fifty years of age, and of very delicate nervous organisation. For the last twenty-three years she has constantly experienced a pain in the sole of one of her feet. The pain is invariably aggravated by warmth and by exercise, which induce redness and swelling. Pressure on any portion of the internal plantar nerve causes her much suffering. She also is compelled to walk on the outside of her foot. She has been treated for this affection by various well-known members of the profession both in London and in the provinces, and has been sent to numerous continental spas. She has also worn cunningly constructed supports and sundry mechanical contrivances, besides being subjected to prolonged courses of bandaging. The affection, however, still remains as intractable as ever. I have been told by this lady that she can at any time estimate pretty accurately the state of her general health by the sensations in her foot. When her foot is painful her health is unsatisfactory.

In one other case of which I have notes, such symptoms as I have described are almost entirely confined to one hand. Long-continued work with the hammer first induced the attacks of pain, which were followed by redness and swelling. The pain is invariably worse when the arm is in a dependent position. Here, as in analogous cases, where the feet were similarly affected, the pain was almost entirely confined to the median nerve, as it is distributed over the palm of the hand and between the fingers.

The affection to which I have here directed attention is, it will be admitted, one of very considerable interest, and in some of its features altogether unique. As Dr. Mitchell remarks, it is a highly intractable disorder, the great majority of the cases remaining more or less stationary for ten, twenty, or thirty years; while in others the upper portions of the limbs affected gradually become involved, and occasionally the spinal cord itself is affected. In three of my cases the internal plantar nerve was the part solely implicated. Along the course of this nerve there were certain well-marked painful spots (points douloureux). One of these points was situated at the inner side of the ankle, where the posterior tibial nerve comes into close contiguity with the posterior tibial artery; another on the inside of the foot, where the internal plantar nerve crosses the common tendon of the flexor longus digitorum

muscle; and others were situated at the inter-digital clefts, between the three inner toes, at that portion of the foot where the internal plantar divides to form the digital nerves. At all these spots, even when the patient was perfectly comfortable in the horizontal position, pain was induced by somewhat firm pressure. Dr. Mitchell, who has very ably discussed everything bearing on this affection, and to whom we are primarily indebted for what we know regarding its etiology and symptoms, speaks of it as "an unrecognised type of spinal or cerebro-spinal disorder." He considers "the vaso-motor phenomena either as caused by the implication of the vaso-motor centres, or as due to a reflex affection caused by the pain or by the implication of the central organs, disease of which gave rise to pain." Dr. Ross, in his valuable "Handbook of the Diseases of the Nervous System," describes the affection as "a plantar neuralgia or neuritis [of the plantar nerves]." No doubt in some respects it does resemble both a neuralgia and neuritis, still there are important points of distinction which separate it to a greater or less extent from both these affections of the nervous system. In the first place, it is very rare to find neuralgia so symmetrical in its distribution as many of these cases are shown to be. It seldom attacks exactly corresponding parts of the two extremities with equal intensity, the pain in neuralgia being usually unilateral. The causes also which induce the paroxysmal attacks differ materially: neuralgia as a rule is aggravated, if not actually excited, by exposure to wet and cold; it is certainly not invariably increased by warm weather and warm clothing, as were the cases I have described. Again, although in neuralgia we frequently observe considerable congestion of the parts affected—as, for example, in "tic douloureux,"—still in these cases the congestion is of an active character, accompanied by heat and by a kind of rose-red blush. In erythro-melalgia, on the other hand, the congestion, especially in the later stages of the attack, is rather of a passive character; often there is no increase of temperature, and the parts affected look rather purple and livid than actually red. Again, although it is true that in cases of neuralgia we frequently observe both vaso-motor disturbances and abnormal secretions, still those symptoms are not so constant an accompaniment of the attack, nor are they so pronounced as in the affection I am discussing. Further, in all varieties of neuralgia, the patient has usually suffered from attacks of pain in various other parts of the body, while very few cases of neuralgia prove so intractable as this wellnigh incurable disorder. If, then, the objections to looking upon these cases as examples of neuralgia be manifold, can they more appropriately be deemed, not a functional, but an inflammatory, affection of the peripheral nerves—in other words, a neuritis? In answer to this question, it must be admitted that some notable symptoms usually observed in neuritis are seldom associated with this affection. Sensation, for example, is rarely perverted, or even blunted. Indeed, in the notes of some of my cases it is distinctly stated that there was not at any time either hyperæsthesia or paræsthesia. Then, again, there is rarely any paresis or fibrillar twitching of the muscles supplied by the affected nerve. Thus, in one of the cases cited by Dr. Mitchell—a case originally recorded by the late Dr. Graves—that acute observer remarks: "It is curious that the long-continued derangement in the circulation of the lower extremities, and the extraordinary pain this patient experienced daily for six years, did not produce any paralysis or any diminution of muscular power." Although, however, there is neither muscular weakness nor perversion of sensibility, still there are other symptoms which we are accustomed to associate with peripheral neuritis. Thus pressure over any part of the affected nerve invariably occasions pain of a more or less severe character, and this pain can be induced not only during the exacerbations (as is so frequently noticed in neuralgic attacks), but even during the remissions, when the nerves are in a state of painless repose. Furthermore, the pain is constantly increased by any change of posture which involves the stretching or tension of the nerves. Moreover, in several of the cases to which I have alluded, reference is made to an ascending neuritis observed in the later stages of the disorder, wherein the inflammatory changes seemed to travel upwards, and in some cases actually involved the spinal cord itself. If, however, the disorder must be looked upon as a neuritis, it is evident that it is a form of neuritis in which only certain parts of the nerve—probably the sheath—are involved. This

affection is, in fact, a peri-neuritis rather than a parenchymatous or interstitial neuritis. Such forms of neuritis are occasionally observed in cases apparently due to toxic or infectious influences, notably where the system is affected by such disorders as syphilis and rheumatism. Dr. Ross, in his work, to which I have already referred, seems to think that the cases of obstinate pain in the soles of the feet described by Dr. Elliotson as following gonorrheal rheumatism really belong to this category. In one of my cases the patient had experienced a severe and somewhat protracted attack of gonorrhœa; Dr. Dreschfeld's patient, also, had at one time suffered from the same complaint. Taking the cases of erythro-melalgia which have been recorded by various writers, it may be stated generally that the great majority of the patients, in detailing the history of the diseases from which they had previously suffered, specially alluded either to rheumatism or syphilis, or gonorrhœa, or gout, or some variety of fever. It would seem, therefore, that these disorders engender within the system certain septic changes, which specially predispose it to the occurrence of some such form of peripheral neuritis as we here observe. From the considerations here urged, we are probably justified in believing that the paroxysms of pain in these cases are due to a peri-neuritis dependent on a cachectic taint in the system, while the vaso-motor phenomena are due to a reflex irritation starting from the affected nerves, and thence transmitted to those vaso-motor centres in the cord with which the experiments of Goltz have made us acquainted—an irritation which, if prolonged, we might not unreasonably expect to be followed by vaso-motor paralysis.

Manchester.

NOTE ON THE KNEE JERK IN DIPHTHERIA.

By W. B. HADDEN, M.D. LOND., F.R.C.P.,

ASSISTANT PHYSICIAN TO ST. THOMAS'S HOSPITAL, AND TO THE HOSPITAL FOR SICK CHILDREN, GREAT ORMOND-STREET.

THE condition of the knee jerk in diphtheria first occupied my attention three years ago. Others, no doubt, have investigated the same point, but very little information exists in medical literature on the subject. I regret to say that my observations have been limited, and that I have been unable to follow out some of the points, to which I shall briefly allude.

It is, of course, a matter of common knowledge that in the paralysis following diphtheria the knee jerk is usually feeble or absent. It is, however, not so well known that, even when the membrane is present in the fauces, the knee jerk occasionally disappears. In a certain proportion of cases, on the other hand, it is exaggerated; but whether this is succeeded by abolition of the phenomenon, as is sometimes the case in diphtheritic paralysis, is a point on which I cannot positively pronounce. The absence of the knee jerk in diphtheria is, no doubt, due to the same nervous derangement (whatever that may be) which causes actual paralysis in the later periods. Paralytic affections of the soft palate in the acute stage, though often slight and transitory, are, I believe, of greater frequency than is usually supposed. Sudden death from cardiac failure is in all likelihood dependent on some paralytic condition affecting the nervous apparatus of the heart. The opinion which I am inclined to entertain is that the nervous system not uncommonly undergoes morbid change, though there may be few objective signs during the first few days of the disease. Post-diphtheritic disorder of the nervous system may be indicated solely by the absence of the knee jerk, and this may persist for many months. In this respect the following case is of interest.

The patient was a little girl aged seven, who was brought to me at the Hospital for Sick Children in May or June last. She had had a severe attack of diphtheria in November, 1887, and was kept in bed until the middle of January. The mother was carefully questioned as to the occurrence of paralysis, but there was no evidence of it. Although during the next few months the child had improved, she still remained thin and emaciated. When I saw her I could find no definite signs of disease, though she appeared listless and without energy.