CHART 3.

Chart illustrating the difference between the temperature in the mouth and that in the rectum (dotted line).

The failure to maintain the gradual increase in weight during the month of March was clearly due to the attack of influenza.

There are two points in this case to which I wish to direct attention. One is the remarkable gain in weight, probably from about 7 st. (certainly from 7 st. 7½ lb.) to 10 st. 2½ lb. This occurred notwithstanding the fact that the disease was not arrested, as is shown by the presence of cough and expectoration and the discovery of numerous tubercle bacilli in the expectoration when the patient left the hospital. This very considerable gain in weight I attribute to the large quantity of food which the patient took and assimilated. That a patient may gain in weight and yet the disease may not be arrested is no new observation, but with an ordinary hospital diet it is, in my experience, a rare event for a patient to gain 3 st. in weight without the disease undergoing arrest. What I observed in this case accords with my experience of others treated on similar lines (rest and hyper-alimentation), and I believe that *the larger the quantity of food administered the less is the significance to be attached to a gain in weight as evidence of the arrest of the disease.* I do not say that a gain in weight is not of importance. On the contrary, it is a fact of the greatest moment, for without it the progress of the disease is almost certain, but it is not by itself, even when remarkable in amount, a proof of arrest, when the quantity of food taken has been unusually large. I so frequently observe reference to the increase in weight in patients submitted to this method of treatment that I think it desirable to draw attention to the fact just stated.

The second point is the difference between the oral and the rectal temperatures (Chart 3). You are doubtless aware that at Nordrach the patients take their own temperatures in the rectum four times daily, and so far as I have been able to learn they do so without expressing any particular objection. The necessity for such a proceeding, which for obvious reasons does not appear at first sight to commend itself, has not hitherto been recognised in this country. The oral temperatures in the evening were from 0.5° to 1° or even 2° F. lower than the rectal. The oral temperatures in the morning were usually slightly above the rectal but sometimes one-fifth of a degree below. There was thus a far greater difference between the temperature in the mouth and that in the rectum at the higher than at the lower level.

The important question whether the patient is to remain at rest or to take exercise and the amount of exercise to be taken is to be decided by the presence or absence of pyrexia. It is therefore of great moment to know the real temperature of the body, not merely that of the mouth. Now it is clear that the difference between the temperature in the mouth and in the rectum may be as much as one degree or one and a half, or even two degrees. It has, I believe, been shown that in order to invariably obtain the true temperature in the mouth it is necessary to close the lips for 20 minutes before inserting the thermometer. If there were a constant difference between the oral and the rectal temperature it is obvious that it would only be necessary to know this and to take it into consideration in reading the temperature as taken in the mouth, but, as these charts prove, the difference is not constant but extremely variable. I am therefore led to the conclusion that the only

trustworthy method of ascertaining the temperature of the body is to take it in the rectum.

Hospitals for consumption have in the past conferred incalculable benefits on sufferers from tuberculosis, but owing to the necessary limitation of the period of treatment it has not been possible here to set before ourselves as an objective in all suitable cases the complete arrest of the disease, and had we possessed a sign-board we should, I fear, have been compelled to imitate the humble craftsman and imprint upon it the words "Repairs neatly executed." Let us hope that in the future by greatly extending the period of treatment it may be possible in a much larger proportion of cases to secure such complete arrest. Herein, indeed, lies the difficulty in dealing with hospital patients, especially such as have families or relatives dependent upon them for support, as they cannot in most cases afford the time necessary to secure permanence of the reparative changes taking place in the lungs. How long a period of complete rest is generally necessary to effect this must have been appreciated by all who have had

an extended experience in the treatment of such cases.

Clarges-street, Mayfair, W.

THE PREVENTION OF SYPHILIS.¹

By FRANCIS H. WELCH, F.R.C.S. ENG.,

SURGEON-COLONEL (RETIRED), MEDICAL STAFF.

IN entering upon the question of the prevention of syphilis there are two features which require full recognition, as upon them this problem rests, one etiological and pathological, physical—the cause, its multiplication in the frame, and mode of transfer from individual to individual; the other socio-moral, ethical—that phase of our civilisation, promiscuous intercourse, by which the cause is mainly spread in the community. It is the relationship of the latter to the former as determining the outcome—prevalence of disease—which compels us to consider the question as a whole and not merely from a hygienic standpoint when deciding on the action to be taken under our present civilisation and polity.

The cause, which everywhere (and the same pertains in time from the earliest recognition of the disease) demonstrates the presence of syphilis through similar manifestations is, we infer, an organic entity—a micro-organism—not yet identified. While capable of transfer to certain animals (though this is denied by some) it is practically a human malady which is communicated through contact (acquired) or through parentage (inherited). As a rule it enters the body by surface inoculation, and in a lesser ratio by the blood or semen. In the first case it sets up active changes at the site and in the contiguous lymph glands suggestive of attempted preclusion and localisation of the virus, yet followed for the major part by a true systemic infection; in the second case it is directly transmitted to the system of the individual or embryo through these media. During the primary and secondary periods the virus is capable of being inoculated, and during the latter period it is capable of being transferred with the blood or semen, but how far beyond the period of active changes it can be transferred to the embryo seems an undetermined point.

Like disease in general the origin of this special malady is obscure. Whether it is primarily a pathogenic germ apart from the human frame, or is primarily innocent yet pathogenically evolved under a favourable human environment, or is derived from normal elements adversely modified; whether, again, it arose in one or several sections of the human race; or whether it is capable of a spontaneous recrudescence in the future if conditions permit—these are not merely points of academic interest but they have an important bearing on the eradication of the cause. For present practical purposes we know of it only as arising from pre-existent disease and as transferred throughout the world at large in the selfsame methods, and while

¹ A paper read before the British Medical Association, August 3rd, 1899.

apparently no special marriage bonds such as monogamy, polygamy, or polyandry, have been connected directly with its presence or diffusion, unrestricted sexual intercourse, promiscuous as to time and person, has everywhere characterised its prevalence. In the local conditions produced by prostitution an environment has been generated favourable to the multiplication and envenoming of the virus and by means of this agency the virus has become extensively dispersed throughout humanity.

With these features clearly borne in mind it is apparent that we can approach the problem of prevention in two ways—through hygiene, having for its object the preclusion of lineal descent of the virus, or through ethics directed towards the extirpation of the great diffusing medium. While both agencies operate towards the suppression of the disease the ethical procedure makes, in addition, for the suppression of the vice. For complete results the first necessitates an abnegation of the liberty of the subject and a strict compliance with stringent sanitary rules, and the second a moral millennium in which the ascendancy of the higher faculties of our composite nature over the animal instinct and physiological impulses shall have been brought about or the brotherhood of humanity be the dominant factor of conduct. It may be said that both these standpoints may be dismissed from the sphere of present discussion on the ground that they are incapable of being fully realised under the conditions which obtain, yet it is no less apparent that they contain the principles on which we must proceed to obtain the present attainable, even though that level be far below what is possible or desirable.

But when this question of prevention comes up for inquiry and solution there is one aspect of the subject which requires prior consideration, bearing as it does on the necessity or desirability of action. I refer to the statement which has been made by some that the natural course of this disease is to retrogress, apparently from an acquired immunity of the race through exhaustion of the necessary nidus by existent and anterior widespread disease. This view is brought forward in the "Encyclopædia Britannica."² After stating that "syphilis in the United Kingdom at the present moment is in the stage of an epidemic in its decline," the writer goes on to say: "A time must come when the soil is practically worn out, when it becomes so poor that the organism grows only in a stunted form producing a mild disease, till in time it ceases to grow altogether;" yet adding the cold comfort that: "After lying fallow for a time the soil may recover its power and the disease be renewed in a more virulent form." As opposed to the first statement we have the *British Medical Journal*³ noting that "those best acquainted with it state that venereal disease is on the increase among the civil population of this country"—an opinion which probably many of those in hospital practice in the metropolis and in the larger towns will support. And in respect to this theory of retrogression, while it seems tenable on the hypothesis that the disease is universally present and that the acquired immunity is indefinitely transmitted, yet this necessary basis is wanting, and, moreover, fresh soil is in production with each succeeding generation to furnish the necessary nidus for the growth and the maintenance of the vigour of the virus. If the natural history of the disease is to self-effacement the nations of the Old World, such as China and India, should by their present state illustrate it or at least closely approximate thereto, but the available data culled from native sources and the returns from troops located in these countries are far from supporting such a view; here frequency and intensity of disease are features certainly not absent. A decline, partial in extent, may be perceptible in this country due to increased knowledge, to personal hygiene, and to improved therapeutics, and possibly not unconnected with sanitary regulations now non-existent, yet these are not innate and constant, but are acquired and fluctuating factors, and I submit that this theory of natural emancipation from this disease is a fragile reed to rest upon. That, like other infectious diseases, it has its ebb and flow, seen in statistical undulations, is a feature deducible from its cause and falling under the one same explanatory theory.

Nor does it seem possible to ignore the hygienic side of the question and to rely upon the extirpation of prostitution through moral and social agencies. Nor does it seem

desirable to rest satisfied with hygienic means and so leave prostitution, the main diffusing agency, free to renew the never-ending task of meeting its dire physical results. Yet we need to remember that it is the comparative hopelessness of assistance from moral and social forces in the near future which constitutes the cogency of the demand for sanitary rules to meet the grave existing facts.

To suppress all disease as far as possible by eliminating the cause and curing the results and so to reduce the sum total of suffering and misery in this world is a duty inseparable from our profession, and it is self-evident that this malady is one which, physically viewed, from its extent and baneful influence upon the individual and the race beyond any other known disease specially claims from us the use of every weapon, natural or otherwise, which Providence has placed in our heads or hands. And ethically viewed, appealing to us as philanthropic men and good citizens, I submit that no socio-moral argument has been advanced sufficiently cogent to justify us in placing this disease in a different category from other diseases and so withholding any power which we possess for its suppression or cure.

Hence, then, this disease being amenable to hygienic principles and claiming from us all possible alleviation what data have we to guide us as to what is practically attainable? We have the principle recognised in this country in other dangerous infectious diseases that not merely actual transfer but acts conducive thereto are rightful grounds for State interference by law, and to go no further than ourselves we have also evidence accruing during the last 35 years to the effect that through surveillance of prostitution and segregation of the affected we can diminish the amount of this special disease and attenuate its virulence. I base these deductions on the conclusions arrived at by official committees, on the statements of public officials, on the comparison of military stations under sanitary rules and those not under sanitary rules during the same period and of districts at different periods when these opposite conditions existed, and on the outcome from limited communities under favourable circumstances for testing purposes. (a) In respect to reduction of prevalence the Home Committees of 1868,⁴ 1869,⁵ 1870,⁶ 1879,⁷ and 1882,⁸ leave no doubt; and the statements of the Government of India on the results following the Cantonment Rules of 1868⁹; of the Calcutta Committee of 1881¹⁰; the Fort St. George (Madras) returns following the Act and the recrudescence of disease in 1877 on its suppression¹¹; the evidence of the Surgeon-General in Bombay in reply to the

⁴ "In places where the Act is in operation not only is the amount of disease less but the cases exhibit a milder character the physical evils have abated." (Report of the House of Lords Committee, 1868.)

⁵ In view of the "decrease of disease" and other recognised good results of the 1866 Act an extension of it was recommended. (Report of the House of Commons Committee, 1869.)

⁶ "Acts have reduced the worst forms of the disease and directly and indirectly promoted the objects sought for." "Primary syphilis had been reduced from 120 per 1000 in 1865 to 54 in 1870, while no such lessening had taken place in the unprotected stations which varied from 99 to 113." (Report of the Royal Commission on the motion for the repeal of the Contagious Diseases Acts.)

⁷ "In stations not under the Act the proportion was 20 against five in those under the Act." "In the first years of the Act the admissions from primary syphilis in certain registered stations were 91 per 1000, reduced to 35 in 1877 and 40 in 1878." Cases of diseases among the prostitutes examined amounted in 1865 to 76.24 per cent. and in 1879 to 7.40 per cent. (Report of the House of Commons Committee, 1879.)

⁸ The House of Commons Committee re-appointed in 1880 reported in 1882: "Has successfully served" for "the diminution of venereal disease." "The actual statistics show that the reduction in secondary syphilis due to the Contagious Diseases Acts has been 39 per cent." "The extent to which the Acts have diminished syphilis in the subjected districts appears of itself to establish the hygienic utility."

⁹ "The introduction of rules into cantonments was followed by a marked decrease in the prevalence of venereal disease among the troops and the decrease continued to manifest itself with more or less fluctuation until about 10 years ago (i.e., 1887) when an increase became observable." "Increase has been materially checked by the operation of the rules in question and by the establishment of lock hospitals." Sir A. D. Home gives the average of five years preceding the operation of the Contagious Diseases Act in Bengal as 314 per 1000 and from 1877 to 1881 as 241 per 1000. (House of Commons Return, East India (Contagious Diseases), 1888.)

¹⁰ In the Calcutta garrison primary syphilis was reduced from 10 per cent. to 1.7 per cent., in women from 25.6 per cent. to 3 per cent., and secondary syphilis "had ceased to exist." When the rules were withdrawn the garrison ratio advanced to 30.24 per cent. In the police force the same features pertained. The Surgeon-General remarks: "It is impossible that any sanitary improvement could have been brought to more absolute demonstration than the vast decrease of syphilis in Calcutta and its practical disappearance from the garrison." (Ibid.)

¹¹ "Its introduction was followed by a great diminution in the ratios of admission from constitutional syphilis." In 1863-68, prior to the Act, the average ratio from primary syphilis was 129.72 per 1000; from 1869 to 1876, during the Act, it was 58.84 per 1000; from 1877 to 1886, on its withdrawal, it was 146.03 per 1000. (Ibid.: Minute of Commander-in-Chief, Madras.)

² "Surgery—Venereal Diseases," *Encyclopædia Britannica* (ninth edition).

³ *Brit. Med. Jour.*, April 22nd, 1899.

Madras Committee of 1884¹²; the information furnished to the Venereal Committee of 1864 by Sir Henry Storks from the Ionian Isles and Malta¹³ and by Mr. Inglott, the Controller of Charities in Malta¹⁴; the official reports to the Colonial Secretary by the Governor of Malta in 1887, including the military and naval experience,¹⁵ and those by the Administrator of Hong-Kong¹⁶ and the Administrator of Singapore and Penang;¹⁷ the results of closure of 10 lock hospitals in India in 1885¹⁸; the 16 years contrast between protected and unprotected stations in this country¹⁹; the statements made by the *British Medical Journal* in the early years of the Acts,²⁰ and the latest returns from India compared with periods under protection,²¹ surely furnish safe grounds for the conclusion, despite the apparent tendencies of some statistics to an opposite inference, that diminution of syphilis in frequency has been the outcome of the sanitary regulations known as the Contagious Diseases Acts. Indeed, as syphilis results in at least 90 out of every 100 cases from impure sexual intercourse and as the Acts undoubtedly prevented many diseased prostitutes from acting as foci of dispersion, how could the results be otherwise than as attested by the above sources of evidence? (b) In respect to virus attenuation, the statements made by Mr. J. R. Lane from lock hospital experience,²² corroborated by Dr.

Payne's experience as supplied to the Calcutta Committee of 1882²³; the contents of Mr. Lane's letter to the *British Medical Journal*²⁴; the summaries of the *British Medical Journal* in 1868²⁵; the statement of Mr. Inglott on Malta²⁶; that of the House of Lords Committee in 1868,²⁷ and that of the Calcutta Committee in 1881²⁸; the opinions of the Madras, Bombay, and Calcutta medical officers referred to by the Commander-in-Chief, Madras²⁹; the figures on duration in hospital and invaliding brought forward in the Indian Viceregal Council, 1897, by the Commander-in-Chief³⁰; the memorandum of the Army Sanitary Commission, March 9th, 1897; the Indian Lock Hospital experience,³¹ that of Hong-Kong,³² that of Japan as stated by D. I. G. Fisher, R.N.,³³ and of Singapore and Penang as stated by the Colonial Surgeon³⁴; and the increased deaths and invaliding in India in later years³⁵; all these concur in the inference that the intensity also of syphilis has been markedly reduced by hygienic means. That by early recognition of the disease, by segregation of the sufferer from all irritating influences, and by local personal hygiene and therapeutics we form an environment curtailing the growth and evolution of the virus, is, I submit, reasonable, and that the practical data support this view is,

against 37; secondary syphilis, 56 against 37; tertiary syphilis, 52 against 55. Mr. Lane remarks: "Of the protected cases the primary sores were milder in character, secondary sores more severe among the ordinary women; early treatment had a most beneficial influence on the character and duration of the diseases." (*Brit. Med. Jour.*, Feb. 15th, 1868.)

²³ "Among the registered women there has been a very marked change in the type of the diseases; in fact, the worst forms have practically disappeared." "In Calcutta the improvement among the European soldiers has in this respect been equally marked." (*House of Commons Return, East India, May 14th, 1888.*)

²⁴ "It is indeed a most remarkable fact that this legislation has not only greatly diminished the quantity of the disease, but has also had a very marked beneficial influence upon its quality, in its secondary no less than its primary stage. They appear to have imbibed so small a dose of the poison or have imbibed it in so unconcentrated a form that they may be safely left to get rid of it by their own unaided powers." (*Brit. Med. Jour.*, Dec. 12th, 1874.)

²⁵ *Brit. Med. Jour.*, Nov. 7th, 1868.

²⁶ "Syphilitic affections being thus timely discovered and forcibly placed under medical treatment are generally primitive in their character and consequently easily overcome. Cases of consecutive syphilis are comparatively rare and none present those horrid forms so common where the disease is under no surveillance." (*Report of the Venereal Committee of 1864.*)

²⁷ Vide Note 3.

²⁸ "The Act has tended to diminish the extent of venereal disease and has so modified the type that the worst forms are now almost exceptional." As to its influence on the general population, there was "An important diminution in the extent of primary syphilis and the type was milder than in previous years." In 1884 following the suppression of the Act "The men (police) have suffered from the most malignant types of the disease, a thing unknown in previous years," and "30 per cent. of the garrison were invalided with syphilis." (*House of Commons Return, East Indies, May 14th, 1888.*)

²⁹ "It is worthy of note that the medical officers in Calcutta, Madras, and Bombay bear testimony to the favourable change in the type of disease effected by the operation of the Act." "Type of such disease as does exist (in Madras) is much less severe than that which formerly existed." (*Ibid.*)

³⁰ "Average stay in hospital for secondary syphilis, in 1887, 26 days; in 1895, 31½; invaliding augmented fourfold." Lord Roberts in the House of Lords stated that the invaliding two years before relaxation amounted to 126 and in 1894-95 it had increased to 648.

³¹ That of Bengal and Madras is not noted, but in Bombay "in all three stations not only did venereal disease greatly increase but the type of the disease was more virulent than in the stations where the hospitals were kept up." (*House of Commons Return, East Indies, May 14th, 1888.*)

³² "Of a type very amenable to treatment, formerly the type of the disease was of the most virulent character." The Senior Naval Medical Officer in 1875 wrote: "In 1849-50-51 I was surgeon of H.M.S. *Reynard* have a vivid recollection of the sufferings of the crew from syphilitic disease in Hong-Kong; a large proportion became affected, hard chancres and large phagedenic sores, secondary symptoms were also of frequent occurrence often requiring invaliding." "The altered state of things is very striking. During the year and three-quarters in which I have had charge of this hospital I have not seen a case of regular Hunterian chancre contracted at Hong-Kong and only a few sores which healed readily." (*House of Commons Return, 1887.*)

³³ "The lock hospital system combined with careful examination of all women belonging to the licensed brothels has simply stamped out the old virulent forms which has for years been associated with the name of Japan." (*Ibid.*)

³⁴ Speaking of the result of temporary closure of the lock hospital the Colonial Surgeon stated that in October, 1882, the percentage of diseased women was 11.34, and that in September, 1883, on re-examination, it was 44 (53 syphilis and 15 gonorrhoea). "Moreover, the type of syphilis and the ravages it had committed was such as required lengthened treatment in hospital and such as was seen only when the Act was introduced."

³⁵ In 1880 secondary syphilis gave ratios per 1000 of strength, deaths 0.06, invaliding 1.62, and in 1896 0.23 and 6.36 respectively. The primary ratio in 1896 was not quite double that of 1880, while the secondary ratio was nearly four times that of 1880 with about the same augmentation for deaths and invaliding. In 1896 it was noted that "the syphilitic cases were of a most virulent type."

¹² Not only have admissions from gonorrhoea and primary syphilis among the British troops been reduced, but the type of the disease has been modified and the constitutional affections have almost disappeared. (*Ibid.*)

¹³ "The disease may be said to have almost disappeared in the islands of Corfu, Zante, and Cephalonia." And in Malta, "in 1864, on April 12th out of a force of 6192 only five were under treatment; on Oct. 21st there were only eight cases in the whole garrison." (*Report of Venereal Committee of 1864.*)

¹⁴ After noting that "the evils arising therefrom have never made their appearance in those terrible forms which are common in many other parts of Europe," Mr. Inglott refers to the year 1859 when in sequence to temporary abeyance of the Contagious Diseases Act "the awful consequences of non-restraint soon became apparent and an ordinance was re-enacted in 1861" which "had the effect of checking public prostitution to a great extent, besides of almost annihilating the disease." "Were it not for the inoculation from imported affections by foreigners and others the disease would long since have been extinct." (*Ibid.*)

¹⁵ Sir T. Simmonds wrote: "There can be no doubt that in Malta the result of the administration of the existing law has been hygienically most successful." The Principal Medical Officer, Army Medical Staff, wrote: "The Acts have attained their object by preventing the spread of the disease and by mitigating its type. In 1884 the Royal Munster Fusiliers, 374 strong, gave no admission from syphilis; the Duke of Connaught's Light Infantry, 900 strong, which on arrival from England landed 28 cases of venereal disease had on Nov. 15th, 1886, only one case of gonorrhoea in hospital; the 'Black Watch,' 740 strong, which arrived from Egypt with 38 cases have at present none in hospital." The Principal Medical Officer, Royal Navy, wrote: "Venereal disease not only few in number but very slight in character." (*House of Commons Return, Sept. 10th, 1887.*)

¹⁶ The laws have had most beneficial effect in preventing the development and communication of a very virulent form of venereal disease for which the colony in former years had an unenviable notoriety. Cases of syphilis now few in number" (*Ibid.*). The Senior Naval Medical Officer, writing in 1870, says: "Judging from my own experience I should say that syphilis has all but disappeared from the colony." Compare these comments with the Army Returns for 1896, which give 65.3 per 1000 admissions for primary syphilis and 106.1 for the secondary form.

¹⁷ "Disease has been largely checked." The Colonial Surgeon states: "The ravages, local and constitutional, formerly made by syphilis can only be thoroughly known and the beneficial effect of the ordinance fully appreciated by anyone who has held a similar position as I have done in Singapore and Penang before and since it came into force some 14 years ago." (*Ibid.*)

¹⁸ From primary syphilis the advance was:—Bengal from 76.9 to 141.6 per 1000; Madras from 65.1 to 109.9 per 1000; and Bombay from 132.2 to 168 per 1000. From the statement of the Commander-in-Chief in the two Madras stations secondary disease increased in one from four to 40 and in the other from 22 to 34. The Government of India Proceedings record "a decided and progressive increase. The percentage compares most unfavourably with the percentage at the stations where these institutions were maintained." (*House of Commons Return, 1888.*)

¹⁹ The primary form was reduced from 130 to 50 per 1000 and returned to 138 per 1000 when protection was withdrawn, while the unprotected ratio from an initial 116 rose to 118 per 1000 and after abolition to 160 per 1000. In the words of Dr. Parkes: "In eight years the State was saved nearly 10,000 cases of syphilis." In secondary disease the protected ratio was 22.5 against 36 per 1000 in the unprotected, and five years afterwards the former reverted to 34 per 1000.

²⁰ For example in 1870: "Thus it is shown beyond all dispute that both for men and women the amount of syphilis is diminished by one-half where the Acts are in operation."

²¹ The Commander-in-Chief at the Viceregal Council, India, 1897, gave the increase as 137 per cent. in primary disease and 188 in secondary disease from 1887 to 1895; and secondary disease in 1895 as four times more prevalent than in 1873. In 1880 the primary ratio was 87.9 per 1000 and the secondary ratio was 25.0 per 1000. In 1896 the ratios were 159.3 and 97.7 per 1000 respectively, while the ratio for all venereal diseases was 249.7 against 444.4 per 1000.

²² Syphilis characterised 80 per cent. of ordinary women, 41.5 per cent. of the Government women; the average duration in hospital was in ordinary cases against Government cases: primary syphilis, 49

to my mind, unquestionable. Indeed, the conclusions drawn on both these points from general sources are in accord with my own personal experience derived from an army practice of over 30 years.

Time does not permit the quotation of all those references mentioned and I limit myself to two, both from public sources, one bearing on the early years and one on the late years of the period of experience, one referring to home and the other to Indian results. The *British Medical Journal*³⁶ stated that "the evidence from all quarters is remarkable that both as regards the men and women not only has the quantity of the disease been diminished but its quality has been modified for the better in the most marked manner. Not only has the proportion of syphilitic disease to the milder affections been much diminished, but the syphilis which has occurred has been of a mitigated character and secondary affections have been rare." The Army Sanitary Commission, reporting on March 9th, 1897, after referring to their former statements that the Acts had been a failure, go on to say: "We now have proof, however, that the measures heretofore adopted in India, imperfectly administered as they often were, nevertheless effected a sensible influence in checking increase in the number of British soldiers admitted into hospital from venereal disease and in mitigating the severity of the disease itself, preventing its assuming that virulent and destructive character which it has now attained."

But the good which is to be obtained from sanitary measures will be dependent on the degree of observance of certain essentials for complete success—(a) an application of them co-extensive with the disease in range, in class, in sex; and (b) a uniform and continuous enforcement. These features were but partially present in the old Acts.

However, this country and most of our dependencies at the present time are in the unfortunate condition of communities in which vice, solicitation, and disease have a free hand. We have no public restrictive measures, and our present home state, be its relative bearing to the original ante-Act state what it may, is beyond doubt bad enough, while the army returns are a conclusive index to many foreign stations. Noting, however, the relation of prostitution to disease and the many contributing elements—social, moral, municipal—I submit that the right future course to adopt in the prevention of syphilis lies not in any one isolated line of action but in bringing to bear concurrently on the disease and its diffusing agency all the powers which our present civilisation places at our disposal; and these are: (a) moral and religious advance—to strengthen the higher elements of our nature; (b) proper housing of the poorer classes—to render possible the growth of modesty and chastity, and rational healthy amusements for mind and body away from our crowded thoroughfares—to curtail the present fostering influences of precocity and animalism; (c) social action—to facilitate marriage for those not having "the gift of continence," to remove the possible resort to selling the body for prostitution as a means of livelihood, and to curtail the twin contributing vice—intemperance; (d) municipal action—to prevent solicitation in the streets and public gathering-places and so remove the present temptations to the youths of both sexes, to eliminate agencies and localities for the pursuit of vice, and to make in all ways prostitution difficult; and (e) hygienic rules—to necessitate from the votaries of prostitution proof (in the shape of a periodical certificate satisfactory to a magistrate) of physical purity as a guarantee that in their illegitimate practice they are not disseminators of disease as well as of vice, to make sufficient and proper hospital accommodation for those not in a position to obtain adequate private treatment, and to furnish the means for compulsory segregation when necessary.³⁷

Yet in furtherance of these ends we have to remember that in this country we cannot go in advance of public opinion and that there is a limit to restrictive measures in the power of the people, and herein lies the important and pressing necessity of educating the public generally (possibly even some of our own profession) on this subject, so that reason and right action may prevail in the end; and this seems to me a primary essential towards any future measure for the prevention of venereal disease.

Lee, S.E.

³⁶ Brit. Med. Jour., Nov. 7th, 1888.

³⁷ This groundwork of action is based on the existence of prostitution and its object is to provide a safeguard against this openly tolerated practice being detrimental to public health as well as to public morals, the burden of proof being thrown on the pursuer of this permitted free trade in immorality.

TEN CASES OF VAGINAL HYSTERECTOMY.

By ARNOLD W. W. LEA, M.D., B.S. LOND.,
F.R.C.S. ENG.,

ASSISTANT LECTURER ON DISEASES OF WOMEN AT OWENS COLLEGE;
ASSISTANT PHYSICIAN TO THE CLINICAL HOSPITAL FOR
WOMEN AND CHILDREN, MANCHESTER.

THE operation of vaginal extirpation of the uterus and appendages has been very extensively practised during the last few years and the results obtained have been very satisfactory. Originally introduced as a radical treatment for cancer, its scope has been extended to other diseased conditions, such as uterine fibroids if below a certain size, chronic inflammation of the uterus with advanced bilateral disease of the appendages, pelvic suppuration, and, lastly, as a remedy in intractable cases of prolapse. The improvements in technique have been followed by great reduction in mortality. The best results have been obtained in extirpation for fibroids and inflammatory conditions, some operators having published series of 40 and 60 cases without a death. In cancer the mortality has been somewhat higher, especially if the disease has advanced beyond the early stage. It has also been clearly shown that for malignant disease extirpation of the uterus can only rank as a palliative procedure since recurrence sooner or later is almost inevitable. There can be little question, however, of the great relief and prolongation of life which is generally afforded by the operation. I have performed vaginal extirpation in 10 cases—in five cases for cancer of the uterus (the cervix, four; the body, one); in three cases for multiple fibroids; and in two cases for chronic inflammatory affections, combined in the latter with removal of the appendages. In this series of cases one death has occurred due to intestinal obstruction nine days after operation; all the other cases made rapid recovery. The notes of the cases are briefly as follows.

CASE 1. *Adeno-carcinoma of the body of the uterus*.—A woman, aged 51 years, was seen first in June, 1896, on account of uterine hæmorrhage which had commenced six months previously. She had had one child 25 years before and the menopause came on in her thirty-ninth year. Examination revealed a narrow senile vagina. The cervix was atrophic; the body of the uterus was bulky and retroflexed. No adhesions could be discovered. The sound passed three inches and the cavity felt large. The uterus was explored under anæsthesia and small hard masses of growth were removed. Microscopically the fragments showed typical adeno-carcinoma and immediate extirpation was advised. The patient would not, however, at first consent, but later—in October, 1896—she was admitted into a home and extirpation of the uterus was performed. The ovaries and tube were normal and they were not removed. The patient rapidly recovered and she left the home in three weeks. The uterus on section showed extensive growth infiltrating the muscular wall; the peritoneum was healthy. The patient remained quite free from recurrence for 12 months. She then began to feel intense pain in the back and down the left leg and examination revealed a recurrence of growth at the side of the pelvis. A little later she developed enlargement of the lumbar glands and ascites. She then commenced with symptoms of lung trouble, hæmoptysis, and cough, the result of extensive secondary growth. She gradually sank and died 18 months after operation.

CASE 2. *Carcinoma of the cervix*.—The patient, a woman, aged 53 years, was sent to me by my friend Dr. Chapman in March of last year. For two months she had noticed bleeding from the vagina and a month previously to the time when she was seen by me she had commenced to have pains over the right hip. The menopause occurred at 43 years of age. The vagina was senile and narrow, the cervix was hard, and a mass of growth could be felt in the canal and invaded the internal os. The fundus was small and the cervix moved freely. Per rectum, however, induration of the right sacro-uterine fold and broad ligament could be plainly made out. The appendages were small. On April 30th vaginal extirpation was performed in a nursing home. The patient rapidly recovered, returning to her house in 18 days. The uterus showed advanced glandular carcinoma commencing in the cervical canal and reaching as high as the internal os. The fundus was healthy. It is in this form that the growth frequently becomes very extensive