of tearing through the pia mater the blood was extravasated between that membrane and the brain substance ; it tracked round the optic chiasma and pons to the side of the medulla and thence poured by way of the iter into the ventricles.

The diagnosis of ruptured cerebral aneurysm was based on the sudden onset with vomiting and occipital headache, together with the remission of the symptoms. The blindness of the right eye appeared to be a disturbing factor in the diagnosis ; it was, presumably, explained post mortem by the pressure of the aneurysm on the right optic nerve. A specimen is preserved in the University College Museum in which the same condition is present. The alternative diagnosis was that of syphilitic meningitis and the improvement in the patient's general state, concurrent with antispasmodic treatment, lent for a time support to that belief. It is probable that the symptoms at the onset were due to a small rupture of the aneurysm ; this furnishes an explanation for the brown staining observed in the neighbourhood of the optic nerves. The aperture was plugged by blood clot and the effused blood was gradually absorbed with temporary relief to the patient's condition ; then, on the sixteenth day, the second rupture took place (perhaps due to straining at stool) and death resulted. Although a history was not obtained the etiology of the aneurysm was presumably syphilitic. This is the most plausible explanation for the atheromatous and degenerative state of the arterial vessels in so comparatively young a subject and many of the recorded cases of cerebral aneurysm have a syphilitic origin.

The case is of interest owing to the rare condition and the period of 16 days that elapsed between the primary rupture of the aneurysm and the fatal issue. With regard to the first point, the anterior cerebral artery holds the fourth place in the order of frequency of aneurysms of the cerebral vessels, the middle cerebral, the basilar and the internal carotid being more commonly affected. Beadles,¹ after an exhaustive review of 555 cases of cerebral aneurysm, states that "in aneurysms of the anterior cerebral and anterior communicating arteries the number in which definite cerebral symptoms occurred is so trifling and so vague that most certainly such an aneurysm is impossible of diagnosis." In this case, however, definite signs, the expression of a grave cerebral lesion, were present. The first symptom of the aneurysm was its partial rupture. This, of course, obtains in many of the cases recorded in the literature, the other types being (a) those in which no symptoms existed during life and the patient succumbed to some intercurrent disease ; and (b) those cases where the apoplectic seizure was preceded by pressure symptoms and signs simulating those given by a cerebral tumour.

As a rule, rupture of the aneurysm is followed by speedy death, and if the terminal event is delayed the interval between the primary and secondary rupture of the aneurysm is not of the duration noted here. In van der Byl's² case of rupture of the posterior cerebral the patient was taken to Middlesex Hospital, having fallen down in the street in a fit and having died while in the waiting room. Dr. J. Rose Bradford³ records a case of aneurysm of the basilar artery in which the patient lived for 40 hours from the onset of his symptoms, and he draws attention to the long survival between the first and second rupture. Optic neuritis and absent knee-jerks are no new features in the physical signs that a cerebral aneurysm may present.

In conclusion, I must express my thanks to Dr. Rose Bradford for his kindness in permitting me to publish the case. Brompton, S.W.

¹ Aneurysms of the Larger Cerebral Arteries, Brain, 1907, p. 285.
² Transactions of the Pathological Society of London, 1855-56. vol. vii., p. 129.
³ Aneurysm of the Basilar Artery, Transactions of the Clinical Society of London, vol. xxviii., 1895.

THE RESEARCH DEFENCE SOCIETY.—A public meeting in connexion with the Research Defence Society will be held in St. William's College, York, on Friday, Dec. 11th, at 3.30 P.M., when a lecture will be given by Mr. Stephen Paget on the work which has been done. The following amongst others have signified their intention of being present :—Lord Deramore, the Dean of York, Canon Watson, Canon Robinson, Rev. C. C. Bell, Mr. W. H. Jalland, Dr. F. Shann, Dr. R. Turner, Dr. Macdonald, Mr. E. R. Dodsworth, and Mr. H. V. Scott. The honorary secretary of the York branch is Mr. Gerald S. Hughes.

A NOTE ON THE BACTERIAL CONTAMINATION OF MILK AS ILLUSTRATING THE CONNEXION BETWEEN FLIES AND EPIDEMIC DIARRHŒA.

By J. T. C. NASH, M.D. EDIN., D.P.H. CANTAB.,
COUNTY MEDICAL OFFICER OF HEALTH OF NORFOLK.

MY views on this subject are well known and on some future occasion I shall revert to the whole matter. The object of the present communication is to record briefly the results of two simple experiments which I made in the summer of 1907.

I. The first was undertaken on a warm day in August, 1907, as follows. Two well-cleansed saucers were three-fourths filled with milk from the ordinary household supply. [This supply is subject to rather stringent conditions under guarantee.] One saucer was then covered over with a clean plate, the other was left exposed, both being placed on the same table in the kitchen. Some flies were observed in the room. Apart from the circumstance that one saucer was covered and the other uncovered, both were under identical conditions as to temperature, &c. Five hours later the uncovered milk was found to contain the bodies of two flies, and I proceeded to subject both milks to bacteriological tests, the results of which may be summarised as follows (both samples being kept covered, after the removal of the two flies which were in the sample which had been exposed for five hours) :—

Sample.	Number of bacteria per c.c. (as estimated from gelatin plate by naked-eye examination after 42 hours' incubation at 22° C. Dilution = 0·0001 c.c. original milk).	Taurocholate broth cultivation 0·0001 c.c. after 18 hours.	Glucose bile salt litmus broth 0·001 c.c. original milk.		Anaerobic milk culture for spores.
			Acid.	Gas.	
A (uncovered) containing two flies.	7,425,000 (after five hours' exposure).	Uniform turbidity.	+	+ $\frac{3}{4}$	-
B (covered over).	3,100,000 (after five hours covered over).	„	+	+ $\frac{1}{4}$	-

These results indicate that in five hours on a warm day milk exposed to flies contained millions more of gas-forming bacteria than the same milk kept covered over.

The following observations are significant of the physical deterioration of the exposed fly-containing milk as compared with the protected milk (both samples being kept covered, after the removal of the two flies which were in the sample which had been exposed for five hours) :—

Sample.	August 15th, after 24 hours.	August 16th, after 48 hours.	August 17th, after 72 hours.
A	Slight putrefactive odour.	Stinking odour more marked; milk clotted.	Odour still more marked; milk acid.
B	No odour.	Milk clotted; ordinary smell of sour milk.	No putrefactive odour; only that of lactic acid fermentation.

II. The second experiment was made in September, 1907. The temperature was 72° F. in the laboratory. Two portions of the same sample of milk hereafter referred to as C and D were placed in the same room in a cottage in which flies were present. Sample C stood exposed from 4 P.M. on the 11th until 8 A.M. on the 12th. Sample D was covered over but otherwise subject to exactly similar conditions. Both samples were subjected to bacteriological tests at 4 P.M. on the 12th. [With regard to the physical condition, the exposed milk—Sample C—contained two flies floating in it. It was observed that the ovipositors of these flies were completely extruded in each fly. The milk had a slight putrefactive odour and already there was a tendency to clot.

The covered milk—Sample D—looked and smelt quite fresh.]
The bacteriological condition was as follows:—

Sample.	Number of aerobic organisms capable of growth on gelatin at room temperature after 40 hours.	Growth in peptone broth at 37° C. 0·0001 c.c.	Glucose broth at 37° C. 0·0001 c.c.		Lactose broth at 37° C. 0·0001 c.c.		Agar slope at 37° C. 0·0001 c.c.	Litmus whey at 37° C.
			Acid.	Gas.	Acid.	Gas.		
C (exposed milk).	72,600,000	Growth + scum.	+	No note.	+	+	70 colonies.	+
D (covered milk).	24,640,000	Growth + scum.	+	No note.	++	+	7 colonies.	No note.

These results again prove a marked and rapid deterioration of milk exposed to flies, there being in the course of 24 hours no fewer than approximately 40,000,000 more bacteria per cubic centimetre in the exposed milk, and ten times as many organisms capable of growing on agar at the body temperature (37° C.). In round numbers the exposed milk showed 700,000 bacteria per cubic centimetre capable of growing at 37° C. against only 70,000 bacteria in the case of the milk protected from flies.

A further experiment showed that under anaerobic conditions the exposed milk (C) contained spores which on development in milk produced both gas and clot, while the covered milk (D) showed clot alone but no gas.

Remarks.—The importance of protecting milk from flies is illustrated by these experiments. It can readily be understood that the larger the number of flies the greater would be the pollution. It is not difficult to realise how in a fly-ridden district diarrhoeal diseases may be in excess unless care is taken to see that food-supplies (especially milk) are protected from the contaminating attentions of flies.

The experiments were undertaken on the lines of natural conditions obtaining in houses, so as to include the natural factors of symbiosis or antagonism of bacteria already existing in an ordinary milk-supply. A point I wish to make is that the number of putrefactive bacteria in milk is of great importance. The chemical changes produced in milk by, say, *thousands* of putrefactive bacteria may perhaps not be sufficient to cause serious diarrhoea, while the chemical changes produced by *millions* of such bacteria may conceivably be very prejudicial and excite acute diarrhoea.

A word as to treatment.—The facts indicate the wisdom of early expulsion of the bacteria and their toxic products by a large dose of castor oil which should, in my judgment, be followed by small doses—say, ten grains—of sulphate of soda every two hours dissolved in sterilised (boiled) water. No food should be given for six or eight hours, and then only *sterilised whey*, commencing with small quantities, gradually increasing same. The utmost care should be taken that all milk used for artificially fed infants should at all times be scrupulously protected from fly contamination, especially in dealing with an infant suffering from or convalescent from epidemic diarrhoea.

Norwich.

A CASE OF TRAUMATIC EXOPHTHALMOS.

By DAVID RANKEN, M.B., B.S. LOND. & DURH.,
M.R.C.S. ENG., L.R.C.P. LOND.,

HOUSE SURGEON TO THE SPECIAL DEPARTMENTS, ROYAL VICTORIA INFIRMARY, NEWCASTLE-ON-TYNE.

THE patient was brought to the Royal Victoria Infirmary, Newcastle-on-Tyne, on August 13th last, with the following history. He was driving a waggon whilst sitting on the shaft when the horse, which happened to be young and restive, lashed out suddenly with one of its hind legs, striking him in the face and knocking him over. His left eye was immediately dislocated forward beyond the lids, and directly afterwards he lost consciousness. About a quarter of an hour afterwards he came to himself and was sick, but felt no pain. The eye still remained protruding from the orbital cavity. Within the next hour he was brought to the hospital and whilst being examined again vomited.

On admission the patient was quite conscious and was able to answer questions intelligently. His pulse was good. There was epistaxis from both nostrils. On the left cheek there were several gashes, one extending into the ala of the nose, and obviously produced by the horse's hoof. The left eyeball protruded well beyond the lids. There was hyphæmia and the bulbar conjunctiva was torn all round in a neat circle quite close to the corneal margin. The sclera was intensely injected and the lids were swollen but not lacerated. No sight whatever remained in the eye. The left cheek was painful to the touch and one was unable to make out by palpation the normal outline of the left side of the face. Crepitus of the nasal bone and the nasal process of the superior maxilla was present. The patient had great pain and difficulty on attempting to open his mouth. The teeth outline was normal and there was no hæmorrhage either from the mouth or ears. Attempts to replace the eye were desisted from because of the pain they caused.

Under chloroform the eye could be easily replaced in the orbital cavity, but slight pressure on the lids was sufficient to luxate it forwards again. One was able to detect with the finger in the left nostril great comminution of the inner wall of the antrum of Highmore. The outline of the orbital plate of the superior maxilla on the face was lost. It was decided to remove the eye, and the recti muscles were divided in the ordinary way. On exposing the optic nerve it was found to be loose, and after dividing the rest of the structures connecting the eyeball with the orbital cavity one was able to draw out exactly two inches of nerve, still firmly connected with the eyeball, but with the cranial end severed as if with a sharp knife. Very little bleeding ensued. The wounds in the cheek were sutured with horse-hair, a sponge was left over the left eyelids, and the nostrils were loosely packed with gauze wrung out of a perchloride of mercury solution.

Recovery was uninterrupted. There was some serous oozing from the eye and nostrils during the next few days. Ten days later the patient left the hospital. He was seen eight weeks after the operation when he stated that he had been working then for some weeks past. The wounds in his cheek were quite healed, his left orbital cavity presented the appearance as after an ordinary excision, and he could open his mouth quite wide.

Fracture of the anterior fossa of the skull with severance of the optic nerve is not a common accident, but when complicated with complete exophthalmos becomes one of extreme rarity. The average length of the optic nerve is 5 centimetres, or about two inches, and hence it must have been severed in this case just at its origin from the optic chiasma, in all probability when passing through the optic foramen. The fracture of the superior maxilla on that side was an additional complication.

This case was under the care of Mr. J. D. Wardale, ophthalmic surgeon to the hospital, and I am indebted to him for permission to publish these notes.

Newcastle-on-Tyne.

EXTENSION OF A PLEURAL EFFUSION TOWARDS THE SOUND SIDE IN A STILL-BORN CHILD

CORRESPONDING TO GROCCO'S PARAVERTEBRAL TRIANGLE.

By ROBERT D. KEITH, M.A., M.D. ABERD.,
LECTURER ON PHYSIOLOGY, THE STRAITS MEDICAL SCHOOL, SINGAPORE;

AND

ARTHUR KEITH, M.D. ABERD., F.R.C.S. ENG.,
CONSERVATOR OF MUSEUM, ROYAL COLLEGE OF SURGEONS OF ENGLAND

THE subject of this note was a still-born child, probably of Chinese parentage, found in the streets of Singapore and investigated by us with a view of discovering the cause of the anasarca condition of its tissues. In this we were not successful; the ductus venosus which was found absent in a similar case was here present; the heart also was normally formed; the tissues of the body, especially the kidneys, were extensively invaded with leucocytes.

Our main reason for recording the case is that it supports those who regard Grocco's paravertebral triangle of dulness as due to a direct extension of the pleural fluid or contents from the diseased towards the sound side. A reference to