of the six wine cases and two of the five tea cases occurred in the autumn. Cases of lead poisoning traced to tea packed in lead occurred at Odessa in 1886, and the safety of so packing it has often been called in question. It is acknowledged by the trade that many teas contain a trace of lead. The tea may apparently get its lead in several different ways. It may no doubt mechanically rub off lead from the lining of the chest, or still more (from the larger proportion of surface exposed) from the lead paper of the small packets. It would be interesting to make comparative analyses of the tea from the corners and sides of the chests and from the middle. The lead lining of the chest is sometimes acted upon either by the tea fermenting inside it or from the chest being made of green wood. The salt produced is basic lead carbonate, an impalpable powder which would readily diffuse itself through the tea. A slow action by which the lead becomes brittle and "perished" goes on in shops and places where the cases are kept for a long time—possibly tea kept long in such coverings might be contaminated in this way. Traces of copper, chips of lead paper, and in one case a piece of brass, found by the analyst in several small packets of tea, would seem to point to great carelessness in packing either in England or abroad. Mr. Angell of Southampton, the county analyst for Hants, to whom I have submitted most of the teas I have had examined, finds by experiment that tea infusion has a powerful solvent action on lead, so that the chips of lead paper would be a source of danger in the teapot. The samples examined were nearly all packet teas of the common sorts. They were packed in ordinary paper, in lead-paper, and in lead paper lined with white paper. Mr. Angell has kindly given me the results of some analyses he made on his own account, so that altogether twenty-two different kinds of tea have been examined, of which seventeen contained traces of lead. The quantity of lead found was in most cases extremely small—of sixteen samples of eleven different brands of tea the lowest quantity found was 1.375 milligrammes per pound, the highest was 18 mgr. per pound, and the average of all 5.962 mgr. per pound. The lead was most unevenly distributed; of two packets of the same brand bought at the same time at the same shop one contained 1.6 mgr. per pound and the other 18 mgr. How far one would expect these small quantities of lead to prove dangerous it is difficult to judge. Data as to quantities which have proved poisonous are few and refer almost exclusively to water. Oliver refers in his Goulstonian Lectures to a case of his where only $\frac{3}{16}$ gr. per gallon was found. Parkes quotes cases of lead poisoning due to water containing only $\frac{1}{16}$ gr. per gallon, and sums up by saying that water with $\frac{1}{16}$ gr. per gallon should be considered dangerous and that some persons may be affected by less. It is of course impossible, either in water or in tea, to say that a found quantity is in any degree constant; still greater uncertainty is introduced by the fact that persons differ very widely in the amount of water that they drink. The quantity of tea is perhaps not quite so variable; a quarter of a pound a week seems a fair average. If one assumes two pints of water to be the average quantity drunk per diem one can make a sort of comparison which may be given for what it is worth. Tea containing 6 mgr. of lead per pound (the average of all the estimations) would be then about equivalent to water containing $\frac{7}{17}$ gr. per gallon; while the tea which had most lead (18 mgr. per lb.) would be equal to water containing $\frac{3}{8}$ gr. per gallons. Possibly these quantities may be peculiar to the year owing to some temporary cause; but, assuming that they are not exceptional, though far too small to cause a widespread epidemic, such tea would probably cause plumbism in a few susceptible individuals and would certainly add to the danger of their being affected from other sources, such as water or home-made wine. Moreover, may not some of the many cases where tea is suspected of causing vague dyspeptic and colicky symptoms be in reality minor cases of lead poisoning? The tea trade is in the hands of the Customs authorities; are they justified in passing tea which contains any lead whatever?

Weyhill, Andover.

PHOTOGRAPHIC CLUB.—As a large number of the medical profession are interested in photo-micrography, we are asked to state that Mr. Andrew Pringle, F.R.M.S., will read a paper on this subject at the Photographic Club, Anderton's Hotel, Fleet-street, on Wednesday, Oct. 3rd. Free tickets of admission may be obtained by sending stamped envelope to the honorary secretary, Mr. James A. Sinclair, 26, Charing-cross-road, W.C.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

"BONNET-PIN IN THE MALE URETHRA."

BY J. J. STACK, L.R.C.P. EDIN., L.R.C.S. IREL.

A FEW years ago, a man, aged about forty-five years, called on me and asked me to remove a pin from his urethra. He said that he passed the beaded head of a bonnet-pin into his "pipe" "to relieve himself from the gravel." He was in a fearful state of mind over the accident, as he was afraid that the pin had slipped into his bladder. On examination I found the beaded end by pressing my fingers well up behind the scrotum, where I could use pressure enough to propel the foreign body forwards. I got an ordinary No. 8 catheter, and cut off the top with the scissors. This I passed into the urethra, with the object, if possible, of ensheathing the point of the pin. My idea, however, failed, and after several trials I was obliged to give it up. I then pressed the beaded end forwards as well as I could, guiding the point with a small forceps, until I got the point within a short distance of the meatus. I next forced the point through the tissues and got the body of the pin exposed. This I pulled on until the head came to the spot where the point was. By pressing the body of the pin downwards and backwards—the bead was in the meatus—and with very little manipulation, I was enabled to thus extract it. This is, so far, exactly like Dr. J. H. Simpson's case, but the presence in the urethra was less mysteriously accounted for. There was a little soreness in the penis for a day or so, with a slight discharge, but beyond this the case progressed favourably. Dr. Simpson's case and mine are very curious coincidences, and one description would nearly suit the two cases.

New Cross-road, S.E.

CURIOUS DEFORMITY OF THE CHEST WALL.

BY S. FLOYD, M.B., L.R.C.P. LOND., M.R.C.S. ENG.

THE following deformity of the chest may be of sufficient rarity to be worth recording. A boy aged thirteen was admitted to the hospital ships, Long Reach, suffering from a mild attack of small-pox. On physical examination the second, third, and fourth ribs and costal cartilages on the right side of the chest were found to be deficient, so that the space wanting in the chest wall measured three and a half inches vertically and two and a half inches horizontally. At the fourth space the right side of the thorax measured twelve and a half inches in circumference, and the left fourteen inches. The unprotected space was markedly depressed on deep inspiration, but the ordinary physical signs elicited upon examination of the thorax were normal, though a little pronounced. The pectoralis major and minor were absent on the right side, with the exception of the clavicular head of the former, which was more developed than on the opposite side. The rest of the pectoralis major was represented by a fibrous band. The right deltoid muscle was hypertrophied, but the boy had almost as much power with the right as with the left arm. In every other respect the chest was quite normal. The costal ends of the third and fourth cartilages could be felt to be joined together, but the other deficient ribs and cartilages terminated separately. The deformity is congenital, and no other member of the family is affected in any way.

Hospital Ships, Long Reach, near Dartford.

A CASE OF HÆMOPHILIA.

BY JAS. WM. WHITE, M.B., C.M. GLASG.

THE patient, a boy aged five years, has since birth been an object of interest and anxiety to his friends. The most moderate blow, and at times even a hand pressure, have sufficed to cause ecchymoses which are slow to disappear. The limbs, therefore, have almost normally a mottled appear-

ance. The articular phase is marked, the knee-joints having several times suddenly filled with blood after trifling injuries. Nevertheless the patient is ruddy, full of life, and enjoys perfect health. Under such circumstances he lately had the two left lower (temporary) molars removed for caries, the operation being performed under nitrous oxide gas. There remained after extraction a slight laceration on the inner aspect of the anterior socket. At the time of operation hæmorrhage was trifling, and the boy returned home well. After a few hours, however, blood began to well from both sockets, and the flow was with the greatest difficulty arrested by graduated compresses. Several times within the next three days hæmorrhage recurred, and the same trouble of arresting it was encountered. Local styptics were of no assistance to local pressure, but ergot seemed to act somewhat beneficially. Matters now appearing to be well, the patient was put to bed; but from the laceration before mentioned blood streamed, and the pillow was soon quite saturated. The hæmorrhage only ceased when fainting occurred, and even afterwards, whenever pressure was relaxed for a brief interval, blood flowed freely. This state of affairs lasted for five or six days, the outbursts being gradually more controllable, and finally bleeding ceased. The gum was atrophied and flattened by the pressure. The case demonstrates the risk of extraction under such circumstances, for in the present instance the fangs were short, being in part absorbed, and bleeding under normal circumstances should have been slight. Hereditary transmission followed the ordinary course.

Regent-park-square, Glasgow.

A Mirror

OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et mor-
torum et dissectionum historias, tum aliorum tum proprias collectas
habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*,
lib. iv. Proœmium.

ROYAL HOSPITAL FOR CHILDREN AND WOMEN, WATERLOO-ROAD, S.E.

TUMOUR OF THE VENTER OF THE SCAPULA INFILTRATING
THE SUBSCAPULARIS MUSCLE IN A BOY AGED TEN;
GROWTH OF THE TUMOUR INTO THE THORACIC
PARIETES; ATTEMPTED REMOVAL; PNEUMO-
THORAX; DEATH; NECROPSY; REMARKS.

(Under the care of Mr. A. MARMADUKE SHEILD.)

THE operative treatment of growths of, or in, the region of, the scapula, of a sarcomatous character, is not very promising, even when the whole of the tumour has apparently been successfully removed. There is a great tendency in such growths to spread into parts around, and so the success of the operation is too frequently more apparent than real. Operation should be performed early. This case presents many interesting features, its position and extent rendering exact diagnosis before removal quite impossible. For the notes of this case we are indebted to Mr. Huntly, house surgeon.

The patient, a boy aged ten, was sent to Mr. Sheild by Mr. Muirhead Little, having been taken to the Orthopædic Hospital on account of atrophy of the muscles about the right shoulder and marked weakness and wasting of the upper extremity. In the axilla there was a hard, rounded, fixed mass the size of an orange, which pushed up the shoulder and obviously compressed the axillary nerves. It was exceedingly difficult to make out whether or not the tumour moved with the scapula, but it appeared to do so slightly. The skin over the tumour was not adherent and was not inflamed. There was no tenderness except on firm manipulation. The measurements of the tumour, taken with the arm abducted as far as it would go, were—(1) from above downwards, two inches and a half, and (2) from before backwards, 2 inches. Looked at from the front there appeared to be marked wasting of the pectoral muscles on the right, there being quite a depression in the subclavicular region as compared with the other side. The deltoid also was much wasted, the head of the right humerus and acromion being

much more prominent. The shoulder was kept from three-quarters of an inch to one inch higher on the right side than on the left, and, as noticed by the mother, the boy seemed to keep his head a little bent towards the right shoulder. The lower part of the anterior margin of the trapezius, where it was forming the posterior border of the posterior triangle of the neck, stood out more prominently on this side also. With the arms held to the side the tumour was only slightly visible, but when the right arm was abducted it was very evident. From behind, the right supra- and infra-spinatus muscles were considerably atrophied, and the tumour was perfectly evident with the arm hanging by the side, forming quite a bulge along the axillary border of the scapula. There was well-marked lateral curvature of the spine in the dorsal region, with the convexity towards the left side, so that the left inferior angle of the scapula stood out prominently. On passing a tape underneath the axillæ and taking the measure of the circumference of the shoulders separately, it was found that on the right side it was about 11½ in. and on the left side 10½ in., the tape passing over the clavicles just internally to the acromion on each side. The right arm, as might be expected, was more wasted than the left, the circumference of the former over the fleshy part of the biceps (six inches from the tip of the acromion) being 5½ in., whereas on the left side it was 6¼ in. The circumference of the forearm one inch below the internal condyle on the right side was 6 in. and on the left side 6¾ in.

Movements of the right shoulder-joint.—On the right side, when asked to abduct his arm (holding the body fixed), it was found that it was only possible to do this to a distance of half a right angle. It was impossible for him to raise his arm higher without causing pain. He could swing his arms backwards and forwards about equally. Rotation was much affected on the right side, more especially external rotation, which was very limited. Internal rotation was better, but not so good as on the left side. Circumduction was impossible. He was able to place his right hand round over the left shoulder almost as far as he could place the left arm over the right shoulder. There was no pain experienced in making any of these movements. Those at the right elbow and the wrist were as good as on the left side. Sensation was everywhere unimpaired.

Family history.—The father is living and healthy. The mother is also a fairly healthy woman. There were seven children in the family, three of them are living. Of the four who are dead one died from tumour on the brain, one from "water on the brain," one was born dead, and in one death was indefinite, but it was not a full-term child; it only lived for one week. There was no history of tubercle to be obtained in any branch of the family.

History of the disease.—About twelve or fifteen months ago the patient was taken to the Royal Free Hospital by his mother, who thought that he had put his right shoulder out. On being asked why she thought so, she said because a small lump was forming in his right armpit. She also said that she was told that the patient was suffering from curvature of the spine and was treated for this. She likewise noticed that the patient always kept his head flexed towards the right shoulder, and held the latter raised towards the head. Up to four or five months previously the patient had fairly good movements about his right shoulder-joint, but thought that he had had a greater tendency to use the left arm than the right since the lump in the armpit had been getting larger. There is no marked history of pain. For the last four or five months the patient had been noticed to keep the right elbow flexed. When not in this position the elbow was extended and the whole arm was held in such a position that the palm of the hand was turned outwards. The patient had been noticed to be often rubbing the arm with his left hand in the region of the elbow-joint, but whether from pain or a feeling of coldness was not quite clear. There had never been starting pains in the arm, nor had it ever been noticed to be blue or swollen. He was said to always join in games with the other boys at school, but he used the left arm whenever he had the opportunity.

The parents of the boy, having had the grave and difficult nature of the case explained to them, wished an operation to be performed, and this was accordingly undertaken on Aug. 13th. The A.C.E. mixture was administered, and proceedings were commenced by making an incision into the axilla, to try more accurately to define the limits of the tumour. It was now quite evident that the tumour was of scapular origin, but the finger could not pass between the