The Effect of Operation and also of Hemorrhage upon the Pregnant Patient.—NEU (*Arch. f. Gynäk.*, 1906, Band lxxx, Heft 2) reports the following cases: Case I, a primipara, between the fifth and sixth months of pregnancy, had repeated bleeding from varicose veins at the entrance to the vagina. Although her condition seemed otherwise good and there was no essential difficulty in the birth, which was spontaneous, the child was stillborn, and no apparent cause could be assigned for this except the repeated hemorrhage. In Case II a pregnant patient was found to have multiple cysts with nodules in the vagina. Excision was practised, followed by free bleeding, which required the use of a tampon to control it. This patient gave birth to a macerated foetus. In Case III the vagina was almost occluded by the scars from a former labor. Operation was performed, and was complicated by hemorrhage. In this case the pregnancy remained without interruption. In Case IV, a multipara between five and six months pregnant had empyema of the gall-bladder, with peritonitis. Incision and drainage were practised and were followed by recovery. Spontaneous labor afterward occurred, after which the gall-bladder was removed and adhesions of the intestines liberated. This patient made a good recovery.

Innervation of the Uterus.—ROITH (*Monats. f. Geburts. u. Gynäk.*, 1907, Band xxv, Heft 1) has studied the nervous supply of the uterus. He found upon minute dissection that the uterine muscle is stimulated by nerve fibers which arise from local ganglia in the connective tissue of the pelvis and also of the cervix.

Placenta Prævia.—BURGER-GRAF (*Monats. f. Geburts. u. Gynäk.*, 1907, Band xxv, Heft 1) contributes an interesting paper upon this subject. In 44,676 births, 344 cases of placenta prævia were observed, being 77 per cent., or 1 in 130. The reports of other clinics give a frequency of from 1 in 58 to 1 in 1564. As regards age, placenta prævia is most frequent between the years of thirty-one and thirty-five. It is very much more common in multipara, 88.3 per cent. of these patients being multiparæ, and 11.7 per cent. primiparæ; 16 per cent. of the multiparæ were in the second pregnancy. Central placenta prævia is most frequent in multiparæ, and the tendency to this complication increases with repeated births. This variety is least frequent in primiparæ. In central placenta prævia only 31.3 per cent. go to full term; 16 per cent. were delivered before viability, and 52.6 per cent. had premature labor. In lateral placenta prævia the patient goes to term most frequently. Next to this comes partial placenta prævia, while central placenta prævia naturally interrupts pregnancy most often. As to the time in gestation when central placenta prævia develops, it is most frequent in the eighth, ninth, and tenth months. Partial placenta prævia is most common in the ninth month, and lateral placenta prævia in the tenth month.

There seems to be some relation between placenta prævia and abnormality of the pelvis, as in 13.7 per cent. of these cases the pelvis was not normal. Twin pregnancy also predisposes to placenta prævia, and eight cases of twins occurred among the patients reported. As regards the presentation of the foetus, the vertex presented in 69.2 per cent., transverse and oblique presentations occurred in 21.1 per cent., and