the rupture of a pus-tube or ovary. If recovery takes place, the parametritis and para-typhlitic exudates can usually be diagnosticated by their characteristic shape and position.

Diseased adnexa are generally recognized from their close relation to the uterus. When these are associated with appendicitis the diagnosis may be exceedingly difficult.

Alexander's Operation.—SIMÕES (*La Gynécologie*, 1897, Nos. 3 and 4) concludes an elaborate paper on this subject with the following summary: The operation should only be performed in cases of movable uterus. Since metritis and prolapsus usually accompany the retro-deviation, curettement, amputation of the cervix, and colpo-perineorrhaphy should supplement Alexander's operation in cases of chronic retroflexion. Extra-peritoneal shortening of the round ligaments is preferable to all the other methods of supporting the uterus in its normal position, as it is the most simple and rational. Finally, statistics have shown that pregnancy is least likely to be interfered with after this operation.

PÆDIATRICS.

UNDER THE CHARGE OF

LOUIS STARR, M.D.,
OF PHILADELPHIA.

ASSISTED BY

THOMPSON S. WESTCOTT, M.D.,
OF PHILADELPHIA.

The Clinical Variability of Rubella—PÉRON (*Revue mensuelle des Maladies de l'Enfance*, October, 1897) records three cases of rubella in a family, one of which, in the person of the father, aged forty-five years, presented almost a typical picture of severe measles.

The first patient affected was a son, aged eleven years, who, after three days of mild prodromes, presented a rapidly spreading typical rubellous eruption, with enlargement of the post-cervical lymphatic chains, slight fever, secondary sore-throat, slight hoarseness, coryza, and infrequent cough. The attack terminated on the third day.

Fourteen days after the appearance of the son's eruption the father was taken ill, with violent headache, lassitude, and chilly sensations. Then followed violent attacks of sneezing, hoarseness, and cough. The latter became very frequent and distressing, and disturbed sleep as markedly as in the prodromal stage of measles. Toward evening of the fourth day the symptoms increased, the patient took to bed, passed a bad night, and awoke with a confluent and generalized eruption. There was slight cervical adenopathy. With the appearance of the eruption there were abundant coryza, lachrymation, and conjunctival injection. Upon the uvula was seen a typical rubellous mottling.